7.1 Urinary Tract Infection

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "It burns when I pass urine and I go very often." **Objective** • Urine cloudy and smells bad • Dipstick positive for nitrites and leukocytes • Tender above the pubis	**ACTUAL** **Impaired urinary elimination** related to bladder infection as evidenced by dysuria, frequency and urgency.	Within 48–72 hours of treatment, says the burning and frequency have gone.	• Collect a midstream urine sample before the antibiotics, as ordered. • Give antibiotics as prescribed and stress finishing the course. • Unless fluids are restricted, encourage 2–3 L/day. • She should pass urine often and empty her bladder fully each time.	Symptoms gone: **goal met**.
Subjective • Gives suprapubic pain 5/10 **Objective** • Tender above the pubis • Uncomfortable when passing urine	**ACTUAL** **Acute pain** related to bladder mucosal inflammation as evidenced by suprapubic pain and burning.	Pain 3/10 or less within 24 hours.	• Give analgesia or a urinary analgesic as prescribed. • If she finds it soothing, put warmth over the lower abdomen.	Pain controlled: **goal met**.
Subjective • "I get this infection every few months." **Objective** • Keeps getting • UTIs • Female, sexually active	**RISK** **Risk for infection (recurrent)** as evidenced by risk factors: history of recurrent UTI.	Can describe how to prevent another infection.	• Teach her to wipe from front to back. • Teach her to pass urine after intercourse. • Encourage plenty of fluids and not holding on to urine.	Describes the measures: **goal met**.