7.13 Urinary Incontinence

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I leak urine when I cough, sneeze or laugh." **Objective** • Stress incontinence • Has had several children • BMI 31 kg/m²	**ACTUAL** **Stress urinary incontinence** related to weak pelvic floor muscles as evidenced by leakage with increased abdominal pressure.	Within 6–12 weeks, leaks less often.	• Teach pelvic floor exercises and check she is doing them right. • She should practise every day and keep going long term. • Advise weight loss, and treating constipation and any chronic cough. • Ask her to keep a bladder diary. • If there is no improvement, refer her to a continence specialist.	Fewer episodes: **goal met**.
Subjective • "I stopped going to family visits because of the smell." **Objective** • Avoids going out • Uses a lot of pads	**ACTUAL** **Social isolation** related to fear of leakage and odour as evidenced by avoiding social activities.	Gets back to at least one social activity.	• Talk about which continence products would suit her. • Plan a toilet visit before she goes out. • Let her talk about how she feels.	Back to an activity: **goal met**.
Subjective • *No specific complaint* **Objective** • Skin always damp • Wears pads	**RISK** **Risk for impaired skin integrity** as evidenced by risk factors: moisture from urine.	Perineal skin stays intact.	• Teach her to wash with a mild cleanser and dry the skin after each episode. • Use barrier cream. • Change pads quickly.	Skin intact: **goal met**.