7.14 Urinary Retention

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I need to pass urine but nothing comes." **Objective** • Bladder distended and dull to percussion • Bladder scan 700 mL • Taking an anticholinergic drug	**ACTUAL** **Urinary retention** related to bladder outlet obstruction or medication effect as evidenced by distended bladder and inability to void.	Within 24 hours the bladder is emptied and he passes urine on his own.	• Try things that help him go: privacy, running water, standing or sitting up. • Catheterise as ordered and record how much drains. • Go through anticholinergic and opioid drugs with the physician. • Do a trial without catheter as ordered and scan for residual.	Voids on his own with a small residual: **goal met**.
Subjective • Gives lower abdominal discomfort 7/10 **Objective** • Restless • Sweaty	**ACTUAL** **Acute pain** related to bladder distension as evidenced by discomfort and restlessness.	Says he feels relief once drained.	• Drain the bladder as ordered. • Check pain again after drainage. • After a large volume drains, watch for blood in the urine.	Relieved: **goal met**.
Subjective • *No specific complaint* **Objective** • Catheterised • Urine left in the bladder	**RISK** **Risk for infection** as evidenced by risk factors: urinary catheter and urinary stasis.	No UTI.	• Aseptic technique for insertion. • Keep the drainage system closed. • Take the catheter out as early as possible.	No infection: **goal met**.