8.13 Diabetes Insipidus

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I drink and pass urine all the time." **Objective** • Passing 6 L of urine a day • Urine specific gravity low • Na^+ 150 mmol/L	**ACTUAL** **Deficient fluid volume** related to excessive dilute urine output as evidenced by polyuria and hypernatremia.	I&O balanced and sodium normal.	• Record urine output and specific gravity every hour. • Give desmopressin as prescribed. • Replace fluids as ordered. • Weigh daily.	Balanced: **goal met**.
Subjective • "I cannot sleep because I go to the toilet." **Objective** • Up at night to pass urine • Tired in the day	**ACTUAL** **Disturbed sleep pattern** related to nocturia as evidenced by waking to void.	Within 1 week of treatment, says he sleeps better.	• Give desmopressin at the prescribed time (often at night). • Keep a urinal by the bed. • Only cut down evening fluids if he is advised to.	Sleeping better: **goal met**.
Subjective • *No specific complaint* **Objective** • On desmopressin • Headache	**RISK** **Risk for electrolyte imbalance** as evidenced by risk factors: desmopressin causing water retention.	Serum sodium stays within the normal range.	• Follow sodium. • Teach him to drink only when thirsty. • Tell him to report headache, nausea or confusion.	Normal: **goal met**.