8.5 Hypoglycemia

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel shaky, sweaty and hungry." **Objective** • Glucose 52 mg/dL • Shaking and sweating • Cannot concentrate • HR 110/min	**ACTUAL** **Acute confusion** related to low blood glucose as evidenced by glucose 52 mg/dL, tremor, sweating and difficulty concentrating.	Glucose above 70 mg/dL within 15 minutes.	• If he is conscious: give 15–20 g of fast-acting glucose, recheck in 15 minutes and repeat if needed. • After that, give a carbohydrate snack or his next meal. • If he is unconscious: IV dextrose or IM glucagon as prescribed, and nothing by mouth. • Work out why it happened.	Glucose normal: **goal met**.
Subjective • "I did not eat lunch but took my insulin." **Objective** • Skipped a meal • Cannot say how to prevent a hypo	**ACTUAL** **Deficient knowledge** related to matching food and insulin as evidenced by taking insulin without eating.	Can explain how to prevent hypoglycemia.	• Teach him to time his meals with his insulin. • Tell him to carry glucose with him.	Explains prevention: **goal met**.
Subjective • *No specific complaint* **Objective** • Has hypos again and again • Drives to work	**RISK** **Risk for injury** as evidenced by risk factors: hypoglycemic episodes with confusion.	No injury from hypoglycemia.	• He should check his glucose before he drives. • Teach the family to spot and treat a hypo, including using glucagon if it is prescribed. • He should wear diabetes identification.	No injury: **goal met**.