9.5 Hip Fracture and Internal Fixation

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My hip hurts after I fell." • Gives the pain 8/10 **Objective** • Leg shortened and turned outwards • Age 80 years • Day 1 after fixation	**ACTUAL** **Acute pain** related to fracture and surgery as evidenced by pain score 8/10.	Pain 3/10 or less and joins in with mobilisation.	• Give analgesia regularly; consider a nerve block as ordered. • Put a pillow between her legs as ordered. • Give analgesia before physiotherapy.	Pain controlled: **goal met**.
Subjective • Confused at night **Objective** • Newly disoriented • Elderly, on opioids	**ACTUAL** **Acute confusion** related to surgery, pain and medication as evidenced by new disorientation.	Back to her usual orientation.	• Screen for delirium every day. • Reorient her and bring the family in. • Treat pain, dehydration and constipation.	Oriented: **goal met**.
Subjective • *No specific complaint* **Objective** • Elderly • Immobile • Had hip surgery	**RISK** **Risk for adult pressure injury** as evidenced by risk factors: older age, immobility and surgery.	Skin stays intact.	• Turn every 2 hours, as far as allowed. • Nurse her on a pressure-relieving mattress. • Check heels and sacrum every shift. • Get her up early, as ordered.	Skin intact: **goal met**.