9.8 Amputation

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I can still feel my foot even though it is gone." **Objective** • Below-knee amputation • Phantom limb pain	**ACTUAL** **Chronic pain** related to phantom limb sensation as evidenced by pain in the absent limb.	Says the phantom pain is less.	• Tell him phantom pain is real. • Give analgesia and neuropathic agents as prescribed. • Encourage massaging the stump and desensitisation.	Less pain: **goal met**.
Subjective • "I am not a whole person now." **Objective** • Will not look at the stump • Tearful	**ACTUAL** **Disturbed body image** related to loss of limb as evidenced by avoidance and distress.	Looks at the stump and talks about rehabilitation.	• Let him grieve and talk about it. • Bring him into the care of his stump. • Arrange for him to meet another amputee or a prosthetist.	Engages with rehabilitation: **goal met**.
Subjective • *No specific complaint* **Objective** • Stump	**RISK** **Risk for disuse syndrome** as evidenced by risk factors: amputation, pain and prolonged flexed positioning of the stump.	Stump keeps its full extension.	• After the first day, no pillows under the stump, as ordered. • If allowed, lie prone for 20–30 minutes several times a day. • Do the exercises as taught.	No contracture: **goal met**.