13.5 Prostate Cancer and Prostatectomy*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I leak urine since the catheter was removed." **Objective** • Dribbling after catheter removal • Uses three pads a day	**ACTUAL** Stress urinary incontinence related to sphincter weakness after prostatectomy as evidenced by urine leakage.	• Using fewer pads within 6–12 weeks.	• Teach pelvic floor exercises and check his technique. • Supply suitable continence products. • Suggest cutting down on caffeine and alcohol. • Explain that continence usually improves over several months.	• Met as continence improves.
Subjective • "Will I be able to have relations with my wife?" **Objective** • Worried about erectile function after surgery • Avoids the subject with his wife	**ACTUAL** Impaired sexual function related to effects of radical prostatectomy as evidenced by expressed concern about erectile function.	• Discusses his concerns and the options available.	• Offer privacy and keep a non-judgmental manner. • Give accurate information about recovery and treatment. • Include his wife if he wishes. • Refer to urology or sexual health services.	• Achieved once options have been discussed.
Subjective *No specific complaint* **Objective** • Pelvic surgery • Malignancy	**RISK** Risk for thrombosis as evidenced by risk factors: pelvic surgery and malignancy.	• No DVT.	• Give prophylactic anticoagulation as ordered. • Mobilize early. • Teach leg exercises and how to check his calves.	• Outcome reached if no DVT develops.

*References for this care plan: [5], [32], [35], [37]. Full entries are listed in tFull entries are listed in the References section of the CPMN book.