17.1 Bronchiolitis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Mother says, "He is breathing fast and not feeding." **Objective** • 4 months old; RR 64/min • Nasal flaring, subcostal recession • Wheeze and crackles; SpO_2 90%	**ACTUAL** Ineffective airway clearance related to excessive mucus and small-airway inflammation as evidenced by retractions, crackles and low SpO_2.	• SpO_2 of 92% or more (or the local target) and less work of breathing within 48 hours.	• Clear the nose with saline drops and gentle suction before feeds. • Give oxygen to the target SpO_2. • Keep his head slightly raised. • Watch RR, SpO_2 and for signs of exhaustion or apnea.	• Considered met when SpO_2 is on target and distress has eased.
Subjective • Mother says, "He takes only a little milk." **Objective** • Taking under half his usual feeds • Fewer wet diapers	**ACTUAL** Ineffective infant feeding dynamics related to dyspnea during feeds as evidenced by reduced intake and fewer wet diapers.	• Stays well hydrated.	• Offer small, frequent feeds. • Give NG feeds or IV fluids as ordered if intake falls short. • Count wet diapers and weigh daily.	• Goal reached while he is hydrated.
Subjective *No specific complaint* **Objective** • Young infant • Respiratory infection	**RISK** Risk for infection [transmission to others] as evidenced by risk factors: contagious viral infection.	• No spread to other babies.	• Use contact and droplet precautions per policy. • Insist on strict hand hygiene. • Teach the parents hand hygiene.	• Met if there is no cross-infection.

*References for this care plan: [35], [38], [113], [117]. Full entries are listed in the References section of the CPMN book.