17.22 Hypospadias Repair*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Crying after surgery **Objective** • Penile dressing • Urinary stent in place • Crying	**ACTUAL** Acute pain related to surgery as evidenced by crying.	• Looks comfortable within 1 hour of analgesia.	• Give weight-based analgesia as prescribed. • Give the prescribed antispasmodic for bladder spasm. • Comfort and distract him.	• Achieved when he is comfortable.
Subjective • Mother asks, "How do I care for the tube at home?" **Objective** • Stent in place • Mother unsure how to put the diaper on	**ACTUAL** Inadequate health knowledge [parental] related to stent and dressing care as evidenced by questioning.	• Parent demonstrates the care plan.	• Teach double-diapering with the stent left free. • Teach the signs of infection.	• Outcome reached when demonstrated.
Subjective *No specific complaint* **Objective** • Surgical site • Urinary stent	**RISK** Risk for infection as evidenced by risk factors: surgical wound and urinary stent.	• Surgical site free of infection.	• Keep the area clean. • Give antibiotics as ordered. • Teach the parents the signs of infection.	• Considered met if no infection develops.

* **References for this care plan:** [35], [38], [49]. Full entries are listed in the References section of the CPMN book.