2.14 Blood Transfusion*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I get tired after walking a few steps." **Objective** • Hb 6.8 g/dL • Pale • HR 110/min on exertion	**ACTUAL** Decreased activity tolerance related to reduced oxygen-carrying capacity as evidenced by exertional fatigue and low hemoglobin.	• After transfusion, manages self-care without breathlessness.	• Space activities with rest. • Give blood as prescribed. • Observe HR and SpO_2 during activity. • Repeat Hb after the transfusion, as ordered.	• Achieved when ADL are completed without fatigue.
Subjective • "I am worried about receiving blood." **Objective** • Tense facial expression • Repeatedly asks whether it is safe	**ACTUAL** Excessive anxiety related to an unfamiliar procedure as evidenced by expressed worry and repeated questioning.	• Before the transfusion starts, she reports understanding and less worry.	• Explain the procedure, its length and the safety checks. • Tell her which symptoms to report. • Stay with her for the first 15 minutes.	• Outcome reached if she is calm and cooperative.
Subjective *No specific complaint* **Objective** • Receiving a blood component • Has had transfusions before	**RISK** Risk for physical injury as evidenced by risk factors: potential transfusion reaction or ABO incompatibility.	• Transfusion completed without a reaction.	• Verify identity and the unit at the bedside with a second nurse. • Record baseline vital signs. • Start slowly and observe closely for the first 15 minutes. • Monitor vital signs per protocol. • In a reaction: stop the unit, keep the line open with saline, call the physician and return the unit. • Complete within 4 hours of removal from storage.	• Considered met if no reaction occurs.

* **References for this care plan:** [35], [37], [70], [114], [130]. Full entries are listed in the References section of the CPMN book.