2.20 Urinary Catheter Care*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I cannot pass urine." **Objective** • Distended bladder • Bladder scan 600 mL	**ACTUAL** Impaired urinary elimination related to urinary retention as evidenced by inability to void and bladder distension.	• Bladder drained; urine output at least 0.5 mL/kg/hour.	• Insert the catheter aseptically, as ordered. • Measure urine output. • Keep the bag below bladder level. • Plan early removal.	• Goal reached when drainage is adequate.
Subjective • "The catheter feels uncomfortable." **Objective** • Restless • Tugging at the catheter	**ACTUAL** Impaired physical comfort related to an indwelling catheter as evidenced by reported discomfort.	• Discomfort reduced within 24 hours.	• Secure the catheter to the thigh. • Check for kinks and traction. • Give perineal care.	• Met when he reports comfort.
Subjective *No specific complaint* **Objective** • Indwelling urinary catheter	**RISK** Risk for infection as evidenced by risk factors: indwelling urinary catheter.	• No catheter-associated UTI.	• Keep the drainage system closed. • Perform daily perineal care. • Encourage oral fluids. • Remove the catheter as soon as it is no longer indicated.	• Achieved if no signs of UTI appear.

* **References for this care plan:** [28], [35], [37], [110]. Full entFull entries are listed in the References section of the CPMN book.