5.18 Delirium*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Family say, "He is not himself since yesterday." **Objective** • Attention comes and goes • Disoriented • Agitated at night	**ACTUAL** Acute confusion related to infection, medication effects and dehydration as evidenced by fluctuating consciousness and disorientation.	• Back to his usual orientation as the cause is treated.	• Work with the medical team to find and treat the cause. • Reorient him often and make sure he wears his glasses and hearing aids. • Encourage the family to be present. • Support sleep at night and activity by day. • Avoid restraints and unnecessary sedatives.	• Goal reached once he is oriented.
Subjective • Family say, "He sleeps all day and is awake at night." **Objective** • Awake at night • Sleeps by day	**ACTUAL** Ineffective sleep pattern related to confusion as evidenced by night waking.	• Sleeping at night and awake by day within 3 days.	• Keep noise and light low at night. • Open the curtains and keep him active during the day. • Avoid waking him at night unless necessary.	• Met as sleep improves.
Subjective *No specific complaint* **Objective** • Agitated • Tries to climb out of bed	**RISK** Risk for adult falls as evidenced by risk factors: confusion and agitation.	• No falls.	• Put fall precautions in place. • Observe him closely. • Keep the call bell and his things within reach.	• Achieved if he does not fall.

* **References for this care plan:** [35], [51], [58], [126]. Full entries are listed in the References section of the CPMN book.