6.12 Hernia Repair*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "The groin wound hurts when I move." • Rates pain 5/10 **Objective** • Groin incision • Guards when repositioned • HR 96/min	**ACTUAL** Acute pain related to the surgical incision as evidenced by a pain score of 5/10 and guarding.	• Pain at or below 3/10 within an hour of analgesia.	• Assess pain regularly and after each intervention. • Give analgesia as prescribed. • Show him how to support the wound with a hand or pillow when coughing or getting up. • Use a scrotal support or ice pack for scrotal swelling if ordered.	• Achieved when pain is 3/10 or less. • Otherwise, review analgesia with the surgeon.
Subjective • "When can I go back to lifting at work?" **Objective** • Manual job • Cannot state his activity limits	**ACTUAL** Inadequate health knowledge related to post-operative activity restrictions as evidenced by questioning and inability to state limits.	• Before discharge, states his lifting limits and wound care.	• Advise no heavy lifting or straining for the time the surgeon advises. • Teach how to avoid constipation with fluids and fiber. • Teach wound care and the signs of infection. • Ask him to report a new bulge or severe pain.	• Outcome reached when he states the restrictions correctly.
Subjective • "I have not passed urine since the operation." **Objective** • Spinal or general anesthetic • Opioid analgesia • Older man	**RISK** Risk for urinary retention as evidenced by risk factors: anesthesia, opioids and inguinal surgery.	• Voids within 6–8 hours of surgery.	• Note the time and volume of the first void. • Palpate or scan the bladder if he has not voided. • Help him stand to void if allowed. • Catheterize as ordered if he cannot void.	• Considered met when voiding is adequate.

*References for this care plan: [14], [32], [35], [93]. Full entries are listed in the References section of the CPMN book.