6.15 Hemorrhoids and Hemorrhoidectomy*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "It hurts to sit and to pass stool." • Rates pain 7/10 **Objective** • Perianal swelling • Shifts his weight when sitting	**ACTUAL** Acute pain related to inflamed hemorrhoids or surgical trauma as evidenced by a pain score of 7/10 and difficulty sitting.	• Pain at or below 3/10 within 24 hours.	• Give analgesia regularly, especially before the first bowel action. • Offer warm sitz baths 2–3 times a day, as ordered. • Provide a soft cushion; suggest lying on his side. • Apply topical treatment as prescribed.	• Met when pain is controlled.
Subjective • "I am afraid to open my bowels." **Objective** • No stool for 3 days • History of hard stools	**ACTUAL** Impaired intestinal elimination related to fear of pain on defecation as evidenced by no stool for 3 days.	• Soft stool passed within 48 hours.	• Give the prescribed stool softener. • Encourage 2–2.5 L of fluid a day and a high-fiber diet. • Advise against straining and sitting long on the toilet.	• Achieved once a soft stool is passed.
Subjective *No specific complaint* **Objective** • Anorectal surgery • One pad soaked since surgery	**RISK** Risk for excessive bleeding as evidenced by risk factors: surgical wound in vascular tissue.	• No significant bleeding.	• Check dressings and pads. • Follow vital signs. • Tell him to report heavy bleeding or clots after discharge.	• Outcome reached if there is no significant bleeding.

*References for this care plan: [32], [35], [37]. Full entries are listed in the References section of the CPMN book.