7.13 Urinary Incontinence*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I leak urine when I cough, sneeze or laugh." **Objective** • Stress incontinence • Several previous births • BMI 31 kg/m²	**ACTUAL** Stress urinary incontinence related to weak pelvic floor musculature as evidenced by leakage with increased intra-abdominal pressure.	• Less leakage within 6–12 weeks.	• Teach pelvic floor muscle training and check her technique. • Encourage daily practice over the long term. • Advise weight loss and treatment of constipation or chronic cough. • Ask her to keep a bladder diary. • Refer to a continence specialist if there is no improvement.	• Achieved as episodes become fewer.
Subjective • "I stopped going to family visits because of the smell." **Objective** • Avoids going out • Uses many pads	**ACTUAL** Inadequate social connectedness related to fear of leakage and odor as evidenced by avoidance of social activities.	• Takes up at least one social activity again.	• Discuss suitable continence products. • Plan toilet visits before outings. • Encourage her to talk about how she feels.	• Outcome reached once an activity is resumed.
Subjective *No specific complaint* **Objective** • Skin often damp • Wears pads	**RISK** Risk for impaired skin integrity as evidenced by risk factors: moisture from urine.	• Perineal skin stays intact.	• Teach gentle washing and drying after each episode. • Apply barrier cream. • Change pads promptly.	• Considered met if the skin stays intact.

* **References for this care plan:** [32], [35], [92], [126]. Full entries are listed in the References section.