8.13 Diabetes Insipidus*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I drink and pass urine all the time." **Objective** • Urine output 6 L a day • Low urine specific gravity • Na^+ 150 mmol/L	**ACTUAL** Inadequate fluid volume related to excessive dilute urine output as evidenced by polyuria and hypernatremia.	• Intake and output balanced; sodium normal.	• Measure urine hourly and check specific gravity. • Give desmopressin as prescribed. • Replace fluids as ordered. • Weigh daily.	• Achieved when balance is achieved.
Subjective • "I cannot sleep because I go to the toilet." **Objective** • Nocturia • Tired by day	**ACTUAL** Ineffective sleep pattern related to nocturia as evidenced by waking to void.	• Sleeps better within 1 week of treatment.	• Give desmopressin at the prescribed time, often at night. • Keep a urinal at the bedside. • Cut evening fluids only if advised.	• Outcome reached as sleep improves.
Subjective *No specific complaint* **Objective** • On desmopressin • Headache	**RISK** Risk for impaired water-electrolyte balance as evidenced by risk factors: desmopressin-induced water retention.	• Serum sodium stays normal.	• Follow sodium. • Teach him to drink only to thirst. • Ask him to report headache, nausea or confusion.	• Considered met if sodium remains normal.

*References for this care plan: [32], [35], [37], [128]. Full entries are listed in the References section of the CPMN book.