9.5 Hip Fracture and Internal Fixation*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My hip hurts after I fell." • Rates pain 8/10 **Objective** • Leg shortened and externally rotated • Aged 80 • Day 1 after fixation	**ACTUAL** Acute pain related to fracture and surgery as evidenced by a pain score of 8/10.	• Pain at or below 3/10 while taking part in mobilization.	• Give regular analgesia; consider a nerve block as ordered. • Place a pillow between her legs as ordered. • Give analgesia before physiotherapy.	• Goal reached when pain is controlled.
Subjective • Confused at night **Objective** • Newly disoriented • Older adult on opioids	**ACTUAL** Acute confusion related to surgery, pain and medication as evidenced by new disorientation.	• Back to her usual orientation.	• Screen for delirium every day. • Reorient her and involve the family. • Treat pain, dehydration and constipation.	• Met once she is oriented.
Subjective *No specific complaint* **Objective** • Older adult • Immobile after hip surgery	**RISK** Risk for adult pressure injury as evidenced by risk factors: advanced age, immobility and surgery.	• Skin intact.	• Turn every 2 hours as permitted. • Use a pressure-relieving mattress. • Inspect heels and sacrum every shift. • Mobilize early as ordered.	• Achieved if the skin stays intact.

*References for this care plan: [19], [32], [35], [58], [60]. Full entries are listed in the References section of the CPMN book.