Notes on Thesis Corrections

Simon Hulse

19/1/2024

General

I have made some stylistic changes to the thesis (specifically, the formatting of the title page and chapter titles). Ralph asked about Oxford formatting guidelines. I am unaware of Oxford having specific thesis format guidelines, and haven't been able to find any guidance on this, so I am assuming that my thesis format is acceptable.

Spaces were added to citations.

In the corrected thesis, corrections which aren't simply typo fixes are in red.

Corrections by page

- Page 3 Specified that only nuclei with odd numbers of protons and/or neutrons possess spin.
- Page 3 Mentioned molecular rotation as an additional source of angular momentum.
- Page 3 Equation 1.1: Squared the reduced Planck constant.
- Page 3 Replaced "with non-zero spin" with "with spin".
- Page 4 Table 1.1: Updated caption to include source of gyromagnetic ratios.
- Page 4 Replaced "which are spin-0" with "which do not possess spin".
- Page 5 Figure 1.1: Mentioned the sign of γ for each nucleus in the caption.
- Page 6 Mentioned that the high temperature approximation is applied in arriving at Equation 1.8. See also Footnote ‡.

- Page 7 Re-worded description of RF pulse. See also Footnote ||.
- Page 7 Equation 1.13: Corrected the expressions for \tilde{i} and \tilde{j} .
- Page 8 Included a more detailed qualitative description of relaxation.
- Page 10 Explicitly mentioned the presence of a vacuum chamber.
- Page 10 Included solid-state NMR as an area that requires very high field strengths.
- Page 11 "inhomogeneities" → "small inhomogeneities".
- Page 11 A few extra details about the probe.
- Page 12 "is sent to" \rightarrow "travels to".
- Page 12 Peter suggested that I mention how data could be treated if there were line-shape distortions (annotation on Section 3.1.2 heading). Footnote ‡‡ mentions reference deconvolution as a means of correcting these in order to yield Lorentzian line-shapes.
- Page 12 Replace "sweep width" with "spectral width". This has been done in numerous places in the thesis.
- Page 13 Equation 1.18b: Correct equation label.
- Page 17 Mentioned that exponential broadening is not the optimal window function for sensitivity enhancement. See also Footnote ¶¶.
- Page 17 Improved comparison of Gaussian vs Lorentzian lineshapes.
- Page 17 Reworded paragraph on truncation artefacts.
- Page 18 Elaborated on Kramers-Kronig relations, and included a citation.
- Page 18 Added Footnote ††† to mention the lock's use of dispersion lineshapes for monitoring field drifts.
- Page 22 Removed footnote discussing consideration of linewidth for T_2 measurement, as this is not reliable.

- Page 26 Ralph commented that iterative methods are employed routinely in ¹³C NMR for metabolomics fingerprinting. I am unaware of this; from what I am aware, the typical method of performing metabolomics fingerprinting is to break up spectrum into small regions (bins), integrate these bins, and then input the integrals into some routine for multivariate analysis, such as PCA. I have added the phrase "like VARPRO and AMARES" to clarify what I mean by an "iterative method".
- Page 27 Improved wording of why holistically analysing a 2D dataset can be better than sequentially analysing 1D increments.
- Page 31 Added citations to make the fact that the routine makes use of previous theory more explicit.
- Page 32 Peter mentioned that a square root was absent in the probability density (what was Equation 2.3 in my pre-viva draft). The square root wasn't present as the expression was the product of the PDFs of the real and imaginary components of a particular datapoint. I have rewritten this to make the origin of the scaling factor more clear.
- Page 34 Added Footnote † to give the definition of a matrix pencil.
- Page 54 Added Footnote ¶¶ to provide a bit more context on why 25 was chosen as the number of iterations between negative amplitude checks.
- Page 55 Brief discussion about limitations on FID size, based on RAM requirements, in the context of Pines' 1997 paper on the MPM.
- Page 56 Figure 2.4: Made caption and y-axis texts equivalent. Indicated where 96MiB appears on the MPM peak memory usage plot, to help with the 1997 paper discussion.
- Page 66 Mentioned that the filtering process could likely be replaced by the simpler method of using a rectangular filter and slicing at the filter boundaries.
- Page 71 Figure 3.1: Replaced landscape figure with portrait version. Included a description of the different peak colours in the caption.
- Page 74 Figure 3.2: Replaced landscape figure with portrait version. Added structure of andrographolide.
- Page 77 Figure 3.3: Added structure of cyclosporin, and edited the caption accordingly.

Page 79 – Replaced "the spin tumbles" with "the molecule that the spin is associated with tumbles".

Page 82 - Figure 3.4: Tweaked the relative widths of pulses and gradients.

Page
$$83 - T_1 \ll T_2 \to T_1 \gg T_2$$
.

Page 83 – Fixed unit of constant *c* in Stejskal-Tanner.

Page 90 – Provided an explicit expression for the errors associated with D, T_1, T_2 measurements. A derivation of a similar expression for FID parameter estimate errors is given in the appendix.

Page 91 – Figure 3.5: Edited panel d; changed the aspect ratio and viewing angle. Hopefully the lineshapes look better with the altered view. If you still think it looks odd, I could simply remove the panel altogether; it is not the most crucial aspect of the figure, though I think it is nice to give the reader an idea of how the oscillators vary across increments.

Page 93 – Figure 3.6: Replaced landscape figure with portrait version. Added structure of andrographolide.

Page 94 – Figure 3.7: Added structures of valine, threonine and major anomeric forms of glucose.

Page 95 – Brief mention of likely cause for difference in diffusion constant between glucose anomers.

Page 102 – Reworded a sentence in the summary.

Page 103 – Peter queried my statement that data sensitivity is proportional to \sqrt{NS} . I believe this to be correct; when the number of scans is quadrupled, a doubling in sensitivity is expected.

Page 105 – Figure 4.1: TODO: perhaps have a figure with a strongly-coupled *IS* spin system? could then be explicit about positions of strong coupling artefacts (Thrippelton, Keeler paper)

Page 118 – Figure 4.6: Added structure of strychnine.

Page 120 – Figure 4.7: Added structure of quinine.

Page 122 – Figure 4.8: Added structure of camphor. Included a description of grey points in the figure.

Page 123 – Figure 4.9: Replaced landscape figure with portrait version. Added structure of dexamethaone. Included a description of grey points in the figure.

Page 125 – Figure 4.10: Added structure of estradiol. Included a description of grey points in the figure. Page 138 – Added a discussion of how multiprocessing could be employed to improve NLP efficiency.