Experiment 8: The Jumping Pot — Macroscopic Work Against Gravity

Experimental Protocol

Container: Tramontina stainless steel triple-bottom leiteira (same vessel as Experiments 3–6). Liquid: ~300 ml water with chalk and detergent, pre-heated to ~75–80°C before transfer to the stainless steel vessel. Heat source: Domestic gas flame, maximum intensity. Thermometry: Fast Digital Thermometer (probe, immersed in liquid) + Handheld IR thermometer (surface). Recording: Continuous smartphone video at 30 fps. Video: youtube.com/shorts/RDqMcJBkYac

CRITICAL DETAIL: The mixture was initially heated in an aluminum pot, then transferred to the stainless steel vessel at ~75–80°C. The stainless steel triple-bottom conducts thermal gradients more abruptly than aluminum (which distributes heat uniformly). This creates a sharper gradient at the steel-liquid interface upon flame ignition — and this sharper gradient is what produces the anomalous mechanical response.

Sub-Experiment A: The Jump (C1)



Figure 42: C1 — THE JUMPING POT: FRAME-BY-FRAME MILLISECOND ANALYSIS. Video: youtube.com/shorts/RDqMcJBkYac. T = 95.5°C, IR = 96.0°C. Flame ON. Four frames spanning 133 milliseconds (frame 118 at 03:900 to frame 122 at 04:033). The liquid surface shows violent displacement — this is NOT internal convection. The entire pot physically lifts off the stove grate and impacts back down. The audio track confirms: a sharp metallic impact sound at the exact moment of the visual displacement (audio peak at –19.0 dB, ~37 dB above ambient — thousands of times louder than background). NEWTON'S THIRD LAW INTERPRETATION: If the pot jumps UP, something pushed it UP. The thermal field attracts the liquid DOWNWARD toward the flame ($F_{thermal} = -k_T m\nabla T$). The reaction force (Newton's Third Law) pushes the container UPWARD. The instantaneous establishment of a steep thermal gradient (~1200°C flame vs ~80°C liquid) across the triple-bottom steel base creates an impulsive attractive force on the liquid mass. The liquid is pulled down; the pot is pushed up. FALSIFICATION: Statistical mechanics does not generate impulsive macroscopic work against gravity. Random molecular motion cannot produce a coherent unidirectional force sufficient to lift a metal pot containing ~300 ml of liquid. This requires a directed force — a field.

Sub-Experiment B: Flame OFF — Vapor Explosion with IR Confirmation (C2)



Figure 43: C2 — FLAME EXTINCTION: VAPOR EXPLOSION AT CONSTANT TEMPERATURE. Eight frames in 4 pairs spanning 2.3 seconds (49:700 to 52:005). [F1, 49:700, T=100.5°C, IR=104.4°C]: Flame extinguishing (red annotation marks dying flame). Vapor: minimal/absent. Active boiling motion visible. [F2, 50:339, T=100.5°C, IR=104.5°C]: Flame OFF. Vapor becoming visible. Temperature

UNCHANGED — probe still 100.5°C, IR still 104.5°C. [F3, 50:433, T=100.5°C, IR=104.5°C]: Vapor clearly visible, expanding. Only 733 ms since flame extinction. [F4, 52:005, T=100.5°C, IR=104.4°C]: Vapor EXPLODING upward — massive visible plume. Temperature STILL 100.5°C. IR STILL 104.4°C. THE TRIPLE FALSIFICATION IN ONE FRAME SET: (1) CONTRADICTION: Temperature unchanged at 100.5°C while motion ceases → temperature does not drive motion. (2) ABSURDITY: Mainstream claims the air above the liquid 'cools rapidly' to condense invisible vapor → but the IR reads 104.4°C at the surface. The surface is HOTTER than the boiling point. Nothing is cooling. (3) REFUTATION: The vapor erupts while the surface is at 104.4°C → the 'cold air condensation' hypothesis is directly measured and falsified by the IR thermometer in the same frame.

Sub-Experiment C: Second Cycle — Reproducibility (C3)

Figure 44: C3 — SECOND FLAME EXTINCTION CYCLE: IDENTICAL PATTERN. Eight frames in 4 pairs spanning 1.2 seconds (1:18:530 to 1:19:750). [F1, 1:18:530, T=100.5°C, IR=105.3°C]: Flame still ON but extinguishing. Vapor minimal. Red annotation marks the dying flame. [F2, 1:18:600, T=100.6°C, IR=105.3°C]: Flame OFF. Vapor OFF. 70 ms after extinction. [F3, 1:19:235, T=100.6°C,

IR=105.6°C]: 705 ms post-extinction. Vapor already clearly visible. IR has INCREASED to 105.6°C — surface getting HOTTER after flame removal. [F4, 1:19:750, T=100.6°C, IR=105.3°C]: Vapor fully developed, expanding plume. REPRODUCIBILITY: Identical pattern in both cycles. Flame OFF → vapor erupts → temperature unchanged/increasing. CONNECTION TO ATMOSPHERIC RETENTION: The vapor behavior demonstrates the thermal component of gravity. During active heating, the thermal field traps vapor within/near the liquid surface (attraction phase). When the field collapses (flame OFF), vapor escapes explosively. At planetary scale, Earth's thermal field (surface ~288K vs space ~2.7K, $\nabla T \approx 285K$ across the atmosphere) performs the same function: retaining atmospheric gases against the vacuum of space. This resolves the Venus-Mars paradox: Venus (extreme surface heat, dense atmosphere) retains its atmosphere via thermal attraction despite lacking a strong magnetic field. Mars (cold surface, thin atmosphere) cannot. Atmospheric retention is primarily thermal, not magnetic — consistent with Gravity = Magnetism + Heat.

Anomaly Analysis — Experiment 8

Anomaly J1: Macroscopic mechanical work against gravity — pot jumps. A stainless steel pot containing ~300 ml of liquid physically lifts off the stove grate upon flame ignition or during active heating. The jump is audible (metallic impact, –19 dB peak vs –58 dB ambient) and visible in frame-by-frame analysis (133 ms event). Statistical mechanics provides no mechanism for random molecular motion to produce coherent unidirectional force lifting a solid container against gravity. The thermal field framework explains it via Newton's Third Law: F_thermal attracts liquid downward → reaction force pushes pot upward.

Anomaly J2: Vapor erupts while surface temperature is 104–106°C. Mainstream explanation for post-extinction vapor: 'the air above the surface cools rapidly, condensing invisible vapor.' IR measurement at the moment of vapor eruption: 104.4–105.6°C. The surface is 4–6°C ABOVE the boiling point. Nothing is cooling. The explanation is falsified by direct measurement in the same frame as the phenomenon.

Anomaly J3: Vapor increases while temperature is constant or increasing. Between C2-F1 (IR 104.4°C) and C2-F4 (IR 104.4°C), vapor goes from zero to explosive — at constant temperature. Between C3-F2 (IR 105.3°C) and C3-F3 (IR 105.6°C), vapor develops while IR INCREASES by 0.3°C. The 'cooling' hypothesis fails in both directions: temperature doesn't decrease, and when it changes, it increases.

Anomaly J4: From contradiction to absurdity to refutation. Step 1 (Contradiction): The metal pot is hotter than the liquid, so the liquid should keep heating — yet motion stops instantly when flame is removed. If temperature drives motion, motion should persist. Step 2 (Absurdity): To explain the vapor, mainstream invokes rapid air cooling above the surface in <0.5 s — while the pot is still radiating heat into that air from all sides. Step 3 (Refutation): IR measurement shows the surface at 104–106°C during vapor eruption. The premise of the explanation (surface cooling) is directly measured and shown to be false.

Systematic Falsification — Experiment 8

Classical Prediction	Observation	Status
Statistical motion cannot lift solids	Pot jumps off grate (audible + visible)	FALSIFIED
Vapor requires surface cooling	IR = 104–106°C during vapor eruption	FALSIFIED
Motion driven by temperature	Motion stops at 100.5°C when flame OFF	FALSIFIED
Air cools surface in <0.5 s	IR increases 105.3→105.6 post-extinction	FALSIFIED
Vapor ∝ heating intensity	Vapor absent with flame, erupts without	FALSIFIED

Connection to Gravity = Magnetism + Heat

Experiment 8 provides three independent lines of evidence for the thermal component of gravity:

(1) The jump: The pot lifts against gravity via Newton's Third Law reaction to thermal attraction of liquid toward the flame. This is the same force observed in Experiments 1–2 (water/butter flowing uphill), now producing macroscopic mechanical work on a solid object. The force is not statistical — it is directional, impulsive, and sufficient to overcome gravity for a metal pot with ~300 ml of liquid.

(2) Vapor trapping and release: During active heating, the thermal field traps vapor near the liquid surface (attraction phase of the thermal cycle). Upon field collapse (flame OFF), vapor escapes explosively. At planetary scale, this mechanism explains atmospheric retention: Earth's thermal gradient (surface ~288K, space ~2.7K) creates a thermal attraction field that retains atmospheric gases. Venus (extreme surface heat) retains a dense atmosphere despite no magnetic field. Mars (cold surface) cannot retain its atmosphere. The thermal component of gravity, not the magnetic component, is the primary mechanism for atmospheric retention.

(3) IR confirmation: The surface being hotter than the bulk (IR > probe) during active heating, combined with vapor suppression during heating and eruption during cooling, demonstrates that the thermal field actively organizes matter distribution (liquid below, vapor trapped). This organization collapses when the field collapses — identical to gravitational field behavior.

Cross-Experiment Unified Analysis

Comparative Parameter Table

Param	Exp 1 Iceberg	Exp 2 Butter	Exp 3 Water	Exp 4 Milk	Exp 5–6 Beans	Exp 7 Ice	Exp 8 Jump
Material	H■O	Lipid	H■O	Milk	H■O+ beans	H■O/ ice	H■O+ chalk
T range	0–100	25–65	72–100	69–101	16–39	–18–25	80–106
Key	Uphill flow	Curved path	27.3°C inv.	0.3°C cycle	–1.5°C 17min↑	Ice contracts	Pot jumps
Falsif.	P1–3	P4–5	P6–9	P8–9	P10–12	P13–18	P19–23

Universal Principle Statement

Eight experiments — water/ice, lipid/butter, water at boiling, milk emulsion, water with beans+lentils (short and long), ice in multiple container materials, and water/chalk/detergent in steel — in different phase regimes, different temperature ranges, different gradient magnitudes, different timescales — produce convergent evidence that: **(1) bulk mechanical and thermal behavior is determined by active thermal flux, not absolute temperature; (2) ice expansion is not intrinsic to water but is mediated by the electromagnetic environment of the container; (3) structured vesicular entities are organized material structures, not gas bubbles; (4) the thermal field performs macroscopic mechanical work against gravity and traps vapor near liquid surfaces.**

Experiments 1–2 demonstrate spatial transport against gravity. Experiments 3–4 demonstrate temporal correlation (motion tracks flame state). Experiments 5–6 demonstrate thermal absorption anomalies (temperature drops with flame, rises without flame) and inverted gradients (bottom colder than top with heating from below). Experiment 7 demonstrates that ice expansion is container-dependent (glass/copper fail, plastic contracts), that SVEs are structured entities surviving ~300 atm, and that the electromagnetic seal state of the container determines SVE behavior during phase transition. Experiment 8 demonstrates macroscopic work against gravity (pot jump), direct IR falsification of the 'air cooling' hypothesis for post-extinction vapor, and provides the empirical basis for thermal atmospheric retention (Venus–Mars paradox resolution).

The thermal force law $F■_{thermal} = -k_T m\nabla T$ unifies all observations. The convective cycle is attraction→torque→repulsion (thermal inverse of magnetic repulsion→attraction). $k_T \approx (4.3–6.4)\times10^{-3}$ $m\cdot s^{-2}\cdot K^{-1}$ (first terrestrial calibration).

Conclusion

Eight independent kitchen-scale experiments falsify twenty-three classical predictions and establish fourteen new empirical facts. The thermal force law $F■_{thermal} = -k_T m\nabla T$ unifies observations from Experiments 1–6 and 8, while Experiment 7 extends the framework to phase transition physics, demonstrating that ice expansion is not an intrinsic property of water but is mediated by the electromagnetic environment of the container. Experiment 8 provides the most dramatic evidence: a metal pot with liquid physically jumps off the stove grate — macroscopic work against gravity produced by a thermal gradient — and IR-confirmed vapor eruption at surface temperatures of 104–106°C directly falsifies the mainstream 'air cooling' hypothesis. The vapor trapping/release mechanism models atmospheric retention, resolving the Venus–Mars paradox.

This framework, combined with the result Gravity = Magnetism + Heat, implies that gravitational attraction contains a thermal component of fundamental significance.

References

1. Clausius, R. (1865). The Mechanical Theory of Heat. London: John van Voorst.
2. Boltzmann, L. (1877). Über die Beziehung... Wiener Berichte, 76, 373–435.
3. Prigogine, I. (1978). Time, Structure, and Fluctuations. Science, 201(4358), 777–785.
4. Maxwell, J.C. (1871). Theory of Heat. Longmans, Green, and Co.
5. Scriven, L.E., & Sternling, C.V. (1960). The Marangoni Effects. Nature, 187, 186–188.
6. Landauer, R. (1961). Irreversibility and heat generation. IBM J. Res. Dev., 5(3), 183–191.
7. Guzzon, A.S. (2026). Gravity = Magnetism + Heat. Zenodo. DOI:10.5281/zenodo.18498514

Supplementary materials: Google Drive · YouTube: @5incere5cience · TikTok: @5incere5cience

Published under Creative Commons Attribution 4.0 International (CC BY 4.0).

Video: https://youtube.com/shorts/aXFAot2Q-QE
Supplementary materials: Google Drive · YouTube: @5incere5cience · TikTok: @5incere5cience

Published under Creative Commons Attribution 4.0 International (CC BY 4.0).