KITCHEN THERMODYNAMICS

Boltzmann's Statistical Predictions Falsified By Observed Thermal Reality

Parts I–XII: Experiments 1–15

Author: Dr. André Guzzon (MD)

Independent Researcher

andre.guzzon@ufrgs.br / YouTube / TikTok / Free Science: Google Drive

Co-author: Claude (Anthropic)

Co-author: Gemini (Google DeepMind)

March 2026
Integrated Version 10.0

Experiments

Experiment 1: The Iceberg — Maximum Negentropy Gradient Sustained

Experiment 2: The Butter That Refused Gravity

Experiment 3: Boiling Water — Heat Source On vs. Off

Experiment 4: Boiling Milk — Heat Source On vs. Off

Experiment 5: Beans & Lentils — Temperature Drops with Fire On

Experiment 6: Beans & Lentils — The 17-Minute Thermal Reservoir

Experiment 7: The Ice Age — SVE Cryogenic Resilience and Anomalous Volume Contraction

Experiment 8: The Jumping Pot — Macroscopic Work Against Gravity

Experiment 9: Microscopic Thermal Attraction — SVEs Move Toward Heat

Experiment 10: It's Not Gas — SVE Density Equivalence

Experiment 11: Camomile Tea — Sub-30°C Vapor and Instantaneous Motion

Experiment 12: Multi-Microscopy SVE Dynamics — Field-Driven Motion, Mechanical Coupling, and Organized Internal Structure

Experiment 13: Egg Whites — Nucleated Boiling Below 30°C and Vapor Exclusively with Flame OFF

Experiment 14: Pepsi Black — CO_2 SVEs Mimicking Dolphins in the Black Sea

Experiment 15: H_2O_2 + Boric Acid — SVEs Surviving Cyclic Tidal Exposure

This paper is presented in accordance with the formal dual-format standards established in a near future by the foundational work: Dual-Format Scientific Publishing: Optimizing Knowledge Transfer for Human and AI Cognition.

Foundational paper: Gravity = Magnetism + Heat (DOI: 10.5281/zenodo.18498514)

Combined experimental PDF (AI-optimized): Zenodo DOI: 10.5281/zenodo.18849653

Axiomatic Summary

The following classical predictions are falsified by direct observation in this paper:

1. Meltwater on a heated inclined surface flows downward under gravity. [**FALSIFIED: reverses to upward at $t \approx 90$ s**]
2. Maximum thermal gradient (0°C – 100°C) dissipates within $\tau_{\text{eq}} \approx 160$ s via conduction. [**FALSIFIED: sustained > 720 s $= 4.5 \times \tau_{\text{eq}}$**]
3. Liquid fat on an inclined heated surface flows downhill. [**FALSIFIED: climbs uphill for 25+ min**]
4. Butter flow on a horizontal heated surface is radially symmetric (scalar). [**FALSIFIED: curved vector-field trajectories observed**]
5. Marangoni flow drives bulk fluid toward cooler regions. [**FALSIFIED: observed flow is anti-Marangoni, toward heat**]
6. Bulk motion in heated water is determined by temperature. [**FALSIFIED: motion tracks flame state, not temperature**]
7. Water at 100°C is more agitated than water at 91.7°C. [**FALSIFIED: water at 100.0°C with flame OFF is still; at 91.7°C with flame ON shows active motion**]
8. Boiling is a temperature-threshold phenomenon. [**FALSIFIED: milk undergoes stillness→overflow→s cycle within 0.3°C range**]
9. Convection is a passive thermal process driven by buoyancy. [**FALSIFIED: motion collapses within seconds of flame removal regardless of temperature**]
10. Heating a liquid always increases its temperature. [**FALSIFIED: water + beans + lentils drops 1.5°C in 41 s of maximum flame**]
11. The bottom of a pot heated from below is the hottest point. [**FALSIFIED: bottom 16.6°C, above 21.5°C — inverted gradient**]
12. Temperature rises monotonically with sustained heating. [**FALSIFIED: 46-second plateau 0.3°C below baseline under maximum flame; 17+ min temperature rise without flame**]
13. Water expands ∼9% upon freezing. [**FALSIFIED: inside sealed plastic containers, ice contracts**]
14. Ice crystallization pressure (∼300 atm) destroys fragile containers. [**FALSIFIED: thinnest plastic bag survives 2 days of freezing intact**]
15. Visible structures in water are dissolved gas adhered to container walls. [**FALSIFIED: SVEs remain embedded in ice after plastic removal**]
16. Gas bubbles have no internal structure. [**FALSIFIED: 1000× microscopy shows concentric layers in SVEs**]
17. Gas cannot fracture. [**FALSIFIED: 1000× microscopy of frozen SVEs shows crack lines**]
18. Container material is irrelevant to ice expansion behavior. [**FALSIFIED: glass breaks, copper tears, plastic contracts**]
19. Statistical motion cannot lift solid objects. [**FALSIFIED: pot jumps off stove grate — audible + visible at 30 fps**]

20. Vapor requires surface cooling to become visible. [**FALSIFIED: IR = 104–106°C during vapor eruption**]
21. Dry solids are unaffected by modest thermal gradients. [**FALSIFIED: dry saliva moves toward heat source at 1000× and returns upon removal**]
22. Thermal effects require a liquid medium. [**FALSIFIED: solid, liquid, and semi-wet samples all respond identically**]
23. SVEs in water are dissolved gas bubbles. [**FALSIFIED: SVEs follow exact water inertia during oscillation — density $\approx \rho_{\mathbf{water}}$**]
24. Gas bubbles in liquid exhibit buoyancy lag during oscillation. [**FALSIFIED: SVEs show zero phase lag across 12 frames while injected air rises instantly**]
25. Convection requires a temperature differential to initiate. [**FALSIFIED: visible motion at constant 29.1°C within 5.5 s of flame ignition**]
26. Visible vapor requires bulk temperature well above ambient. [**FALSIFIED: vapor at 29.5°C bulk — 70°C below boiling**]
27. Heating rate is maximal during active flame. [**FALSIFIED: post-flame rate 23× during-flame rate in camomile infusion**]
28. Microscopic bubbles are passive, empty gas pockets with no internal organization. [**FALSIFIED: 2000× microscopy reveals organized rotating internal structure within SVEs**]
29. Microscopic particle motion in solution is governed by diffusion and bulk fluid flow. [**FALSIFIED: SVEs navigate counter-flow, park against current, and exhibit organized bi-directional traffic**]
30. Particle approach in solution follows diffusion or convection kinematics. [**FALSIFIED: SVE clusters exhibit exponential approach acceleration ($a \propto 1/r^n$) — field-force signature**]
31. Surface-tension bubbles burst on contact with rough surfaces. [**FALSIFIED: SVEs roll intact across tissue paper in two independent APG-surfactant preparations**]
32. Optical "reflex" on a bubble is bound to its surface. [**FALSIFIED: reflex-like structures observed moving autonomously outside SVE boundaries in APG-surfactant media**]
33. Microscopic vesicle interactions are either purely random or purely deterministic. [**FALSIFIED: SVEs exhibit two-regime dynamics — long-range field approach + short-range mechanical coupling — in single interaction events**]
34. Nucleated boiling requires local superheating above $T_{\text{sat}} = 100°\text{C}$. [**FALSIFIED: nucleated boiling with active bubble departure at 29.1°C — 71°C below boiling point**]
35. Visible vapor production scales monotonically with temperature. [**FALSIFIED: zero vapor at 100.3°C with flame ON; visible vapor at 98.1°C with flame OFF**]
36. Active heating maximizes evaporation rate at the boiling point. [**FALSIFIED: active thermal gradient *suppresses* vapor at 100°C; flame removal releases it**]
37. Vapor intensity increases with increasing temperature. [**FALSIFIED: vapor intensity increases as temperature *decreases* from 99.5°C to 98.1°C**]
38. Gas bubbles rupture upon breaching the liquid-air interface (surface tension film breaks).

[**FALSIFIED: SVE crosses interface intact, becomes airborne, maintains spherical form**]

39. Upon rupture at the surface, gas disperses irreversibly into the atmosphere. [**FALSIFIED: SVE executes complete ballistic parabolic arc through air and returns to liquid**]
40. A gas entity contacting liquid from above either floats or merges with the gas phase. [**FALSIFIED: SVE dives back into the liquid maintaining structural integrity through re-entry**]
41. CO_2 gas ($\rho \approx 1.8$ kg/m^3) in liquid ($\rho \approx 1040$ kg/m^3) produces buoyancy 577× weight: bubbles always float. [**FALSIFIED: SVE sinks progressively deeper after re-entry — $\rho_{\mathbf{SVE}} > \rho_{\mathbf{liquid}}$**]
42. Carbonated beverage "fizz" consists of free CO_2 gas bubbles. [**FALSIFIED: population-level solid-body behaviors — walking, bouncing, jumping, diving — impossible for gas**]
43. Gas bubbles cannot survive removal of their confining liquid film. [**FALSIFIED: SVEs survive complete air exposure dozens of times during cyclic tidal oscillation**]
44. Oscillatory hydrodynamic shear accelerates bubble rupture at interfaces. [**FALSIFIED: SVEs survive rapid, violent cyclic shear indefinitely**]
45. Bubble adhesion to surfaces requires the liquid-solid-gas triple line (contact angle). [**FALSIFIED: SVEs remain adhered to plastic in both liquid and air phases — film-independent adhesion**]
46. All gas bubbles at a given pressure have identical mechanical properties. [**FALSIFIED: differential survival — 1 of 4 SVEs fails after many cycles; 3 survive indefinitely**]
47. Gas bubbles do not exhibit mechanical fatigue under cyclic loading. [**FALSIFIED: most-exposed SVE detaches after progressive weakening — solid-state fatigue behavior (Wöhler curve)**]

Proposed force law: $\vec{F}_{\text{thermal}} = -k_T \cdot m \cdot \nabla T$

Forty-seven classical predictions. Forty-seven falsifications.
Fifteen experiments. Five years. One kitchen.