Optical Anomalies in Macroscopic Vesicles: Empirical Evidence of Non-Gaseous Internal Architecture in Common Fluids

Dr. Andre Guzzon (MD)

Independent Researcher, Brazil

andre.guzzon@ufrgs.br

January 2026

Abstract

The classical physical model of ephemeral foam bubbles defines them as gas-filled cavities within a liquid matrix, governed by surface tension and strictly adhering to Plateau's laws [1]. According to standard geometric optics, such spherical interfaces under a single light source should exhibit a predictable, singular virtual image (specular reflection) on the convex surface [2]. This study presents empirical data that challenges the universality of this model in common organic and inorganic fluid mediums (e.g., coffee infusions, salivary emulsions). Through a comparative analysis using macroscopic photography and high-magnification optical microscopy (up to 2000x), we identify a distinct class of vesicles herein termed **Structured Vesicular Entities (SVEs)**. Unlike simple gas bubbles (Type-A), SVEs (Type-B) exhibit the "Multi-Focal Paradox"—simultaneous, non-concentric points of high luminosity within a single vesicular envelope—and a distinct internal opacity inconsistent with a homogeneous gas phase. This paper establishes a visual classification system distinguishing these two forms, proposing that SVEs represent a ubiquitous, self-organizing phenomenon possessing complex internal architecture.

1 Introduction

1.1 The Prevailing Paradigm

Fluid dynamics and surface chemistry have long categorized foam bubbles as simple physical phenomena: pockets of gas (air, CO_2, water vapor) trapped within a thin film of liquid. The stability of these structures is traditionally attributed to the Marangoni effect [3, 4] and the minimization of surface area, creating characteristic polyhedral geometries in dense foams [5]. In this established view, the interior of a bubble is a transparent void of structural complexity—a passive volume of gas governed by surface tension [1].

1.2 The Optical Anomaly

While this model accurately describes "ideal" soap bubbles under controlled laboratory conditions, it fails to account for optical anomalies observed in naturally occurring foams and emulsions. A fundamental principle of ray optics dictates that a smooth, spherical convex mirror (the outer surface of a bubble) reflects a single primary light source as a single focal point [2]. However, systematic observation of vesicles in everyday fluids reveals a frequent deviation from this rule: the presence of multiple, distinct, and spatially stable bright spots within the boundary of a single vesicle. This phenomenon, which we classify as the **Multi-Focal Paradox**, implies that light is interacting with internal structures capable of independent reflection, rather than merely passing through a homogeneous gas medium.

1.3 Objective

This paper serves as the foundational entry in a series of investigations into these Structured Vesicular Entities (SVEs). We aim to:

1. Establish a rigorous visual classification system distinguishing "Type-A" (Simple/Empty) vesicles from "Type-B" (Structured) entities.
2. Provide microscopic evidence of the internal architecture corresponding to the macroscopic optical anomalies.

We posit that relying solely on established theoretical models has obscured the observation of a complex, self-organizing phenomenon hidden in plain sight. This investigation adopts a radically empirical approach, prioritizing direct observation over theoretical assumption.

2 Methodology

2.1 Sample Selection

To demonstrate the ubiquity of the observed phenomena, samples were selected from common organic and inorganic fluids characterized by spontaneous vesicle formation. Key distinct mediums were analyzed:

- **Control Sample (Type-A):** Standard commercial surfactant foam ("Espuma Mágica"), known to produce gas-filled cavities.
- **Test Sample (Type-B):** Organic infusions (Coffee) and Biological fluids (Saliva), where optical anomalies are frequently observed.

2.2 Imaging Protocol

Macroscopic observations were conducted using high-resolution mobile photography to identify optical anomalies (multi-focal reflections) visible to the naked eye. Microscopic analysis was performed using a digital optical microscope with magnification ranging from 500x to 2000x. Transillumination and incident light techniques were alternated to reveal internal opacity vs. transparency. Dynamic behavior was recorded in high-definition video to document the stability of the internal structures against fluid motion.

3 Results

3.1 Visual Classification: Type-A vs. Type-B

Microscopic analysis revealed two fundamentally distinct vesicular morphologies co-existing in fluid mechanics, which we have classified as follows:

Type-A (The Empty Vesicle): Observed primarily in the control surfactant samples. These structures exhibit thin, transparent walls that often share lamellae with neighboring bubbles (polyhedral stacking), compliant with Plateau's laws. Internally, they are optically empty, allowing light to pass through without internal scattering or secondary reflection.



Figure 1: Type-A vesicles demonstrating classical foam morphology with shared walls and complete internal transparency.

Type-B (The Structured Vesicular Entity - SVE): Observed in organic and biological samples. These entities maintain a strict spherical autonomy, rarely sharing walls even when clustered. Most notably, they possess a defined **Reflective Core Assembly (RCA)**—a central or eccentric internal density that is opaque under transillumination and highly reflective under incident light.



Figure 2: Type-B vesicles (SVEs) showing spherical autonomy and distinct internal reflective cores.

3.2 The Multi-Focal Paradox (Correlated Findings)

The internal architecture of Type-B entities provides the physical basis for the macroscopic optical anomalies. The multiple bright reflection spots observed in the test samples consistently align with the positioning of the Reflective Core Assembly and the thick, refractive outer cortex. The high refractive index of the core suggests it acts as a distinct optical interface within the vesicle, separate from the outer film, a feature entirely absent in Type-A vesicles.



Figure 3: Structural Persistence Comparison. Unlike gas bubbles which collapse into a bulk liquid state, Type-B entities (right) demonstrate structural persistence during the drying phase, maintaining spherical integrity and internal organization compared to the amorphous residue of Type-A foam (left).

3.3 Microscopic Detail

High-magnification analysis further distinguishes the physical boundaries of these entities. Type-A vesicles present as a continuous lattice, whereas Type-B entities present as discrete, autonomous objects.



Figure 4: Type-A Morphology (Detail). High magnification transillumination of standard foam. Note the shared walls and complete transparency, characteristic of a gas-liquid interface minimizing surface area.



Figure 5: Type-B Morphology (Detail). High-magnification of SVEs. Note the thick, independent cortical walls and the prominent Reflective Core Assembly (white centers), indicating substantial internal matter interacting with light.

4 Discussion

4.1 The Impossibility of the Void

The data presented challenges the fundamental assumption that all vesicles in common fluids are merely gas-filled voids. The Laws of Reflection dictate that a transparent, spherical interface cannot produce multiple, non-concentric specular reflections from a single light source. The observation of the "Multi-Focal Paradox" serves as optical proof that the interior of these Structured Vesicular Entities (SVEs) contains reflective material interfaces.

4.2 A Distinct State of Organization

The contrast between Type-A and Type-B vesicles is not merely one of content, but of structural integrity. Type-A vesicles behave as passive gas pockets governed by surface tension minimization (See Supplementary Video S1). Type-B entities behave as autonomous units with a stabilizing internal architecture that resists simple coalescence (See Supplementary Video S2). Whether this organization represents a novel state of soft matter or a biological phenomenon remains to be defined in future studies. However, visually and optically, it is incompatible with the definition of a simple air bubble.

5 Conclusion

This study establishes a critical distinction in fluid mechanics regarding macroscopic vesicles:

- **Type-A Vesicles:** Consistent with gas-filled cavities (empty, transparent).
- **Type-B Entities (SVEs):** Characterized by complex internal architecture and multifocal reflection.

We conclude that many structures colloquially identified as "bubbles" in organic fluids are, in fact, complex entities with a defined, persistent anatomy. This finding necessitates a reevaluation of micro-structures in everyday fluids, suggesting that a ubiquitous, structured phenomenon has been overlooked due to reliance on simplified theoretical models. Future papers in this series will investigate the specific morphology of the internal core and the thermodynamic behavior of these entities.

Supplementary Material

Video S1: Dynamic microscopy of Type-A vesicles (Control), demonstrating passive Plateau borders and lack of internal structure.

Video S2: Dynamic microscopy of Type-B SVEs, demonstrating spherical autonomy and persistent internal reflective cores.

References

[1] Plateau, J. A. F. (1873). *Statique expérimentale et théorique des liquides soumis aux seules forces moléculaires.* Paris: Gauthier-Villars.

[2] Hecht, E. (2017). *Optics* (5th ed.). Boston: Pearson Education.

[3] Marangoni, C. (1865). *Sull'espansione delle goccie d'un liquido galleggianti sulla superficie di altro liquido.* Pavia: Tipografia dei Fratelli Fusi.

[4] Scriven, L. E., & Sternling, C. V. (1960). "The Marangoni Effects". *Nature*, 187, 186–188.

[5] Weaire, D., & Hutzler, S. (1999). *The Physics of Foams.* Oxford: Clarendon Press.

[6] Isenberg, C. (1978). *The Science of Soap Films and Soap Bubbles.* Clevedon: Tieto Ltd.