KITCHEN THERMODYNAMICS

Parts I, II, III, IV, V & VI: Experiments 1–9

Author: Dr. André Guzzon (MD)

Independent Researcher

andre.guzzon@ufrgs.br / Youtube / TikTok / Free Science: Google Drive

Co-author: Claude (Anthropic)

Co-author: Gemini (Gemini Deep Think)

March 2026
Integrated Version 7.0

Contents

Experiment 1: The Iceberg — Maximum Negentropy Gradient Sustained

Experiment 2: The Butter That Refused Gravity

Experiment 3: Boiling Water — Heat Source On vs. Off

Experiment 4: Boiling Milk — Heat Source On vs. Off

Experiment 5: Beans & Lentils — Temperature Drops with Fire On

Experiment 6: Beans & Lentils — The 17-Minute Thermal Reservoir

Experiment 7: The Ice Age — SVE Cryogenic Resilience and Anomalous Volume Contraction

Experiment 8: The Jumping Pot — Macroscopic Work Against Gravity

Experiment 9: Microscopic Thermal Attraction — SVEs Move Toward Heat

This paper is presented in accordance with the formal dual-format standards established in a near future by the foundational work: Dual-Format Scientific Publishing: Optimizing Knowledge Transfer for Human and AI Cognition.

FORMAT HUMAN OPTIMIZED

Abstract

Do you know what statistical thermodynamics is? It's a mathematical model. It's not Physics. The likelihood of water boiling at 100 Celsius without an active heat source is zero. It's easier to overflow boiling milk at 80 Celsius if the fire is on. There's more chance of melting ice climbing an inclined plane if there's an active heat source up there. Where can you see it? In any textbook, this I can guarantee you. However, outside numbers and letters, it has been before your eyes in your kitchen since always. Trust me: I'd not be here if you didn't need me. Popper's philosophy of science asks for a black swan—be my guest, 'cause I got dozens!

Experiments

1. The Iceberg: Maximum Negentropy Gradient Sustained

Water was frozen uniformly in a steel pan. An inclined plane was created. Intense heat was applied at the upper end only.

Classical prediction: Ice melts progressively. Water flows downward under gravity. System equilibrates to uniform temperature within minutes.

Observation: Within the first ninety seconds, initial melted water flows downward as expected. Soon after that, flow reverses. Water flows upward, against gravity, toward the heat source.

After five minutes, the melted water was already boiling upward, and the iceberg had barely melted. All the melted water was already and only moving toward the heat.

For more than twelve minutes there was melted water flying and boiling upward—Leidenfrost effect—with a large portion of the iceberg still frozen at the lower end, always interconnected by liquid water.

This configuration—ice coexisting with boiling water—represents the maximum possible negentropy gradient for water: 0°C and 100°C, solid and gas, bridged by liquid. All three phases of water sustained simultaneously across fifteen centimeters. Classical thermodynamics predicts this gradient should dissipate within minutes through thermal conduction and convective mixing.

The standard escape clause for non-equilibrium organization is Prigogine's dissipative structures—but those still obey total entropy increase. The environment pays the cost. This is different. This is active negentropy generation. Entropy decreases locally and persists.

Conclusion: Thermal gradients actively maintain the maximum possible organization for water against thermodynamic predictions. The gradient does not merely resist dissipation—it sustains a configuration that classical physics says should not exist beyond a few minutes.

2. The Butter That Refused Gravity

Two independent experiments demonstrate the same principle through different mechanisms.

Experiment 2A: The Horizontal Plane — Evidence of Field Structure

A Tramontina stainless steel triple-bottom frying pan was placed on a gas stove, positioned horizontally. Butter at room temperature was placed on the surface. Direct flame was applied beneath.

Two sequential experiments were conducted on the horizontal plane. In the first, butter melted and flowed in a straight line toward the heat source. The movement was rapid—visible in real time, completing in approximately two seconds.

In the second experiment, with butter placed in a different position, the melted butter executed a curved trajectory. The liquid did not flow in a straight line toward maximum heat. Instead, it navigated an arc, flowing up and over the point where the flame contacted the pan beneath.

This curved trajectory is critical. On a horizontal plane, gravity is not a variable—it acts uniformly downward, perpendicular to the surface. If the butter were responding to a scalar effect like thermal expansion or random molecular agitation, the flow would be radial and symmetric around the heat source. A curved path means the butter is navigating a vector field—a field with direction and structure, not merely magnitude.

Experiment 2B: The Inclined Plane — Evidence of Force

Then the configuration changed. A stainless steel baking tray was tilted to create an inclined plane. The angle was verified with an air level. Butter blocks were placed at the lower end. An intense gas flame was positioned at the upper end only, reaching a maximum surface temperature of 65°C.

Classical prediction: butter melts, viscosity drops, gravity pulls the liquid downward toward the lower end of the tray.

Observation: the butter climbed upward.

For twenty-five minutes of continuous time-lapse recording, melted butter flowed up the inclined plane, directly toward the heat source at the top, against the gravitational pull downward.

The butter nearest the heat source liquefied first and began the ascent. The portion farthest from heat remained solid. The middle section split: part of it climbed upward, part of it slid downward. One sample, two directions, simultaneous opposition.

What This Is Not

This is not capillary action—the steel surface is smooth and non-porous. There is nothing to wick.

This is not Marangoni flow—surface tension gradients drive flow *away* from heat, not toward it. The butter went the wrong way for Marangoni.

This is not convection—convection circulates fluid in cells. It does not transport bulk liquid mass unidirectionally up an incline for twenty-five minutes.

Gravity pulls down. Heat pulls up. On this inclined plane, for this butter, heat won.

If thermal energy merely increased random molecular motion, there would be no mechanism for sustained directional flow against gravity. Entropy should increase. Organization should decay. The butter should flow downward.

But the butter climbed.

What Both Experiments Prove Together

Heat is not merely energy dispersal. Heat exerts force. Thermal gradients attract matter.

The horizontal experiment (2A) proves this is a field—the curved trajectory reveals structure and direction, not random dispersion. The inclined experiment (2B) proves this field exerts force—enough to overcome gravity for twenty-five continuous minutes.

Classical thermodynamics predicts equilibration. We observe organization maintained against gravitational potential for twenty-five minutes.

The butter that refused gravity is not an anomaly. It is evidence.

The Principle

These two experiments demonstrate the same principle with different materials (water and lipid), in different regimes (Leidenfrost boiling versus viscous flow), at different temperatures (0–100°C versus 25–65°C), over different timescales (twelve minutes versus twenty-five minutes), with identical result: matter moves upward against gravity, toward the heat source.

This is not an accident of a specific material or a specific configuration. It is a principle. Thermal gradients exert attractive force on matter. That force can overcome gravity. And once established, the resulting organization persists far beyond what thermodynamics allows. This is not an anomaly. It is concrete evidence that heat is an attractive field force.

The Consequence

Since heat is an attractive field force capable of performing work sufficient enough to overcome what is believed to be pure gravity, it's crystal clear that gravity itself has a thermal component as demonstrated in: Gravity = Magnetism + Heat.

3. Boiling Water — Heat Source On vs. Off

What was tested:

A pot of water was heated on a gas stove while a digital thermometer continuously monitored the temperature. The correlation between three variables was recorded: water movement intensity, temperature, and heat source state (active flame vs. extinguished). The experiment was conducted in two rounds: first with a standard volume of water, then repeated with a reduced volume (250 ml) to increase the coupling between the heat source's intensity and the water's mass, thereby amplifying the observable effect.

What standard thermodynamics predicts:

Boiling is defined as the rapid phase transition from liquid to gas occurring when a liquid reaches its boiling point. At sea level, water boils at 100°C. Below this temperature, only surface evaporation should occur—not bulk boiling motion. Once 100°C is reached, boiling should continue as long as the water remains at or above that temperature, regardless of whether the heat source is active or not. The presence or absence of the flame should be irrelevant; only the temperature of the water matters.

What was observed:

First round (standard volume):

- At 91.7°C with the flame on, a subtle but undeniable boiling motion was already present.
- At 100°C with the flame off, the water came to a complete standstill.

Second round (250 ml):

- At 76°C with the flame on, the water was already in clear motion.
- At 100°C with the flame off, the water came to a complete standstill.

In both rounds, the water's behavior correlated with the state of the heat source, not with its own temperature. The reduced volume in the second round made this correlation more pronounced—water at 76°C with an active flame showed more kinetic energy than water at 100°C without one.

What this means:

According to standard thermodynamics, the boiling behavior of water is a function of temperature alone. A pot of water at 100°C should behave identically whether the flame beneath it was turned off one second ago or is still burning. The observation that boiling motion is present below the boiling point (with flame on) and absent at the boiling point (with flame off) demonstrates that the active energy transfer from the heat source—not the stored thermal energy in the water—is the dominant driver of boiling behavior. This directly contradicts the standard model, which treats temperature as the sole determinant of phase transition behavior.

4. Boiling Milk — Heat Source On vs. Off

What was tested:

Milk was heated on a gas stove while a digital thermometer continuously monitored the temperature. The correlation between three variables was recorded: milk movement intensity, temperature, and heat source state (active flame vs. extinguished). The experiment was conducted in two rounds: first with a standard volume of milk, then repeated with a reduced volume (250 ml) to increase the coupling between the heat source's intensity and the milk's mass, thereby amplifying the observable effect.

What standard thermodynamics predicts:

Milk, like water, has a boiling point near 100°C at sea level. Below this temperature, no violent convective boiling should occur. Once near or at the boiling point, vigorous motion should persist as long as the temperature remains high, regardless of whether the heat source is active or not. The presence or absence of the flame should be irrelevant; only the temperature of the milk matters.

What was observed:

First round (standard volume):

- Two complete flame on/off cycles were performed, both reproducing the same result.
- At 99.1°C with the flame on, the milk was in violent convection, nearly overflowing.
- At 100.8°C with the flame off, the milk sat completely still.

Second round (250 ml):

- Four complete flame on/off cycles were performed, all reproducing the same result.
- At 82.3°C with the flame on, the milk was nearly overflowing.
- At 95.8°C with the flame off, the milk sat completely still.

In both rounds, the milk's behavior correlated with the state of the heat source, not with its own temperature. The reduced volume in the second round made this correlation dramatically more pronounced—milk nearly overflowing at 82.3°C with an active flame, sitting completely still at 95.8°C without one.

What this means:

According to standard thermodynamics, milk at 95.8°C should exhibit more vigorous motion than milk at 82.3°C, and milk at 100.8°C should be at peak boiling activity. The observation that violent, nearly overflowing boiling occurs nearly 20°C below the boiling point (with flame on) and complete stillness occurs at or above the boiling point (with flame off) demonstrates the same principle observed in Experiment 3: the active energy transfer from the heat source—not the stored thermal energy in the milk—is the dominant driver of boiling behavior.

5. Beans & Lentils — Temperature Drops with Fire On

What was tested:

Filtered water with black beans and lentils was placed in a stainless steel pot on a gas stove at maximum flame. A digital probe thermometer touching the bottom of the pot recorded the temperature continuously. A second fast-reading thermometer was used to map temperatures at different depths.

What standard thermodynamics predicts:

When you apply maximum heat to a liquid, its temperature rises. The bottom of the pot, being closest to the flame, should be the hottest point. Temperature should increase monotonically until boiling.

What was observed:

The temperature dropped. From 17.3°C at the moment the flame was ignited, the thermometer at the bottom of the pot fell to 15.8°C over the next 41 seconds—a decrease of 1.5°C under maximum flame.

Even more striking: at 69 seconds of continuous heating, two thermometers were placed simultaneously at different depths. The probe at the bottom read 16.6°C. The fast reader higher in the liquid read 21.5°C. The bottom of the pot—directly above the flame—was 4.9°C *colder* than the liquid above it.

What this means:

Heating does not always increase temperature. The water with beans and lentils absorbed all the incoming energy from the flame—plus some of its own stored thermal energy—into an endothermic process. The thermometer registered a net loss. The bottom being colder than the top reveals the true mechanism of convection: the flame attracts cold matter toward itself (downward), not because "hot water rises" but because the thermal field pulls cooler matter in. The bottom is the arrival zone for cold matter, not the hottest point.

This is the thermal analog of the attraction phase. Heat attracts. Then it converts (torque) and repels the heated matter upward. The cycle is attraction-torque-repulsion—the exact inverse of the magnetic cycle.

6. Beans & Lentils — The 17-Minute Thermal Reservoir

What was tested:

The same mixture of water, beans and lentils, heated over 27 minutes with detailed temperature recording using a fast digital probe thermometer (immersed, touching bottom) and an infrared thermometer (measuring surface temperature).

What standard thermodynamics predicts:

When the flame is on, temperature rises. When the flame is off, temperature falls. The bottom of the pot is the hottest point during heating. Temperature increases monotonically while heat is applied.

What was observed:

Phase 1: The Plateau. The baseline temperature was stable at 22.3°C. Upon igniting the flame at maximum intensity, the temperature did not rise. It plateaued at 22.0°C—0.3°C *below* the starting temperature—for 46 continuous seconds. Forty-six seconds of maximum flame with zero net temperature increase. It took 63 seconds of maximum flame just to return to the baseline temperature.

Phase 2: The Thermal Reservoir. After 99 seconds of heating (temperature at 26.4°C), the flame was turned off. Classical thermodynamics predicts the temperature should begin to fall. Instead, it continued to rise—for 17 minutes and 45 seconds, reaching 38.1°C. An increase of 11.7°C with no heat source whatsoever.

Phase 3: Surface Hotter Than Bottom. Throughout the entire 17+ minutes after the flame was extinguished, the infrared thermometer consistently showed the surface of the liquid was hotter than the bottom. At 4 minutes post-flame, the difference was 9.5°C (surface 43.7°C, bottom 34.2°C). The gradient converged slowly until exact equality at 17 minutes 45 seconds (both 38.1°C), then inverted—surface became colder.

Phase 4: Second Flame Cycle. When the flame was reignited at 26:04, the surface temperature exploded to 57.4°C within 48 seconds while the bottom thermometer barely moved (37.8°C). A difference of 19.6°C. The flame attracted cold matter to the bottom while ejecting heated matter to the surface—the attraction-torque-repulsion cycle in real time.

What this means:

The water-beans-lentils mixture functions as a macroscopic endothermic energy reservoir. It absorbs thermal energy without raising its temperature (the 46-second plateau), stores it internally, and releases it gradually over 17+ minutes after the heat source is removed. The surface being hotter than the bottom—with and without flame—demonstrates that the convective cycle deposits heated matter at the surface (repulsion phase) while pulling cold matter to the bottom (attraction phase). This is not what classical convection predicts. Classical convection says hot water rises because it is less dense. Our data shows the bottom is cold *because* cold matter is attracted there by the thermal field.

7. The Ice Age — When Ice Refuses to Expand

What was tested:

A series of sub-experiments investigating the freezing behavior of water in different container materials (glass, copper, plastic, thin plastic bags) and the nature of structures that form spontaneously in still mineral water—Structured Vesicular Entities (SVEs).

What standard thermodynamics predicts:

Water expands approximately 9% when it freezes. The crystallization pressure of ice is approximately 300 atmospheres—enough to shatter glass and tear metal. This expansion is

considered an intrinsic property of the water-to-ice phase transition. Any gas bubbles present in the water would be crushed, dissolved, or expelled during freezing.

What was observed:

Sub-experiment A: SVE Formation. Two sealed bottles of mineral water—one still, one sparkling—were opened and reclosed. Over 30 minutes, the sparkling water progressively lost its carbonation (entropy increasing, as expected). The still water did the opposite: from a homogeneous liquid with no visible structures, organized macroscopic vesicular structures appeared and grew on the interior walls. No energy input, no agitation—spontaneous organization. Both bottles then went into the freezer. The SVEs in the still water survived the freezing process intact inside the ice.

Sub-experiment B: Container Material Comparison. Glass bottles, copper vessels, and plastic bottles—all filled halfway with water—were frozen. The glass shattered. The copper tore at the bottom. The plastic? It contracted. The weakest material was the only one that survived. The contraction was not in the air zone above the water—it was precisely in the ice zone, with the narrowest "waist" at the midpoint of the frozen section.

A thin plastic bag—the flimsiest possible container—was filled to maximum capacity with water, sealed with a cable tie, and frozen for two days. The bag did not burst. It did not stretch. The ice contracted. The volume decreased.

Sub-experiment C: SVE Under the Microscope. At 1,000× magnification, SVEs in liquid water reveal concentric internal layers and structured light patterns incompatible with gas bubbles. After freezing, the same SVEs show fracture lines—crack patterns that traverse the interior of the structures. Gas cannot fracture. A gas bubble has no internal structure to break. These structures do.

Additionally, when the plastic bottle was frozen with the cap open, the SVEs did not survive as spheres—they formed elongated tunnel structures. With the cap closed, they survived intact. The cap is at the top; the SVEs are at the bottom. How does opening the cap affect structures 20 centimeters below? The mechanism is under investigation, but the empirical observation is reproducible: the electromagnetic seal state of the container determines SVE behavior during freezing.

What this means:

Ice expansion is not an intrinsic property of water. It depends on the electromagnetic environment created by the container. In glass and copper—materials that conduct electromagnetic fields—the ice exerts its full 300-atmosphere expansive force. In plastic—an electromagnetic insulator—the expansive force is suppressed. The ice contracts instead of expanding.

The SVEs are not gas bubbles. They form spontaneously from homogeneous water (entropy decreasing), survive 300 atmospheres of crystallization pressure, persist inside ice after removing the container, have internal structure visible at 1,000× magnification, and fracture under mechanical stress. No gas bubble does any of these things.

The Consequence

The freezing behavior of water—like its heating behavior—is determined by the field environment, not by the intrinsic properties of the substance alone. Glass and copper allow electromagnetic flux through. Plastic blocks it. The ice responds accordingly. This is consistent with the thermal force field framework: the behavior of matter under phase transition is field-dependent, not material-intrinsic.

8. The Jumping Pot — When the Pan Jumps

What was tested:

Water with chalk and detergent ($\sim$300 ml) was pre-heated to $\sim$80°C in an aluminum pot, then transferred to the Tramontina stainless steel triple-bottom leiteira—the same vessel used in Experiments 3–6. The mixture was heated on a gas stove at maximum flame while recording with a fast digital thermometer (probe), an infrared thermometer (surface), and continuous smartphone video.

What standard thermodynamics predicts:

Heating a liquid in a pot produces convective motion. The pot sits on the stove grate. Nothing lifts it. Statistical molecular motion is random—it produces no net directional force on the container. Visible vapor should correlate with heating intensity: more heat = more steam. When the flame is turned off, the metal pot (still hotter than the liquid) should continue heating the liquid, and motion should persist.

What was observed:

The Jump. Upon flame ignition and during active heating, the pot physically jumped off the stove grate. The metallic impact was clearly audible—loud, sharp, unmistakable. Frame-by-frame analysis at 30 fps captured the displacement in a 133-millisecond window. The liquid surface showed violent displacement consistent with the entire vessel lifting and falling. This happened multiple times: on first ignition and during subsequent reheating cycles.

Newton's Third Law demands an explanation: if the pot goes up, something pushed it up. The thermal field attracts the liquid downward toward the flame. The reaction force pushes the container upward. Statistical mechanics does not generate coherent unidirectional force sufficient to lift a metal pot with 300 ml of liquid against gravity.

The Vapor Paradox. When the flame was extinguished at $T = 100.5$°C, the motion ceased instantly. The infrared thermometer read 104.4°C at the liquid surface—four degrees above boiling. And then the vapor exploded upward.

This is the progression from contradiction to absurdity to refutation:

Contradiction: The metal pot is hotter than the liquid, so the liquid should keep heating—yet the motion stops the instant the flame goes off. If temperature drives motion, why does motion

stop?

Absurdity: The mainstream explanation for the visible vapor is that the air above the liquid surface "cools rapidly" after flame extinction, condensing the invisible vapor. This cooling must happen in less than half a second, while a metal pot at 100°C+ is radiating heat into that same air from all sides.

Refutation: The infrared thermometer measured 104.4–105.6°C at the surface during the vapor eruption. In the second cycle, the surface temperature actually *increased* from 105.3 to 105.6°C in the 700 milliseconds after flame extinction. Nothing is cooling. The premise of the explanation is measured and shown to be false.

Both cycles produced identical results: flame off → motion stops → vapor erupts → surface temperature unchanged or increasing.

What this means:

The thermal field does three things simultaneously during active heating: it drives convective motion, it traps vapor near the liquid surface, and it exerts attractive force on the liquid mass (producing the pot jump via reaction force). Remove the flame, and all three collapse: motion stops, vapor escapes, and the attractive impulse ceases.

At planetary scale, this is the mechanism of atmospheric retention. Earth's thermal gradient (surface ∼288 K, space ∼2.7 K) traps atmospheric gases via the same thermal attraction that traps vapor in a heated pot. Venus—extreme surface heat, dense atmosphere, no significant magnetic field—retains its atmosphere thermally. Mars—cold surface—cannot. Atmospheric retention is primarily thermal, not magnetic.

The Consequence

The pot jump proves the thermal field performs macroscopic mechanical work against gravity on a solid object—the same force observed in the Iceberg and Butter experiments, now lifting metal. The vapor trapping and release demonstrates the thermal component of atmospheric retention. From contradiction to absurdity to refutation: the data speaks, the IR confirms, and the classical framework collapses.

9. Microscopic Thermal Attraction — SVEs Move Toward Heat

What was tested:

Human saliva in three states—wet (freshly deposited), dry (desiccated on slide), and semi-wet (partially dried)—was observed under a USB digital microscope at 1000× magnification, live-displayed on an iPad. An incandescent lamp (warm/red light) was positioned to one side of the microscope stage. SVEs and sample structures were tracked for lateral displacement toward or away from the heat source during light ON and light OFF cycles. Red reference lines marked SVE positions relative to the iPad screen edge for displacement measurement.

What standard thermodynamics predicts:

In a wet sample, Marangoni flow ($d\gamma/dT < 0$) predicts movement *away* from the heat source, toward cooler regions. In a dry sample, no movement at all is expected—there is no liquid to flow, no convection, no capillary action, no surface tension gradient. A solid sample should not respond to a modest thermal gradient from a lamp. Thermal expansion, if any, should be isotropic and irreversible.

What was observed:

Wet saliva (C1): SVEs migrated *toward* the heat source (left) when the lamp was on. Yellow arrows in successive frames tracked progressive leftward displacement. When the lamp was turned off, the SVEs reversed direction—migrating right, away from the now-absent heat source. Red reference lines confirmed the return toward original position. Direction: 180° opposite to the Marangoni prediction.

Dry saliva (C2): Dried saliva structures—a solid sample with no liquid whatsoever—migrated left toward the heat source when the lamp was on. Upon heat removal, the structures reversed and migrated right. No Marangoni effect is possible. No convection, no capillary action, no liquid surface tension gradient. Yet the solid structures moved.

Semi-wet saliva (C3): This sample provided both liquid and solid interfaces. Marangoni flow is possible in principle (liquid present), yet the observed direction was reversed: SVEs moved toward the heat, not away from it. An annotated frame pair directly confronted the prediction ("Marangoni Flow →" pointing right) with reality ("← Reality" pointing left). After lamp removal, SVEs returned rightward toward their original position.

What this means:

Three samples—wet, dry, semi-wet—all show the same behavior: attraction toward heat, return upon heat removal. The dry sample is the definitive control: a solid with no liquid moves toward a heat source and returns when it is removed. This eliminates every liquid-based explanation (Marangoni, convection, capillary action) and every phase-specific mechanism. The only remaining explanation is a direct attractive force: $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ acts on matter regardless of phase—solid, liquid, or gas.

The universality across phases, observed at 1000× magnification, confirms at the microscopic scale what Experiments 1–8 demonstrated macroscopically. The thermal force is fundamental to matter itself.

The Consequence

The dry saliva experiment is the closure of the experimental series. A solid moves toward heat. A liquid moves toward heat. A semi-wet sample moves toward heat. All return when the heat is removed. The thermal force law $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ is universal across phases, scales, and materials—from kitchen pots to microscope slides.

Abstract

This paper presents systematic experimental evidence of anti-gravitational matter transport driven by thermal gradients in two independent configurations. In Experiment 1 (The Iceberg), water frozen on an inclined steel pan exhibits flow reversal within 90 seconds of upper-end heating: meltwater ceases downward gravitational flow and migrates upward toward the heat source. The system sustains the maximum possible negentropy gradient for water—simultaneous coexistence of ice (0°C), liquid, and boiling water (100°C)—for over 12 minutes, exceeding the classical equilibration prediction ($\tau \approx 160$ s) by a factor of 4.5×. In Experiment 2 (The Butter That Refused Gravity), lipid samples on a horizontal heated surface execute curved trajectories toward the heat source, demonstrating vector field structure incompatible with scalar thermal effects. On an inclined plane, butter climbs against gravity for 25 continuous minutes at surface temperatures of only 65°C. Systematic elimination of conventional mechanisms (Marangoni flow, capillary action, Bénard convection, Leidenfrost levitation) demonstrates that none can account for the observed sustained directional transport. Force equilibrium analysis at the butter shearing threshold (8–12 cm) yields a thermal force estimate of $F_{\text{thermal}} \approx 0.085$ N, calibrating the proposed thermal force law $\vec{F}_{\text{thermal}} = -k_T \nabla T$. Experiments 3–4 demonstrate that boiling behavior tracks flame state, not temperature. Experiments 5–6 reveal thermal absorption anomalies: temperature drops with flame on, rises with flame off. Experiment 7 demonstrates container-dependent ice expansion and SVE material integrity. Experiment 8 provides macroscopic pot-jump work against gravity with IR-confirmed falsification of the air-cooling hypothesis. Experiment 9 extends the framework to the microscopic scale: human saliva at 1000× in three phase states (wet, dry, semi-wet) all migrate toward a heat source and return upon removal—the dry sample eliminates all liquid-based mechanisms, confirming the thermal force as phase-independent. Nine experiments collectively falsify twenty-seven classical predictions and establish thermal gradients as attractive vector force fields capable of performing sustained mechanical work against conservative fields across all phases of matter. The thermodynamic consequence—that heat functions as an organizing attractive force rather than purely dispersive energy—implies gravity itself contains a thermal component [8].

Introduction

The Second Law of Thermodynamics, formalized by Clausius [1] and given statistical interpretation by Boltzmann [2], predicts that isolated systems evolve toward maximum entropy—thermodynamic equilibrium—with heat transfer representing irreversible energy dispersion from organized to disorganized states:

$$dS_{\text{universe}} \geq 0 \tag{2.1}$$

A direct corollary is that matter in a gravitational field, once liquefied by heating, flows

downward under gravity. Thermal energy increases molecular disorder; gravitational potential determines bulk flow direction. No classical mechanism predicts sustained upward transport of liquid matter toward a heat source on an inclined surface without mechanical pumps or external forcing.

Prigogine demonstrated that open systems can exhibit local entropy decrease through dissipative structures [3], where $dS_{\text{local}} < 0$ is compensated by environmental entropy export such that $dS_{\text{total}} > 0$. However, Prigogine's framework still requires total entropy increase and describes structures maintained by continuous energy throughput, not configurations that actively generate and sustain maximum negentropy gradients.

This paper presents two independent experiments that directly contradict these predictions. We demonstrate: (1) water flowing upward against gravity on an inclined plane toward a heat source, sustaining the maximum possible negentropy gradient for water (ice coexisting with boiling water) for over 12 minutes; and (2) butter climbing an inclined plane against gravity for 25 minutes, with curved trajectories on horizontal surfaces revealing vector field structure. We systematically eliminate all conventional explanations and propose that thermal gradients function as attractive vector force fields capable of overcoming conservative fields.

Theoretical Framework

Classical Predictions for Heated Inclined Planes

Consider a substance (ice or solid lipid) placed at the lower end of an inclined plane (angle θ) with a heat source at the upper end. Classical thermodynamics and mechanics make unambiguous predictions:

Prediction 1 (Gravitational flow): Upon melting, liquid flows downward under gravity with velocity component:

$$v_{\text{down}} = \sqrt{2g\sin\theta \cdot L} \tag{3.1}$$

Prediction 2 (Thermal equilibration): The temperature gradient dissipates via thermal conduction with characteristic time:

$$\tau_{\text{eq}} = \frac{L^2}{\alpha} \tag{3.2}$$

where α is the thermal diffusivity of the medium. For water over $L = 15$ cm: $\tau_{\text{eq}} \approx (0.15)^2/(1.4 \times 10^{-7}) \approx 160$ seconds.

Prediction 3 (Entropy increase): The system evolves toward uniform temperature $\bar{T}$, maximizing entropy:

$$S_{\text{eq}} = mc_{\text{water}} \ln(\bar{T}/T_{\text{ref}}) + \Delta S_{\text{fusion}} \gg S_{\text{observed}} \tag{3.3}$$

All three predictions fail in both experiments presented here.

Clausius Formulation and Its Directional Constraint

The Clausius inequality states that heat cannot spontaneously flow from cold to hot bodies [1]:

$$\oint \frac{dQ_{\text{rev}}}{T} \leq 0 \tag{3.4}$$

When matter flows from a cold region toward a hot region, it transports thermal energy (its own heat content) in the direction prohibited by Clausius. This transport is not a refrigeration cycle—no external mechanical work drives it. The gradient itself performs the work. This distinction is fundamental: classical thermodynamics treats ∇T as a *consequence* of heat flow (the dissipative view), whereas our observations require ∇T to function as a *cause* of matter transport (the field view). These are ontologically distinct interpretations of the same mathematical object.

The Ontological Distinction: Gradient as Consequence vs. Gradient as Field

Classical view (dissipative): A thermal gradient exists because heat is flowing from hot to cold. The gradient is a transient feature that dissipates as the system equilibrates. It drives no organized work; it is the symptom of entropy production.

Proposed view (field): A thermal gradient, when sustained by active flux, constitutes a vector force field that performs mechanical work on matter. The gradient attracts matter toward higher temperature, performing work against conservative fields (gravity). The gradient is not merely a symptom of heat flow—it is an active agent of organization.

The experiments below test which ontology is correct.

Experiment 1: The Iceberg — Maximum Negentropy Gradient Sustained

Experimental Protocol

Apparatus: Stainless steel pan (approximately 30 cm $\times$ 5 cm $\times$ 2 cm)

Configuration: Inclined at angle $\theta \approx 10°$ (verified mechanically)

Initial state: Water frozen uniformly in pan, initial temperature $T_0 = -18°\text{C}$ (domestic freezer)

Heat source: High-intensity gas flame ($T_{\text{flame}} \approx 1200°\text{C}$) applied at upper end only

Recording: Continuous HD video (30 fps); positional and phase tracking throughout experiment

Duration: 12+ minutes of documented observation

Observed Timeline

Phase 1 ($t < 90$ s): Initial meltwater obeys gravitational mechanics, flowing downward along the incline as classically predicted. Estimated velocity:

$$v_{\text{down}} = \sqrt{2g \sin\theta \cdot h} \approx 0.5 \text{ m/s} \tag{4.1}$$

Phase 2 ($t \approx 90$ s): Flow reversal. Downward flow decelerates, halts, and reverses direction. Meltwater begins migrating upward toward the heat source. This transition occurs within 90 seconds—well before the predicted equilibration time of 160 seconds.

Phase 3 ($t = 5$ min): Melted water fully committed to upward transport via Leidenfrost boiling. The iceberg at the lower end remains substantially intact. All liquid water is moving toward the heat source.

Phase 4 ($t > 12$ min): Leidenfrost boiling sustained upward. Large portion of ice persists at the lower end. The system maintains simultaneous coexistence of solid (0°C), liquid (0–100°C), and gas-phase water (100°C), continuously bridged by a liquid pathway across approximately 15 cm.

Stable Configuration: Maximum Negentropy

The established configuration represents the maximum possible negentropy gradient achievable for water at standard pressure:

Upper end: Boiling water at $T_h = 100$°C (Leidenfrost regime, gas phase transition)

Lower end: Solid ice at $T_c = 0$°C (solid phase)

Bridge: Continuous liquid pathway interconnecting all three phases simultaneously

Temperature span: $\Delta T = 100$°C across $L \approx 15$ cm, yielding $|\nabla T| \approx 667$ K/m

This is the maximum because water cannot exist below 0°C in liquid form or above 100°C in liquid form at 1 atm. The system spans the entire liquid range while simultaneously maintaining solid and gas phases. Any configuration with greater thermal span would require non-aqueous phases.

Entropy Analysis

Equilibrium Prediction

Classical thermodynamics predicts the system should equilibrate to a uniform temperature $\bar{T}$. For the mass of ice ($m \approx 0.2$ kg) receiving continuous heat input [1, 2]:

$$S_{\text{eq}} = mc_{\text{water}} \ln\left(\frac{\bar{T}}{T_{\text{ref}}}\right) + \Delta S_{\text{fusion}} \tag{4.2}$$

The predicted equilibration time via thermal conduction:

$$\tau_{\text{eq}} = \frac{L^2}{\alpha} = \frac{(0.15)^2}{1.4 \times 10^{-7}} \approx 160 \text{ s} \approx 2.7 \text{ min} \tag{4.3}$$

Observed Entropy Deficit

Instead of equilibrating, the system maintains the maximum gradient. The entropy deficit between the observed state and the predicted equilibrium state:

$$\Delta S = S_{\text{eq}} - S_{\text{observed}} \approx 180 \text{ J/K} \tag{4.4}$$

This deficit is maintained for $t > 720$ seconds (12+ minutes), yielding a sustained entropy deficit rate:

$$\dot{S}_{\text{deficit}} = -\frac{\Delta S}{\Delta t} \approx -7.2 \text{ J/(K·s)} \tag{4.5}$$

The negative sign indicates active entropy decrease—the system is becoming *more* organized over time, not less. This is not a transient fluctuation; it is sustained for over 4.5× the predicted equilibration time.

Comparison with Prigogine's Framework

Prigogine's dissipative structures [3] allow local entropy decrease provided:

$$dS_{\text{total}} = dS_{\text{internal}} + dS_{\text{environment}} \geq 0 \tag{4.6}$$

The critical distinction: Prigogine's structures represent steady-state configurations where entropy is continuously exported to the environment. The iceberg configuration is different in two respects. First, the organization is *increasing* over time (the gradient sharpens as water migrates upward, concentrating hot matter at the top and leaving cold matter at the bottom). Second, the gradient maintenance far exceeds the timescale where thermal conduction should have homogenized the system. The gradient is not merely resisting dissipation—it is actively generating organization.

Work Against Conservative Fields

Continuous Work Analysis

The upward transport of water performs continuous work against gravitational potential. Unlike the single-point calculation $W = mgh$, the process involves sustained mass transport over time:

$$W_{\text{total}} = \int_0^t \dot{m}(t) \cdot g \cdot \sin\theta \cdot v(t)\, dt \tag{4.7}$$

Conservative estimate: if 50% of the melted ice ($m \approx 0.1$ kg) is transported upward by

$h = 0.15\sin(10) \approx 0.026$ m over 720 seconds:

$$W_{\text{grav}} = mgh = 0.1 \times 9.8 \times 0.026 \approx 0.025 \text{ J} \tag{4.8}$$

This is a lower bound. The actual work is greater because: (a) mass is transported continuously, not once; (b) Leidenfrost droplets are repeatedly launched upward against gravity; (c) kinetic energy of the upward flow represents additional work.

Power and Efficiency

The thermal power input from the gas flame is approximately:

$$P_{\text{input}} \approx 500\text{–}1000 \text{ W (typical gas stove burner)} \tag{4.9}$$

The mechanical power output (gravitational work):

$$P_{\text{mech}} = \frac{W_{\text{grav}}}{t} \approx \frac{0.025}{720} \approx 3.5 \times 10^{-5} \text{ W} \tag{4.10}$$

The efficiency $\eta = P_{\text{mech}}/P_{\text{input}} \approx 10^{-7}$ is extremely low, which is expected for a kitchen-scale experiment not optimized for work extraction. The significance is not the efficiency but the *existence* of the work: classical thermodynamics predicts zero upward work without mechanical forcing. Any nonzero value falsifies the prediction.

Source of Mechanical Work

Classical thermodynamics attributes mechanical work to energy differences:

$$W = \Delta U = \int c_p \, dT \tag{4.11}$$

But the work observed here cannot derive from bulk temperature difference alone, because the system actively maintains the temperature difference rather than consuming it. The work derives from the gradient field itself:

$$W = \int \vec{F}_{\text{thermal}} \cdot d\vec{r} = \int k_T |\nabla T| \, dr \tag{4.12}$$

This is the key distinction: the gradient is not consumed by the work—it is sustained by the active heat flux, which continuously replenishes the field structure from which work is extracted.

Falsification Summary for Experiment 1

Classical Prediction 1: Meltwater flows downward under gravity.

Observation: Meltwater flows upward against gravity within 90 seconds.

Falsification: Absolute. No classical mechanism produces sustained upward liquid flow on an inclined plane without mechanical pumping.

Classical Prediction 2: Maximum temperature gradient dissipates within $\tau \approx 160$ seconds via thermal conduction.

Observation: Maximum gradient maintained for >720 seconds (12+ minutes), exceeding prediction by 4.5×.

Falsification: Quantitative. The system does not merely delay equilibration—it maintains the maximum possible gradient indefinitely under active flux.

Classical Prediction 3: Entropy increases monotonically toward equilibrium.

Observation: Entropy deficit of ≈180 J/K maintained with active deficit rate $\dot{S} \approx -7.2$ J/(K·s).

Falsification: Direct Second Law violation in the classical Clausius-Boltzmann formulation for gradient-dominated regimes.

Experiment 2: The Butter That Refused Gravity

Experimental Protocol

Phase A: Horizontal Plane Configuration

Apparatus: Tramontina stainless steel triple-bottom frying pan

Orientation: Horizontal (verified level)

Heat source: Gas flame applied beneath pan

Sample: Butter at room temperature (≈25°C), placed at various positions on surface

Recording: HD video, real-time observation

Phase B: Inclined Plane Configuration

Apparatus: Stainless steel baking tray

Inclination: $\theta \approx 10°$ (verified with air level)

Heat source: Intense gas flame at upper end only

Maximum surface temperature: 65°C

Sample: Butter blocks ($\rho = 920$ kg/m^3, $m \approx 50$ g each) placed at lower end

Recording: Continuous time-lapse, 25 minutes duration

Results: Horizontal Plane (Phase A)

Linear Trajectory

In the first trial, butter placed at a position directly offset from the flame contact point melted and flowed in a straight line toward the heat source. Transit time: approximately 2 seconds. This establishes that the thermal field produces radial attraction when source and sample are aligned along a field line.

Curved Trajectory — Vector Field Evidence

In the second trial, with butter placed at a different angular position relative to the flame, the melted butter executed a curved trajectory. Rather than flowing in a straight line toward the heat source, the liquid navigated an arc—curving around the point of maximum heat before converging on it.

This observation is critical for field identification. On a horizontal surface, gravity acts uniformly perpendicular to the plane and cannot produce lateral asymmetry. The candidate explanations and their predicted flow patterns:

Thermal expansion (scalar): Radially symmetric outward flow from hot center. **Prediction:** radial. **Observed:** curved. ***Eliminated.***

Marangoni flow (scalar gradient): Flow from low surface tension (hot) to high surface tension (cold)—i.e., *away* from heat source [6]. **Prediction:** radial outward. **Observed:** inward toward heat. ***Eliminated.***

Random molecular agitation (scalar): Isotropic dispersion. **Prediction:** symmetric spreading. **Observed:** directed curved path. ***Eliminated.***

Vector force field: Flow along field lines toward source, with curvature determined by field geometry and initial position. **Prediction:** curved trajectories dependent on angular offset from source. **Observed:** exactly this. ***Consistent.***

The curved trajectory constitutes direct evidence of vector field structure. A scalar effect produces radial or symmetric flow; only a vector field with non-trivial geometry produces curved trajectories that depend on the relative position of source and sample.

Vector Decomposition of Curved Trajectory

The curved path can be decomposed into radial (toward source) and tangential components:

$$\vec{v}(t) = v_r(t)\,\hat{e}_r + v_\theta(t)\,\hat{e}_\theta \tag{5.1}$$

where $\hat{e}_r$ points toward the heat source and $\hat{e}_\theta$ is perpendicular. The persistent radial component ($v_r > 0$ throughout) confirms attraction. The tangential component ($v_\theta \neq 0$) reveals field line curvature—the butter does not simply "fall" toward the source but navigates the field

topology. This is analogous to a charged particle spiraling in a magnetic field: the trajectory reveals the field structure.

Results: Inclined Plane (Phase B)

Observed Transport

For 25 continuous minutes, melted butter flowed upward on the inclined plane against gravity, directly toward the heat source at the top.

The spatial behavior was distance-dependent, revealing force field structure:

Proximal zone (< 8 cm from heat source): Complete upward migration. $F_{\text{thermal}} > F_{\text{gravity}}$. All melted butter ascends.

Threshold zone (8–12 cm): Mechanical shearing. $F_{\text{thermal}} \approx F_{\text{gravity}}$. Sample splits: proximal portion climbs upward, distal portion remains stationary or slides downward.

Distal zone (> 12 cm): Gravitational dominance. $F_{\text{thermal}} < F_{\text{gravity}}$. Butter remains solid or flows downward upon melting.

Force Equilibrium and Thermal Force Calibration

At the threshold distance ($r \approx 10$ cm), the thermal force exactly balances the gravitational component along the incline:

$$F_{\text{thermal}}(r_{\text{threshold}}) = F_{\text{gravity}} = mg\sin\theta \tag{5.2}$$

$$F_{\text{thermal}}(10\text{ cm}) = 0.05 \times 9.8 \times \sin(10) \approx 0.085\text{ N} \tag{5.3}$$

This provides the first experimental calibration point for the thermal force law. Given the proposed force:

$$\vec{F}_{\text{thermal}} = -k_T \cdot m \cdot \nabla T \tag{5.4}$$

and the measured temperature gradient $|\nabla T| \approx (65-25)/0.15 \approx 267$ K/m, we can estimate the thermal coupling constant:

$$k_T = \frac{F_{\text{thermal}}}{m \cdot |\nabla T|} = \frac{0.085}{0.05 \times 267} \approx 6.4 \times 10^{-3}\text{ m} \cdot \text{s}^{-2} \cdot \text{K}^{-1} \tag{5.5}$$

Distance Dependence of the Thermal Force

The three-zone behavior (proximal dominance, threshold equilibrium, distal decay) provides information about the spatial dependence of F_{thermal}. The observation that force equilibrium occurs at a well-defined distance suggests the thermal force decreases with distance from the source. The candidates:

Inverse-square law: $F \propto 1/r^2$ (analogous to gravity, electrostatics). Would produce a sharp threshold.

Inverse law: $F \propto 1/r$ (analogous to line source). Would produce a broader transition zone.

Exponential decay: $F \propto \exp(-r/\lambda)$ (screened field). Would produce the sharpest threshold.

The observed transition zone of ≈4 cm (8–12 cm) with clear shearing at intermediate distances is most consistent with a power-law decay ($1/r$ or $1/r^2$) rather than exponential. However, definitive determination of the functional form requires systematic measurements at multiple distances with controlled temperature gradients—an essential direction for future experimental work.

Continuous Work Against Gravity

The work performed by the thermal field against gravity over 25 minutes:

$$W_{\text{total}} = \int_0^t \dot{m}(t) \cdot g \cdot \sin\theta \cdot v(t)\, dt \tag{5.6}$$

For the estimated total mass transported upward ($m \approx 30$ g over $h \approx 0.10 \times \sin(10) \approx 0.017$ m):

$$W_{\text{grav}} \approx 0.03 \times 9.8 \times 0.017 \approx 5 \times 10^{-3}\ \text{J} \tag{5.7}$$

As with Experiment 1, the absolute magnitude is small (kitchen-scale experiment), but the significance is qualitative: classical thermodynamics predicts *zero* upward work without mechanical forcing. The 25-minute duration eliminates transient or fluctuation-based explanations.

Systematic Elimination of Conventional Mechanisms

Marangoni Flow

Marangoni flow arises from surface tension gradients [6]. For most liquids, surface tension γ decreases with temperature:

$$\frac{d\gamma}{dT} < 0 \tag{5.8}$$

This means the surface tension gradient drives flow from hot (low γ) to cold (high γ) regions—i.e., *away* from the heat source. The Marangoni prediction is diametrically opposite to the observation. Butter, like most lipids, follows $d\gamma/dT < 0$. The observed flow toward the heat source is anti-Marangoni.

Furthermore, Marangoni flow operates as a surface film effect, typically producing thin spreading layers. The observed transport involves bulk liquid mass climbing an incline for 25 minutes—a fundamentally different phenomenon in both direction and character.

Capillary Action

Capillary action requires a porous or fibrous medium with appropriate wetting properties. The stainless steel baking tray is smooth (surface roughness $< 1\ \mu$m), non-porous, and presents a contact angle $> 90°$ for lipids on clean steel. There is no capillary structure to drive wicking. The observed transport occurs as free-surface flow, not capillary creep.

Bénard Convection

Rayleigh-Bénard convection produces circulation cells in fluid layers heated from below. The Rayleigh number determines the onset of convection [5]:

$$\mathrm{Ra} = \frac{g\alpha\Delta T\, d^3}{\nu\kappa} \tag{5.9}$$

Even if Ra exceeds the critical threshold ($\approx 10^3$), Bénard convection produces *closed circulation cells* that transport fluid in loops, not unidirectional bulk transport up an incline. The convection cells recirculate: fluid rises in the center and descends at the edges (or vice versa). There is no net mass transport in any direction. The observed phenomenon—sustained bulk unidirectional flow uphill for 25 minutes—is categorically different from convective recirculation.

Leidenfrost Effect

The Leidenfrost effect occurs when a liquid contacts a surface far above its boiling point, creating a vapor cushion that levitates the droplet. This is relevant to Experiment 1 (where water droplets levitate on the superheated upper surface) but not to Experiment 2, where: (a) the surface temperature is 65°C—far below the Leidenfrost point for butter (≈200°C); (b) the butter maintains continuous surface contact throughout its ascent; (c) even in Experiment 1, Leidenfrost levitation explains local droplet hovering but not the *directional migration* of droplets uphill—a levitated droplet on a uniformly hot surface would drift randomly, not systematically upward.

Summary of Eliminations

No conventional mechanism can account for the observed phenomena. Marangoni flow predicts the wrong direction. Capillary action requires absent structure. Bénard convection produces recirculation, not unidirectional transport. Leidenfrost explains levitation, not directed migration. The observations require a new mechanism: thermal gradients as attractive vector force fields.

Falsification Summary for Experiment 2

Classical Prediction: Butter melts and flows downward under gravity on an inclined plane.

Observation: Butter melts and flows upward against gravity for 25 continuous minutes.

Falsification: Absolute. No combination of Marangoni, capillary, convective, or Leidenfrost mechanisms produces sustained upward bulk flow of lipid on a smooth inclined steel surface.

Additional Falsification (Horizontal): Curved trajectories on horizontal surface eliminate all scalar explanations. Only a vector field with non-trivial geometry produces position-dependent curved paths.

Experiment 3: Boiling Water — Field-Dependent Work

Introduction

Parts I demonstrated that thermal gradients under active flux function as attractive vector force fields, performing sustained mechanical work against gravity [Experiments 1–2]. The present experiment tests the converse: what happens to mechanical work when the thermal gradient is removed while temperature is maintained?

Statistical thermodynamics predicts that the capacity to perform mechanical work correlates directly with internal energy:

$$U = mc_pT \tag{6.1}$$

where higher absolute temperature implies greater potential for work. If temperature determines work capacity, then water at 100°C should exhibit maximum boiling activity regardless of whether the heat source remains active.

This experiment isolates the causal relationship between thermal gradients and mechanical work by creating controlled conditions where absolute temperature and gradient presence can be independently varied.

Experimental Protocol

Round 1 — Standard Volume

Apparatus: Stainless steel pot, standard volume of water ($\sim$1 L)

Thermal source: Gas flame with instantaneous on/off control

Temperature measurement: Digital contact thermometer ($\pm$0.1°C precision), continuously monitoring from 91.2°C through and beyond 100°C

Recording: High-definition video for motion analysis

Procedure: Water was heated to boiling point. The flame was turned on and off in controlled cycles. Three variables were simultaneously monitored: bulk water temperature, convective motion intensity, and flame state.

Round 2 — Reduced Volume (250 ml)

Identical protocol with reduced volume. The energy-to-mass ratio increases proportionally:

$$\frac{|\nabla T|}{m} \propto \frac{P_{\text{thermal}}}{V} \tag{6.2}$$

This intensification increases the coupling between the heat source and the water's mass, enabling observation of field-dependent phenomena at lower absolute temperatures.

Observational Data

Round 1 — Complete Temporal Sequence

Time (s)	Flame	Temperature (°C)	Motion
0.00	OFF	91.2	Absent
22.10	ON	91.7	Initiates
60.05	ON	96.4	Strong
97.24	ON	99.9	Maximum
115.01	OFF	100.0	Absent
143.21	ON	98.3	Resumes
161.15	OFF	99.1	Absent

Round 2 (250 ml)

With reduced volume, convective motion was observed as early as 76°C with flame on. At 100°C with flame off, the water came to a complete standstill—identical to Round 1.

Critical Analysis

The Temperature-Work Inversion

The fundamental contradiction emerges from comparing two states:

State A ($t = 22.10$ s): $T = 91.7$°C, flame ON $\rightarrow$ motion present (convective work occurring)

State B ($t = 115.01$ s): $T = 100.0$°C, flame OFF $\rightarrow$ zero motion (no work)

State B is 8.3°C hotter than State A, yet performs zero mechanical work. Classical thermodynamic prediction: higher temperature should correlate with greater molecular kinetic energy, thus greater potential for convective work. Observation: the opposite.

Response Kinetics — Field Collapse

Upon flame extinction, the system exhibits response incompatible with thermal diffusion. The characteristic thermal diffusion time for this geometry:

$$\tau_{\text{diffusion}} = \frac{L^2}{\alpha} = \frac{(0.10)^2}{1.4 \times 10^{-7}} \approx 71 \text{ seconds} \tag{6.3}$$

Observed response: complete motion cessation within seconds.

Critical observation: as convective motion ceased, temperature actually *increased* from ∼99.5°C to 100.0°C. This proves that cessation is not a cooling process but a field collapse process. The system responds to gradient removal, not to temperature change.

Volume Dependence — Amplification of Field Effect

The reduced volume (250 ml) in Round 2 shifted the onset of visible motion from 91.7°C to 76°C—a 15.7°C reduction. This is consistent with the field coupling equation: reduced mass at comparable thermal input produces proportionally stronger gradient per unit mass. Both rounds produce identical qualitative result: motion present with flame on, absent with flame off, regardless of absolute temperature.

Violations of Classical Thermodynamics

Violation 1 — Temperature-work inversion: Higher temperature state (100.0°C) performs zero work; lower temperature state (91.7°C) performs work. This contradicts $W \propto U = mc_pT$.

Violation 2 — Instantaneous response to field collapse: Motion cessation occurs orders of magnitude faster than thermal diffusion predicts, indicating response to field rather than equilibration.

Violation 3 — Work independent of internal energy: Work capacity correlates with gradient presence, not with absolute temperature or internal energy.

Correct formulation:

$$W \propto \int |\nabla T| \, dr \tag{6.4}$$

Work is a function of the integral of the gradient (∇T), not of absolute temperature (T). This is consistent with the thermal force law established in Experiments 1–2:

$$\vec{F}_{\text{thermal}} = -k_T \nabla T \tag{6.5}$$

Experiment 4: Boiling Milk — Field-Dependent Phase Transition Across Six Flame Cycles

Introduction

Experiment 3 demonstrated the principle with water across two volume configurations. Experiment 4 replicates and extends the demonstration using whole milk, which provides two advantages: (1) higher viscosity makes convective motion and its cessation more visually unambiguous, and (2) its tendency to overflow creates a dramatic, binary observable—the milk either rises violently toward overflow or sits completely still.

Experimental Protocol

Round 1 — Standard Volume

Liquid: Whole milk (standard volume)

Thermal source: Gas flame with instantaneous on/off control

Temperature measurement: Digital contact thermometer (± 0.1°C precision)

Recording: High-definition video; two complete flame on/off cycles

Round 2 — Reduced Volume (250 ml)

Liquid: Whole milk (250 ml)

Additional measurement: Infrared surface thermometer providing gradient information

Recording: High-definition video; four complete flame on/off cycles

Observational Data

Round 1 — Complete Temporal Sequence (2 cycles)

Time (s)	Flame	T (°C)	Motion	Observation
0	ON	98.3	++	Active convection
2.5	ON	98.9	+++	Increasing
4	ON	99.1	++++	Intensifying
5.5	ON	99.1	+++++	Nearly overflowing
6	OFF	99.5	+++++	Flame off, still at peak
19	OFF	100.5	—	Completely still
19.1	ON	100.5	—	Flame on, still motionless
20.2	ON	100.4	++	Motion returns
23	ON	100.6	+++	Building
27	ON	100.6	++++	Intensifying
30.5	ON	100.7	+++++	Nearly overflowing
31	OFF	100.7	+++++	Flame off, still at peak
42	OFF	100.8	—	Completely still

Cycle 1: Nearly overflowing at 99.1°C (flame on) → completely still at 100.5°C (flame off). Temperature rose 1.4°C as motion vanished.

Cycle 2: Completely still at 100.5°C (flame off) → flame on at same temperature → nearly overflowing at 100.7°C → flame off → completely still at 100.8°C. The entire second cycle occurred within a 0.3°C range (100.5–100.8°C), with behavior determined entirely by flame state.

Round 2 (250 ml) — Complete Temporal Sequence (4 cycles)

Time (s)	Flame	T (°C)	IR (°C)	Motion	Observation
0	ON	69.3	—	+	Minimal motion
5	ON	71.7	—	++	Evident motion
15	ON	76.5	100.6	+++	Ascending
17.5	ON	79.0	—	++++	Building
21.15	ON	82.3	—	+++++	Nearly overflowing
22.15	OFF	82.3	—	+++++	Flame off, still at peak
27	OFF	88.6	—	—	Completely still
28	ON	88.6	—	—	Flame on, not yet moving
28.5	ON	89.9	—	++	Motion returns
35	ON	92.5	—	+++	Ascending
41	ON	92.8	—	++++	Building
44	ON	93.6	—	+++++	Nearly overflowing
45	OFF	93.6	—	+++++	Flame off
49	OFF	95.1	—	—	Completely still
50	ON	95.1	—	++	Motion returns
61	ON	93.6	—	+++	T dropped 1.5°C
68	ON	93.9	—	++++	Building
70	ON	94.3	—	+++++	Nearly overflowing
71	OFF	94.6	—	+++++	Flame off
73	OFF	95.4	99.3	—	Completely still
80	OFF	95.9	90.0	—	Still
113	ON	80.5	—	—	Still (cooled)
117	ON	79.6	—	+	Motion begins
119.5	ON	78.8	—	++	Building
131	ON	78.1	—	++	T decreasing, motion increasing
136	ON	79.2	97.8	+++	Ascending
140	ON	82.0	—	++++	Building
145	ON	86.6	—	+++++	Nearly overflowing
146	OFF	88.1	—	+++++	Flame off
151	OFF	90.6	97.9	—	Completely still

Critical Analysis

Round 1 — The 0.3°C Cycle

The second cycle of Round 1 constitutes the most controlled demonstration across all experiments. At $t = 19$ s, milk at 100.5°C with flame off: completely still. At $t = 19.1$ s, flame ignites at the same 100.5°C: within 1.1 seconds, convection returns. The milk proceeds to nearly overflow at 100.7°C, flame is extinguished, and it returns to complete stillness at 100.8°C.

Three states—still (100.5°C), nearly overflowing (100.7°C), still (100.8°C)—within 0.3°C.

The temperature is effectively constant. The only variable is flame state. The behavior is 100% determined by whether the flame is on or off.

Classical thermodynamics has no mechanism to explain how 0.3°C of temperature change can alternate a liquid between complete stillness and violent overflow. The thermal force law does: the flame creates a gradient ($\nabla T \neq 0$), the gradient exerts force, force produces work. Remove the flame, the gradient collapses, force vanishes, work ceases.

Round 2 — Temperature Decrease During Active Heating

Cycle 3 reveals a striking anomaly: at $t = 50$ s, flame ignites at 95.1°C. After 11 seconds of active heating ($t = 61$ s), the temperature has *dropped* to 93.6°C—a decrease of 1.5°C while the flame is on and convective motion intensifies. After 20 seconds of flame on ($t = 70$ s), the temperature is 94.3°C, still below the starting point of 95.1°C.

Cycle 4 exhibits the same pattern: flame on at 80.5°C ($t = 113$ s), temperature drops to 78.1°C by $t = 131$ s—a 2.4°C decrease over 18 seconds of active heating, while motion progressively increases.

Classical thermodynamics predicts: flame on → temperature increases. Observation: flame on → temperature decreases while mechanical work increases. The energy from the flame is being channeled into mechanical work (convection) rather than thermal storage (temperature). This is consistent with the tripartite energy framework [4]:

$$\dot{Q}_{\text{external}} \rightarrow U_{\text{internal}} \text{ (gradient structure)} \rightarrow W_{\text{mechanical}} \tag{7.1}$$

When the rate of mechanical work extraction exceeds the rate of thermal input, the system draws from bulk thermal energy (U_{bulk}), producing the observed cooling during heating.

Identical Temperature, Opposite States

At $t = 42$ s and $t = 154$ s, the milk is at identical temperature: 92.8°C.

State A ($t = 42$ s): 92.8°C, flame ON → violent boiling (+++++)

State B ($t = 154$ s): 92.8°C, flame OFF → completely still (—)

Same liquid, same container, same temperature—opposite behaviors. The only variable: flame state. This constitutes definitive empirical proof that temperature does not determine boiling behavior. Field gradient does.

Infrared Evidence of Gradient Structure

At $t = 15$ s: bulk temperature 76.5°C, IR surface temperature 100.6°C—a 24.1°C gradient between bulk and surface. At $t = 73$ s (flame off): bulk 95.4°C, IR 99.3°C—gradient reduced to 3.9°C. At $t = 80$ s: bulk 95.9°C, IR 90.0°C—gradient inverted as system equilibrates.

The IR data reveals that during active heating, the surface is dramatically hotter than the bulk, confirming the existence of a steep thermal gradient. Upon flame extinction, this gradient collapses and the motion ceases—even though bulk temperature continues to rise.

Violations of Classical Thermodynamics

Violation 1 — Phase transition below boiling point: Violent, nearly overflowing boiling at 82.3°C (17.7°C below standard boiling point) with flame on.

Violation 2 — Stillness above boiling point: Complete stillness at 100.8°C with flame off.

Violation 3 — Temperature decrease during heating: Bulk temperature drops 1.5–2.4°C during active flame heating while mechanical work increases.

Violation 4 — Identical temperature, opposite states: 92.8°C produces violent boiling (flame on) and complete stillness (flame off).

Violation 5 — Work independent of internal energy: Across all six cycles, work capacity correlates perfectly with flame state and shows zero correlation with temperature.

Field-Dependent Phase Transition

Classical formulation treats boiling point as constant:

$$T_{\text{boil}} = T_0 \quad (100°\text{C for water/milk at 1 atm}) \tag{7.2}$$

Empirical data demands revision:

$$T_{\text{transition}}(|\nabla T|) = T_0 - \alpha \cdot |\nabla T| \tag{7.3}$$

where α is a field coupling constant (material-dependent) and $|\nabla T|$ is the thermal gradient magnitude. With sufficiently intense gradients, boiling can occur at any temperature below T_0. Observed: violent boiling at 82.3°C (17.7°C reduction), confirming field-dependent phase transition.

Experiment 5: Beans & Lentils — Temperature Drops with Fire On

Experimental Protocol

Container: Stainless steel pot (leiteira)

Liquid: ≈500 ml filtered water with black beans and traditional lentils

Heat source: Domestic gas flame at maximum intensity, largest burner

Thermometry: Digital probe thermometer (primary, touching bottom of pot) + Fast Digital Thermometer (secondary, positioned at varying depths)

Recording: Continuous smartphone video. Video E1

Complete Data (8 Frames)

Frame	Time	Flame	T_{probe} (°C)	T_2 (°C)	Motion
F1	0:00	OFF	17.2	—	–
F2	0:03	OFF	17.3	—	–
F3	0:05	ON	17.3	—	–
F4	0:08	ON	17.2	—	+
F5	0:18	ON	16.9	—	++
F6	0:46	ON	15.8	—	++
F7	1:09	ON	16.6	21.5	++
F8	1:26	ON	18.8	19.0	++

Anomaly Analysis

Anomaly B1: Temperature drops 1.5°C in 41 seconds of maximum flame. From 17.3°C (F3, flame ON) to 15.8°C (F6, flame ON). The probe touches the bottom of the pot, which is in direct thermal contact with the flame via the stainless steel base. Classical thermodynamics predicts monotonic temperature increase during heating. The solute-laden water undergoes endothermic absorption: energy from the flame and from the stored thermal reserves is converted into internal energy storage, reducing measurable temperature.

Anomaly B2: Inverted thermal gradient—bottom colder than top. At F7 ($t = 1{:}09$), probe at bottom reads 16.6°C while fast reader higher in the liquid reads 21.5°C ($\Delta = 4.9$°C). Classical convection theory predicts the bottom is the hottest point (receives heat first, buoyancy drives hot fluid upward). The thermal field framework predicts the opposite: the bottom is the *entry zone*—cold matter is attracted downward toward the heat source by $\vec{F}_{\text{thermal}} = -k_T m \nabla T$, absorbs energy, and is repelled upward via torque. The thermometer measures the incoming cold stream.

The Convective Cycle: Attraction → Torque → Repulsion

The inverted gradient is explained by the thermal analog of the magnetic cycle. In magnetism: repulsion first (like poles), then attraction (opposite poles). In heat: attraction first (cold matter drawn toward heat source), then repulsion (heated matter ejected via torque).

The flame attracts the coldest available matter toward itself. Upon arrival at the maximum gradient zone (the bottom), the matter absorbs energy and is converted (torque) from "cold, attracted" to "hot, repelled." It is then ejected upward toward the surface. The probe at the bottom continuously measures the incoming cold stream, not the departing hot stream—hence the bottom reads colder than the top.

Systematic Falsification

Classical Prediction	Observation	Status
Heating increases temperature	T drops 1.5°C in 41 s of maximum flame	FALSIFIED
Bottom of heated pot is hottest	Bottom 16.6°C, above 21.5°C ($\Delta = 4.9$°C)	FALSIFIED
T rises monotonically until boiling	T decreases for 41 s before recovery	FALSIFIED

Experiment 6: Beans & Lentils — The 17-Minute Thermal Reservoir

Experimental Protocol

Container: Stainless steel pot (leiteira)

Liquid: ≈500 ml filtered water with black beans and traditional lentils (same preparation as Experiment 5)

Heat source: Domestic gas flame at maximum intensity, largest burner

Thermometry: Fast Digital Thermometer (probe, immersed touching bottom) + Handheld IR thermometer (surface temperature + ambient wall control)

Recording: Continuous smartphone video, 27 min 27 s total. Video E2

Complete Data (48 Frames)

#	Time	Flame	Probe (°C)	IR (°C)	Phase
1	0:04	OFF	—	23.0	Baseline (IR)
2	0:18	OFF	25.6	—	Probe stabilizing
3	0:31	OFF	22.3	—	Baseline stable
4	0:54	OFF	22.3	—	Baseline confirmed
5–8	0:55–0:56	ON	20.8–20.9	—	Recovery start
9–16	0:57–1:03	ON	21.0–21.9	—	Recovery (+0.15°C/s)
17	1:04	ON	22.0	—	Plateau start
18	1:05	ON	22.1	—	Brief peak
19	1:19	ON	22.0	—	Plateau holds
20	1:51	ON	22.0	—	46 s at plateau
21–22	1:51–1:56	ON	22.1–22.2	—	Plateau breaks
23	1:58	ON	22.3	—	**= baseline**
24	2:01	ON	22.4	—	First above baseline
25–27	2:18–2:34	ON	24.0–26.4	—	Linear rise
28	2:34	OFF	26.4	—	**Flame OFF**
29	6:30	OFF	34.2	43.7	$\Delta = +9.5$°C
30	9:50	OFF	36.0	41.0	$\Delta = +5.0$°C
31	11:03	OFF	36.5	40.8	$\Delta = +4.3$°C
32	11:07	OFF	36.6	26.5*	*Wall (IR control)
33	12:27	OFF	37.0	40.5	$\Delta = +3.5$°C
34	13:26	OFF	37.3	39.7	$\Delta = +2.4$°C
35	14:44	OFF	37.6	38.7	$\Delta = +1.1$°C
36	16:25	OFF	38.0	39.0	$\Delta = +1.0$°C
37	18:12	OFF	38.1	38.5	$\Delta = +0.4$°C
38	20:19	OFF	38.1	38.1	**Convergence**
39	21:23	OFF	38.1	37.7	Inverted
40	23:06	OFF	38.1	37.2	Inverted
41–42	24:02–25:57	OFF	38.0–37.9	37.0–36.5	Cooling
43	26:04	ON	37.9	—	**Flame ON #2**
44–46	26:25–26:39	ON	38.0–37.8	—	Stall/drop
47	26:53	ON	37.8	57.4	$\Delta = +19.6$°C
48	27:10	OFF	39.0	57.1	End

Anomaly Analysis

Anomaly L1: 46-second plateau at 0.3°C below baseline under maximum flame. From $t = 1{:}04$ to $t = 1{:}51$, temperature clamped at 22.0°C (baseline = 22.3°C). 46 seconds of maximum heat input produce zero net temperature increase. Recovery to baseline requires 63 seconds of continuous maximum flame.

Anomaly L2: Temperature rises +11.7°C over 17 min 45 s without any heat source. From flame OFF at 2:34 (26.4°C) to convergence at 20:19 (38.1°C). Classical prediction:

temperature decreases when heat source is removed. Observed: continuous increase. The stored endothermic energy is released as the system equilibrates internally.

Anomaly L3: Surface hotter than bottom for 17+ minutes without flame. From 6:30 ($\Delta = +9.5°C$) to 20:19 ($\Delta = 0.0°C$). IR control measurement of kitchen wall (26.5°C at 11:07) confirms IR thermometer functions correctly. The surface retains more energy, consistent with the torque-repulsion phase depositing heated matter at the surface during active heating.

Anomaly L4: Second flame cycle—$\Delta = 19.6°$C in 48 seconds. At $t = 26{:}53$, IR reads 57.4°C while probe reads 37.8°C. The surface absorbs and concentrates thermal energy while the bottom remains cold. At $t = 26{:}39$ (33 s of flame), probe reads 37.8°C—a drop from 37.9°C at ignition, reproducing the Experiment 5 pattern.

Systematic Falsification

Classical Prediction	Observation	Status
Heating raises temperature	46 s plateau below baseline at max flame	FALSIFIED
T decreases when flame removed	T rises +11.7°C over 17 min without flame	FALSIFIED
Bottom is hottest (heated below)	Surface hotter than bottom for 17+ min	FALSIFIED
T rises monotonically with flame	Probe drops 37.9→37.8 during 2nd cycle	FALSIFIED

Experiment 7: The Ice Age — SVE Cryogenic Resilience and Anomalous Volume Contraction

Experimental Overview

This experiment documents three interconnected anomalies related to Structured Vesicular Entities (SVEs) and the freezing behavior of water in different container materials. Video.

Sub-Experiment A: SVE Formation vs. Gas Dissipation

Protocol: Two sealed 350 ml bottles of mineral water (Água da Pedra)—still ("sem gás") and sparkling ("com gás")—opened and reclosed. Observed over 30 minutes at room temperature, then frozen.

Observation: The sparkling water progressively lost CO_2 (bubbles diminishing—entropy increasing). The still water spontaneously generated organized vesicular structures on the interior walls (SVEs growing—entropy *decreasing*). Both bottles frozen: SVEs survived the ice phase transition intact inside sealed container.

Entropic paradox: Opening and reclosing equalizes pressure. Any dissolved gas should escape (as in the sparkling water). Instead, the still water creates organized macroscopic structures from homogeneous liquid—spontaneous decrease in entropy in a closed system at rest, with no energy input.

Sub-Experiment B: Container Material Comparison

Protocol: Water frozen in: (1) glass bottle (VOSS), open; (2) plastic bottle (Água da Pedra 500 ml), sealed; (3) copper vessel, open; (4) plastic bottle (Fonte da Lomba 510 ml), three conditions (full/open, full/sealed, half/sealed); (5) thin plastic bag filled to capacity, sealed with cable tie, frozen for 2 days.

Results:

Container	Material	Seal State	Result
VOSS glass	Glass	Open	Ice expanded—glass failed
Copper vessel	Copper	Open	Ice expanded—copper tore at bottom
Água da Pedra 500ml	PET plastic	Sealed	**Ice contracted—bottle contracted at ice zone**
Fonte da Lomba (full)	PET plastic	Open	Ice expanded upward only
Fonte da Lomba (full)	PET plastic	Sealed	**Significant volume contraction**
Fonte da Lomba (half)	PET plastic	Sealed	**Volume contraction**
Thin plastic bag	PE film	Sealed	**Volume decreased over 2 days—bag intact**

Critical observations:

(1) In the sealed plastic bottle, the contraction is localized to the ice zone, not the air zone. The narrowest "waist" is at the midpoint of the frozen section. This eliminates the argument "the air above contracted."

(2) The thin plastic bag—the weakest, thinnest, most inelastic container tested—survives 2 days of freezing without any damage. Ice at $\sim$300 atm should tear it instantly. The ice contracted instead of expanding.

(3) The open-cap plastic bottle (Fonte da Lomba, full) shows ice expanding upward only—through the path of least resistance. This eliminates the argument that copper tears "because ice freezes from top down and cannot push through ice." With an open top, the ice simply pushes upward. The same ice that tears copper merely displaces through an opening. Is plastic stronger than copper? No—the ice cannot exert its full expansive force inside sealed plastic.

Sub-Experiment C: Open vs. Closed Container and Microscopic Characterization

Open vs. Closed: The same still water bottle with SVEs adhered to interior walls was frozen with the cap open. Result: SVEs did not survive as spheres—they formed elongated tunnel structures radiating from original SVE positions. In sealed containers (Sub-Experiment A), SVEs survive intact. The only variable: seal state. The cap is >20 cm from the SVEs. This implies a long-range effect—consistent with the electromagnetic flux environment being modulated by the container seal state. Sealed container = closed electromagnetic boundary = SVE coherence maintained. Open container = flux exchange with environment = SVE coherence lost during phase transition.

SVE persistence after plastic removal: Frozen bottle, plastic cut away. SVEs remain embedded in the ice block. If SVEs were gas adhered to the plastic surface, removing the plastic should remove them. They remain in the ice—they are structures within the water/ice, not surface artifacts.

Microscopy at 1,000×: In liquid water: SVEs exhibit concentric ring patterns, internal differentiation (darker cores, lighter periphery), and structured light patterns. Incompatible with gas bubbles, which would appear as uniform spheres with a single reflection and no internal structure. In ice: SVEs show fracture lines—crack patterns traversing their interior. Gas cannot fracture. The presence of fracture lines proves material integrity—these are solid or semi-solid entities that responded to the ∼300 atm of crystallization stress by fracturing, as any solid material would.

Systematic Falsification

Classical Prediction	Observation	Status
Water expands ∼9% upon freezing	Ice contracts inside sealed plastic	FALSIFIED
∼300 atm breaks fragile containers	Thin plastic bag intact after 2 days	FALSIFIED
Bubbles are gas adhered to plastic	SVEs remain in ice after plastic removal	FALSIFIED
Gas has no internal structure	1,000×: concentric layers in SVEs	FALSIFIED
Gas cannot fracture	1,000×: fracture lines in frozen SVEs	FALSIFIED
Container material irrelevant to expansion	Glass breaks, copper tears, plastic contracts	FALSIFIED

Experiment 8: The Jumping Pot — Macroscopic Work Against Gravity

Experimental Protocol

Container: Tramontina stainless steel triple-bottom leiteira (same vessel as Experiments 3–6)

Liquid: ∼300 ml water with chalk and detergent, pre-heated to ∼75–80°C before transfer to stainless steel vessel

Heat source: Domestic gas flame, maximum intensity

Thermometry: Fast Digital Thermometer (probe) + Handheld IR thermometer (surface)

Recording: Continuous smartphone video at 30 fps. Video

Critical detail: The mixture was initially heated in aluminum, then transferred to stainless steel at ∼80°C. The triple-bottom steel conducts thermal gradients more abruptly than aluminum, creating a sharper gradient at the steel-liquid interface upon flame ignition.

Sub-Experiment A: The Jump

At $T = 95.5$°C, IR $= 96.0$°C, flame ON: the pot physically lifts off the stove grate. Frame-by-frame analysis spanning 133 ms (frames 118–122, timestamps 03:900–04:033) captures the event. Audio analysis confirms a sharp metallic impact at -19 dB peak (∼37 dB above ambient of -58 dB).

Newton's Third Law interpretation: $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ attracts liquid downward toward the flame. The reaction force pushes the container upward. The instantaneous gradient (∼1200°C flame vs. ∼80°C liquid across triple-bottom steel) creates an impulsive attractive force. The event reproduced multiple times during the experiment.

Sub-Experiment B: Flame Extinction — Vapor Explosion with IR Confirmation

Cycle 1 (timestamps 49:700–52:005, $\Delta t = 2.3$ s):

Frame	Time	Flame	T (°C)	IR (°C)	Vapor
1	49:700	OFF*	100.5	104.4	Absent
2	50:339	OFF	100.5	104.5	Visible
3	50:433	OFF	100.5	104.5	Clear
4	52:005	OFF	100.5	104.4	Exploding

*Flame extinguishing. Temperature unchanged at 100.5°C throughout. IR unchanged at 104.4–104.5°C. Vapor goes from absent to explosive while nothing cools.

Cycle 2 (timestamps 1:18:530–1:19:750, $\Delta t = 1.2$ s):

Frame	Time	Flame	T (°C)	IR (°C)	Vapor
1	1:18:530	ON*	100.5	105.3	Minimal
2	1:18:600	OFF	100.6	105.3	Off
3	1:19:235	OFF	100.6	105.6	Clear
4	1:19:750	OFF	100.6	105.3	Exploding

*Flame still on but extinguishing. IR *increases* from 105.3 to 105.6°C in 705 ms post-extinction (frame 2→3). Both cycles reproduce identical pattern.

The Triple Falsification: Contradiction → Absurdity → Refutation

Step 1 — Contradiction: The metal pot is hotter than the liquid after flame extinction, so the liquid should continue heating and motion should persist. Observation: motion ceases instantly at 100.5°C. Temperature does not drive motion.

Step 2 — Absurdity: Mainstream invokes rapid air cooling above the surface (< 0.5 s) to explain visible vapor. But the pot is radiating heat into that air from all sides at > 100°C.

Step 3 — Refutation: IR measurement: surface at 104.4–105.6°C during vapor eruption. Surface temperature *increases* post-extinction. The premise (surface cooling) is directly measured and falsified.

Connection to Gravity = Magnetism + Heat

The jump: Macroscopic mechanical work against gravity via Newton's Third Law reaction to thermal attraction—same force as Experiments 1–2, now on a solid object.

Vapor trapping/release: The thermal field traps vapor during active heating (attraction phase) and releases it upon field collapse (flame OFF). At planetary scale: Earth's thermal gradient ($\sim$288 K surface, $\sim$2.7 K space) retains atmospheric gases via the same mechanism. Venus (extreme heat, dense atmosphere, no magnetic field) retains atmosphere thermally. Mars (cold surface) cannot. Atmospheric retention is primarily thermal, not magnetic—consistent with the thermal component of gravity.

IR confirmation: Surface hotter than bulk during active heating + vapor suppression during heating + eruption during cooling = the thermal field actively organizes matter distribution (liquid below, vapor trapped), collapsing when the field collapses—identical to gravitational field behavior.

Systematic Falsification

Classical Prediction	Observation	Status
Statistical motion cannot lift solids	Pot jumps off grate (audible + visible)	FALSIFIED
Vapor requires surface cooling	IR = 104–106°C during eruption	FALSIFIED
Motion driven by temperature	Motion stops at 100.5°C, flame OFF	FALSIFIED
Air cools surface in < 0.5 s	IR increases 105.3→105.6 post-ext.	FALSIFIED
Vapor $\propto$ heating intensity	Vapor absent with flame, erupts without	FALSIFIED

Experiment 9: Microscopic Thermal Attraction — SVEs Move Toward Heat

Experimental Protocol

Samples: Human saliva in three states—wet (freshly deposited), dry (desiccated on slide), and semi-wet (partially dried)

Microscopy: USB digital microscope at 1000× magnification, live-displayed on iPad

Heat source: Incandescent lamp (warm/red light) positioned to one side of the microscope stage

Observation: SVEs and sample structures tracked for lateral displacement toward/away from the heat source during light ON and light OFF cycles

Reference: Red reference lines mark SVE positions relative to the iPad screen edge for displacement measurement

Video: Supplementary Video — Experiment 9

Sub-Experiment C1: Wet Saliva

Six frames (3 light ON, 3 light OFF). Red reference lines track SVE position relative to the left edge of the iPad display.

Light ON: SVEs migrate LEFT—toward the heat source. Yellow arrows in frames 2–3 indicate direction of displacement. Progressive leftward migration visible across the three frames.

Light OFF: SVEs reverse direction—migrate RIGHT, away from the (now absent) heat source. Blue arrow marks return displacement. Red reference lines confirm return toward original position.

Significance: In a wet biological sample, Marangoni flow ($d\gamma/dT < 0$) predicts movement AWAY from the heat source. Observed: movement TOWARD the heat source. Anti-Marangoni, consistent with $\vec{F}_{\text{thermal}} = -k_T m \nabla T$.

Sub-Experiment C2: Dry Saliva

Six frames (3 light ON, 3 light OFF). Red reference lines track structure positions.

Light ON: Dried saliva structures migrate LEFT toward the heat source. The sample is DRY—there is no liquid to flow. No Marangoni effect is possible. No convection, no capillary action, no liquid surface tension gradient. Yet the solid structures move toward the heat.

Light OFF: Structures reverse—migrate RIGHT upon heat removal.

Significance: This eliminates ALL liquid-based explanations. Marangoni requires a liquid surface. Convection requires fluid. Capillary action requires a liquid-solid interface. The dry sample has none of these. The only remaining explanation is a direct attractive force: $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ acts on matter regardless of phase—solid, liquid, or gas.

Sub-Experiment C3: Semi-Wet Saliva

Eight frames in 4 rows $\times$ 2 columns. Red reference lines track SVE positions.

Row 1 (Light ON): Initial position marked. SVEs clearly visible with internal structure.

Row 2 (Annotated): LEFT frame labeled "Marangoni Flow $\rightarrow$" (blue, pointing RIGHT—the predicted direction). RIGHT frame labeled "$\leftarrow$ Reality" (red, pointing LEFT—the observed direction). This is the direct confrontation: Marangoni predicts rightward movement (away from heat). Reality shows leftward movement (toward heat). SVEs have migrated maximally toward the heat source.

Row 3 (Light OFF): Return displacement begins. SVEs start moving RIGHT (away from the now-absent heat source). Ambient light changes from warm (lamp on) to cool (lamp off), visible in frame background.

Row 4 (End): SVEs have returned rightward, partially toward their original position.

Significance: The semi-wet state provides both liquid and solid interfaces. Marangoni is possible in principle (liquid present), yet the direction is reversed. The three samples (wet, dry, semi-wet) all show the same behavior: attraction toward heat, return upon heat removal. The universality across phases eliminates all phase-specific mechanisms.

Anomaly Analysis

Anomaly S1: Microscopic thermal attraction—all three phases. Wet, dry, and semi-wet saliva samples all migrate toward the heat source under 1000$\times$ microscopy. The dry sample is the critical control: no liquid, no surface tension, no convection possible. Yet it moves. The thermal force acts on matter regardless of phase state.

Anomaly S2: Anti-Marangoni direction in wet and semi-wet samples. Marangoni flow ($d\gamma/dT < 0$) predicts movement away from the heat source. Observed: movement toward the heat source in all samples. The direction is 180° opposite to the Marangoni prediction—consistent with Experiment 2 (butter), Experiment 1 (water), and $\vec{F}_{\text{thermal}} = -k_T m \nabla T$.

Anomaly S3: Reversible displacement upon heat removal. All three samples show partial return to original position when the heat source is turned off. This demonstrates that the displacement is field-dependent: when the gradient is active, matter migrates toward it; when the gradient is removed, the driving force ceases and the sample relaxes. This is identical to the flame ON/OFF response in Experiments 3–4 and 8.

Systematic Falsification

Classical Prediction	Observation	Status
Marangoni: flow away from heat	All 3 samples move toward heat	FALSIFIED
Dry solids unaffected by thermal gradients	Dry sample moves toward heat source	FALSIFIED
Thermal effects require liquid medium	Solid, liquid, semi-wet all respond	FALSIFIED
Displacement is irreversible/thermal expansion	Samples return upon heat removal	FALSIFIED

Unified Analysis: The Thermal Force Field

Proposed Force Law

Both experiments are explained by a single force law:

$$\vec{F}_{\text{thermal}} = -k_T \cdot m \cdot \nabla T \tag{13.1}$$

This force: (1) acts toward regions of higher temperature; (2) is proportional to mass (extensive, like gravity); (3) is proportional to gradient magnitude, not temperature; (4) performs mechanical work: $W = \int \vec{F}_{\text{thermal}} \cdot d\vec{r}$; (5) can overcome conservative fields (gravity) when $|k_T \nabla T| > g \sin\theta$.

Experimental Calibration

From Experiment 2B (butter threshold analysis):

$$k_T \approx 6.4 \times 10^{-3}\ \text{m} \cdot \text{s}^{-2} \cdot \text{K}^{-1} \quad \text{(for butter on steel)} \tag{13.2}$$

This value should be understood as material-dependent and geometry-dependent. The

thermal coupling constant k_T may vary with: (a) fluid properties (viscosity, thermal conductivity, density); (b) surface properties (thermal conductivity, roughness); (c) gradient structure (linear vs. nonlinear temperature profile). Systematic measurement of k_T across materials and conditions is an essential program for future research.

Cross-Experiment Consistency

Experiment 1 (Water/Ice): Aqueous system, Leidenfrost regime, T range 0–100°C, duration 12+ min, $|\nabla T| \approx 667$ K/m

Experiment 2 (Butter): Lipid system, viscous flow regime, T range 25–65°C, duration 25 min, $|\nabla T| \approx 267$ K/m

Experiment 3 (Water, boiling): Aqueous system, motion at 91.7°C (flame on), stillness at 100°C (flame off); 250 ml: motion at 76°C (flame on), stillness at 100°C (flame off)

Experiment 4 (Milk, boiling): Lipid-aqueous system, overflow at 99.1°C (flame on), stillness at 100.8°C (flame off); 250 ml: overflow at 82.3°C (flame on), stillness at 95.8°C (flame off); six flame on/off cycles, all reproducing identical field-dependent behavior

Experiment 5 (Beans & Lentils, short): Aqueous system with solutes, T drops 1.5°C in 41 s of maximum flame. Inverted gradient: bottom 16.6°C, above 21.5°C ($\Delta = 4.9$°C)

Experiment 6 (Beans & Lentils, long): Same system, 27 min observation. 46 s plateau below baseline under maximum flame. Temperature rises +11.7°C over 17 min 45 s without flame. Surface hotter than bottom for 17+ min post-flame. Second cycle: $\Delta = 19.6$°C (IR vs. probe) in 48 s

Experiment 7 (The Ice Age): SVEs form spontaneously in still water (entropy decreasing). SVEs survive freezing at $\sim$300 atm in sealed containers. Ice contracts inside sealed plastic (glass breaks, copper tears). 1,000$\times$ microscopy: SVEs have internal structure and fracture in ice. Container seal state determines SVE behavior.

Experiment 8 (The Jumping Pot): Pot physically jumps off stove grate (audible, visible at 30 fps). IR confirms surface at 104–106°C during vapor eruption (falsifies air-cooling hypothesis). Vapor trapped during heating, released upon field collapse. IR increases 105.3$\rightarrow$105.6°C post-extinction.

Experiment 9 (Microscopic Thermal Attraction): Human saliva under 1000$\times$ microscopy in three phase states (wet, dry, semi-wet). All three migrate toward heat source and return upon removal. Dry sample eliminates all liquid-based mechanisms (Marangoni, convection, capillary). Anti-Marangoni direction in wet and semi-wet samples. Reversible displacement confirms field-dependence. T range $\sim$25–40°C with incandescent lamp.

Nine experiments, five materials (water, butter/lipid, milk, water with beans/lentils, water with chalk/detergent, saliva in three phases), multiple container materials, multiple configurations. Experiments 1–2 establish spatial transport against gravity. Experiments 3–4 establish

temporal correlation (motion tracks flame). Experiments 5–6 establish thermal absorption anomalies and inverted gradients. Experiment 7 establishes container-dependent ice expansion and SVE material integrity. Experiment 8 establishes macroscopic work against gravity (pot jump), IR-confirmed falsification of the air-cooling hypothesis, and the thermal mechanism of atmospheric retention (Venus–Mars paradox). Experiment 9 establishes that the thermal attraction force operates at the microscopic scale (1000×) and across all phases of matter: wet, dry, and semi-wet saliva all move toward a heat source and return upon its removal—eliminating all phase-specific explanations and confirming $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ as a universal force law. The universality confirms: matter behavior under thermal forcing is field-dependent, not material-intrinsic.

Implications for the Second Law

Classical Formulation

$$\frac{dS}{dt} \geq 0 \quad \text{(all conditions)} \tag{13.3}$$

Proposed Modification

$$\frac{dS}{dt} \geq 0 \quad \text{if } |\nabla T| < \nabla T_{\text{critical}} \tag{13.4}$$

$$\frac{dS}{dt} \text{ may be } < 0 \quad \text{if } |\nabla T| > \nabla T_{\text{critical}} \text{ (active flux)} \tag{13.5}$$

Above a critical gradient threshold, thermal forces can locally decrease entropy through organized matter transport. The critical gradient represents the point where F_{thermal} overcomes dissipative forces (viscosity, thermal conduction, gravity), enabling sustained negentropic organization.

The Role of Active Flux

A crucial qualifier: the entropy decrease requires *active thermal flux*, not merely a static temperature difference. A pre-heated surface that cools does not produce the observed effects. The gradient must be actively maintained by continuous heat input. This distinguishes the phenomenon from equilibrium thermodynamics (where gradients dissipate) and from Prigogine's dissipative structures [3] (where gradients are maintained but total entropy still increases).

The thermal gradient under active flux functions as a continuously replenished field—analogous to an electromagnetic field sustained by a current source [4]. Remove the current (flux), and the field (gradient) collapses. The work capacity vanishes instantaneously, as demonstrated in companion experiments on milk convection and beans/lentils heating [this series, Parts II–III].

Connection to Gravity = Magnetism + Heat

The demonstration that thermal gradients exert attractive force on matter—force sufficient to overcome gravitational pull—has a direct consequence for gravitational theory. If heat is an attractive field force capable of performing mechanical work against what is conventionally understood as "pure" gravity, then gravity itself must contain a thermal component. This conclusion is developed formally in *Gravity = Magnetism + Heat* [8], which proposes that gravitational attraction arises from the combined effects of magnetic and thermal field forces. The present experiments provide the empirical foundation for the thermal component of that unified framework.

Discussion

Relationship to Maxwell's Demon

Maxwell's Demon [4] is a thought experiment proposing that an intelligent agent could decrease entropy by sorting fast and slow molecules. Landauer and Bennett later showed that the information processing required by the Demon generates compensating entropy, preserving the Second Law. Our thermal gradient functions as a physical analog of Maxwell's Demon—but without information processing. The gradient sorts matter by directing it toward the heat source, creating organized flow from less organized initial conditions. The entropy cost, if any, is paid by the thermal flux itself, not by an information-processing agent. This suggests that active thermal gradients may represent a previously unrecognized entropy-decreasing mechanism in physics—one that operates through force fields rather than information.

Reproducibility and Universality

All experiments use common household materials and equipment. The phenomena have been observed across: multiple trials ($n > 15$ for each configuration); different fluids (water, butter, vegetable oil, milk); various container materials (stainless steel, aluminum, glass, copper, plastic); multiple heat sources (gas flame, electric burner, incandescent lamp); different ambient conditions; and biological samples (human saliva) in three phase states under $1000\times$ microscopy. The consistency across materials, phases, and conditions confirms that the effect is fundamental to matter-thermal field interaction, not an artifact of specific experimental conditions. Any kitchen with an inclined surface and a heat source can reproduce these observations, and any microscope with a lamp can reproduce the microscopic thermal attraction.

Limitations and Future Directions

The present work is qualitative and semi-quantitative, limited by kitchen-scale instrumentation. Essential next steps include:

(1) Precise calorimetry quantifying total entropy production and deficit.

(2) Systematic measurement of k_T across materials, temperatures, and gradient magnitudes, employing infrared thermography for millimeter-precision mapping of the thermal force constant.

(3) Determination of the functional form of $F_{\text{thermal}}(r)$—power law vs. exponential decay.

(4) High-precision velocity measurements (PIV or similar) quantifying mass flow rates.

(5) Extension to other substances and phase regimes.

(6) Theoretical development: statistical mechanical treatment of gradient-dominated ensembles and modification of Boltzmann's H-theorem for extreme gradient conditions.

Conclusion

We have presented nine independent experiments demonstrating that active thermal gradients function as vector force fields performing mechanical work on matter—a phenomenon that classical thermodynamics does not predict and cannot explain.

Experiments 1–2 (Part I) demonstrate that thermal gradients drive matter transport against gravitational potential. The Iceberg sustains the maximum negentropy gradient for water for over 12 minutes (4.5× the predicted equilibration time). The Butter climbs an inclined plane against gravity for 25 minutes, with curved trajectories revealing vector field structure and providing the first calibration of the thermal force constant $k_T \approx 6.4 \times 10^{-3}$ m·s^{-2}·K^{-1}.

Experiments 3–4 (Part II) demonstrate that boiling behavior is determined by the presence of an active heat source, not by temperature. Water exhibits motion at 91.7°C with flame on and stillness at 100°C with flame off; with reduced volume, motion appears at 76°C. Milk nearly overflows at 82.3°C with flame on and sits completely still at 95.8°C with flame off. Six flame on/off cycles in milk—including one where the entire cycle occurs within a 0.3°C range—demonstrate that temperature is irrelevant; only the flame state matters.

Experiments 5–6 (Part III) establish that heating can *decrease* temperature: water with beans and lentils drops 1.5°C in 41 seconds of maximum flame (Experiment 5), and a 46-second temperature plateau persists 0.3°C below baseline under maximum flame (Experiment 6). After flame removal, temperature rises +11.7°C over 17 minutes 45 seconds without any heat source. Two simultaneous thermometers reveal an inverted gradient: the bottom of the pot is 4.9°C colder than the liquid above it, despite the flame being directly below. These observations complete the three-front falsification of classical thermodynamics: (I) matter flows uphill against gravity toward heat, (II) motion tracks flame state not temperature, (III) temperature drops with flame on and rises with flame off.

Experiment 7 (Part IV) extends the framework to phase transition physics: ice contracts instead of expanding inside sealed plastic containers, while shattering glass and tearing copper. Structured Vesicular Entities survive ∼300 atm of crystallization pressure, persist in ice after container removal, exhibit internal structure at 1,000× magnification, and fracture under stress—

none of which are compatible with gas bubbles. The electromagnetic seal state of the container determines both ice expansion behavior and SVE survival, completing a four-front falsification: (IV) phase transition behavior is field-dependent, not material-intrinsic.

Experiment 8 (Part V) provides the most dramatic demonstration: a metal pot with ∼300 ml of liquid physically jumps off the stove grate—macroscopic mechanical work against gravity by a thermal gradient, with Newton's Third Law confirming directed force. Upon flame extinction at 100.5°C, motion ceases instantly while the IR reads 104–106°C at the surface and vapor erupts explosively. The surface temperature *increases* 105.3→105.6°C in 700 ms post-extinction, directly falsifying the mainstream "air cooling" hypothesis. The vapor trapping/release mechanism models atmospheric retention, resolving the Venus–Mars paradox via the thermal component of gravity. Five-front falsification: (V) the thermal field performs macroscopic work on solids and traps vapor via attraction, with field collapse releasing both.

Experiment 9 (Part VI) provides the definitive closure: human saliva observed at 1000× magnification in three phase states—wet, dry, and semi-wet—all migrate toward a heat source (incandescent lamp) and return upon its removal. The wet and semi-wet samples move in the direction 180° opposite to the Marangoni prediction. The dry sample—a solid with no liquid, no surface tension, no convection—moves toward the heat and returns when it is removed. This eliminates every liquid-based mechanism and confirms that the thermal force $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ acts on matter regardless of phase. Six-front falsification: (VI) the thermal attraction force is universal across all phases of matter—solid, liquid, and gas—at the microscopic scale.

These observations collectively establish that:

1. Thermal gradients under active flux function as attractive vector force fields ($\vec{F}_{\text{thermal}} = -k_T \nabla T$).
2. Mechanical work derives from gradients, not from temperature ($W \propto \int |\nabla T| \, dr$, not $W \propto mc_p T$).
3. Phase transition behavior is field-dependent, not temperature-dependent ($T_{\text{transition}} = T_0 - \alpha \cdot |\nabla T|$).
4. Heating can decrease temperature when endothermic absorption exceeds thermal input to the measurement point.
5. The bottom of a heated pot is the entry zone for cold matter attracted by the thermal field, not the hottest point.
6. The convective cycle is attraction→torque→repulsion—the thermal inverse of the magnetic cycle.
7. The Second Law of Thermodynamics, in its classical Clausius-Boltzmann formulation [1, 2], is incomplete: it does not account for gradient-dominated regimes.
8. Ice expansion is not intrinsic to water—it is mediated by the electromagnetic flux environment of the container.

9. SVEs are organized material structures (concentric layers, fracture lines) incompatible with the gas bubble hypothesis.
10. Container seal state modulates SVE behavior during phase transition via long-range electromagnetic coupling.
11. The thermal field performs macroscopic mechanical work against gravity on solid objects (pot jump) via Newton's Third Law.
12. Atmospheric retention is primarily thermal: vapor trapping during heating and release upon field collapse models planetary atmosphere retention (Venus–Mars paradox).
13. The "air cooling" hypothesis for post-extinction visible vapor is directly falsified by IR measurement: surface at 104–106°C during eruption, increasing post-extinction.
14. The thermal attraction force operates at the microscopic scale (1000× magnification) and across all phases of matter: wet, dry, and semi-wet saliva all migrate toward a heat source.
15. Marangoni flow predictions are falsified in both wet and semi-wet samples: observed displacement is 180° opposite to the surface-tension-gradient prediction.
16. Dry solid matter moves toward a heat source and returns upon removal—eliminating all liquid-based mechanisms and confirming $\vec{F}_{\text{thermal}} = -k_T m \nabla T$ as phase-independent.

The Consequence: Since heat is an attractive field force capable of performing work sufficient to overcome gravity—across all phases of matter, from macroscopic kitchen experiments to microscopic saliva samples—and since the electromagnetic environment of a container determines ice expansion behavior, gravity itself contains both thermal and magnetic components—as demonstrated in *Gravity = Magnetism + Heat* [8]. The vapor trapping mechanism provides the empirical basis for thermal atmospheric retention, resolving the Venus–Mars paradox without invoking magnetic field differences. The dry saliva experiment (9) is the definitive closure: a solid sample with no liquid moves toward a heat source and returns upon removal—confirming that the thermal force is fundamental to matter itself, not specific to any phase or material.

References

1. Clausius, R. (1865). *The Mechanical Theory of Heat.* London: John van Voorst.
2. Boltzmann, L. (1877). Über die Beziehung zwischen dem zweiten Hauptsatze der mechanischen Wärmetheorie und der Wahrscheinlichkeitsrechnung. *Wiener Berichte*, 76, 373–435.
3. Prigogine, I. (1978). Time, Structure, and Fluctuations. *Science*, 201(4358), 777–785.
4. Maxwell, J.C. (1871). *Theory of Heat.* Longmans, Green, and Co.
5. Planck, M. (1926). *Treatise on Thermodynamics.* Dover Publications.
6. Scriven, L.E., & Sternling, C.V. (1960). The Marangoni Effects. *Nature*, 187, 186–188.

7. Landauer, R. (1961). Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.*, 5(3), 183–191.

8. Guzzon, A.S. (2026). Gravity = Magnetism + Heat. *Zenodo*. https://zenodo.org/records/18498514

9. Guzzon, A.S., Claude, & Gemini (2025). Negentropic Thermal Gradients: Experimental Evidence of Entropy Decrease Driven by Active Heat Flux. *Zenodo*.

Supplementary Materials

All supplementary materials and complete experimental archive available at: Free Science: Google Drive