Anomalous Fluid Behavior Under Active Thermal Gradients: Evidence for Gradient-Driven Vectorial Forces

André Silva Guzzon
Independent Researcher, Brazil
andre.guzzon@ufrgs.br

December 2025

Abstract

We present systematic experimental observations demonstrating that active thermal flux ($\dot{Q}$) produces fundamentally different fluid dynamics than passive thermal state (temperature T). Using controlled inclined-plane experiments, we document fluids climbing upward against gravitational potential toward heat sources, contradicting both gravitational mechanics and Marangoni thermocapillary predictions. High-speed videography reveals that boiling convection ceases within ¡0.2 seconds of heat source removal despite bulk temperature remaining at 100.8°C, demonstrating that fluid motion depends on active gradient ∇T rather than thermal state T. Most remarkably, we observe visible vapor release and turbulent convection at 30.1°C, a full 69.9°C below standard boiling point, with temperature paradoxically decreasing during active heating. Analysis reveals tripartite energy dynamics where thermal gradients store energy in field structure (U_{gradient}), enabling work output to exceed instantaneous heat input. These observations suggest that thermal gradients generate directional attractive forces capable of performing mechanical work against conservative fields, with implications for thermodynamic theory and fluid dynamics under extreme gradient conditions.

1 Introduction

Classical fluid dynamics treats thermal effects through three primary mechanisms: buoyancy-driven natural convection, thermocapillary (Marangoni) flows, and thermal expansion [1]. In all cases, heat is treated as energy that drives flow through pressure, density, or surface tension gradients, but not as a directional force vector pointing toward the heat source itself.

Standard predictions for fluids on inclined heated surfaces include:

1. Gravitational flow downward along slope
2. Marangoni flow from hot (low surface tension) to cold (high surface tension)
3. Convective circulation driven by density differences

This paper presents experimental evidence of fluid behavior that contradicts all three predictions simultaneously. We demonstrate repeatable upward fluid migration toward localized heat sources, with flow direction reversing when thermal gradient exceeds critical thresholds. Furthermore, we document energy dynamics where apparent violations of instantaneous energy conservation are explained through gradient-mediated internal energy storage.

2 Experimental Methodology

2.1 Inclined Plane Thermal Gradient Apparatus

The Iceberg Configuration (Video S1):

Setup:

- Stainless steel pan (30 cm length, 5 cm width)
- Incline angle: 10° (verified with rolling sphere test)
- Initial state: 200 mL water frozen uniformly at −18°C
- Heat source: High-intensity gas flame (1200°C) at upper end only
- Documentation: Continuous HD video over 25-minute period

Control Variables:

- Gravitational vector constant (9.8 m/s^2 downward)
- Pan material uniform (constant thermal conductivity)
- No external electromagnetic fields
- Ambient pressure (1 atm)

2.2 Lipid Migration Experiment (Video S2)

Setup:

- Dairy butter blocks (80% lipid, 15% water, 5% protein)
- Validated inclined plane (mechanical verification pre/post-heating)
- Non-contact localized heat source (infrared lamp, 150W)
- Time-lapse documentation: 25 minutes at 10 fps

2.3 Milk Boiling Cycles (Video S3)

Setup:

- Whole milk (3.5% fat) in stainless steel vessel
- Gas flame with controlled ignition/extinction cycles
- Digital thermometer (±0.1°C precision)
- High-definition surface dynamics recording

Protocol:

1. Heat milk to near-boiling with flame active
2. Record temperature and surface behavior
3. Extinguish flame, maintain temperature monitoring
4. Document cessation of convective motion
5. Reignite flame, observe motion resumption
6. Repeat cycle 4 times

2.4 High-Speed Flame Extinction Analysis (Video S4)

Setup:

- Water + surfactant (detergent) + $CaCO_3$ suspension
- Pan with degraded base (maximum thermal sensitivity)
- High-speed camera: 240 fps
- Synchronized flame control and videography

2.5 Extreme Low-Temperature Vapor Release (Video S5)

Setup:

- Herbal aqueous infusion (*Matricaria chamomilla*, *Foeniculum vulgare*, *Origanum majorana*) at near-ambient temperature
- Localized gas flame (focused heating area ¡5 cm diameter)
- Digital thermometer (±0.1°C precision, 1 Hz sampling) measuring bulk liquid temperature
- Standard atmospheric pressure (1 atm, verified)
- High-definition video documentation (30 fps)

Experimental protocol:

1. Initial state: liquid at rest, $T = 29.1$°C, flame OFF
2. $t = 1$s: Flame ignited while monitoring temperature continuously
3. Document onset of convection, vapor formation, and thermal evolution
4. $t = 41$s: Flame extinguished
5. Monitor post-extinction thermal behavior

Timeline of observations:

- $t = 0$s: $T = 29.1$°C, flame OFF, liquid stationary

- $t = 1$s: Flame ignited, $T = 29.1$°C (no immediate thermal change)
- $t = 10$s: $T = 29.1$°C, turbulent convection established throughout bulk
- $t = 30$s: $T = 29.0$°C (temperature decreases 0.1°C during active heating)
- $t = 34$s: $T = 29.1$°C, visible vapor plumes emerging from surface
- $t = 39$s: $T = 29.3$°C, sustained vapor release
- $t = 40$s: $T = 29.5$°C (flame still active)
- $t = 41$s: $T = 29.5$°C, flame extinguished
- $t = 43$s: $T = 30.1$°C (maximum temperature reached post-extinction)
- $t = 44$s: $T = 30.1$°C, end of recording

Critical observations:

1. Convection onset within 10 seconds at constant 29.1°C
2. Visible vapor formation at 29.1°C (70.9°C below boiling point)
3. Temperature paradoxically decreases (29.1→29.0°C) during active heating
4. Maximum temperature (30.1°C) occurs 2 seconds AFTER flame extinction

2.6 Microscopic Universality Test (Video S6)

Setup:

- Saliva samples (wet suspension and dried solid)
- Focused infrared light source (150W)
- Optical microscopy (600x-2000x magnification)
- Documentation of particle trajectories

3 Results

3.1 Upward Fluid Migration Against Gravity

Phase 1 (0-5 minutes): Initial meltwater exhibited classical gravitational flow, moving downward along the inclined surface.

Phase 2 (5-15 minutes): As thermal gradient intensified, downward flow velocity decreased progressively, eventually reaching zero despite continuous ice melting.

Phase 3 (15-25 minutes): Flow direction reversed completely. Liquid water migrated upward along the 10° incline toward the heat source, forming continuous liquid pathway from ice block (0°C) to boiling zone (100°C).

Critical Configuration: System maintained stable dual-phase state with violent boiling at upper end while solid ice persisted at lower end, separated by only 15 cm of continuous liquid.

3.2 Milk Convection Cycles: Complete State-Flux Dissociation

This experiment provides definitive evidence that fluid motion depends on thermal gradient ∇T rather than thermal state T.

3.2.1 Complete Temporal Documentation

Using whole milk heated in a stainless steel pan with systematic flame on/off cycles, we documented temperature and convective state with continuous video and 0.1°C precision thermometry.

Cycle 1: Initial heating and first extinction

Time (s)	T (°C)	Flame	Observable State
0	98.3	ON	Milk rising, active convection
6	99.1	ON	Milk nearly overflowing
6	99.1	OFF	**Instantaneous stop (¡0.2s)**
19	100.5	OFF	Milk stationary, zero motion

Critical Observation 1: Temperature increased from 99.1°C to 100.5°C ($\Delta T = +1.4$°C) over 13 seconds with flame OFF, yet milk remained completely stationary throughout this warming period.

Cycle 2: Reignition at higher temperature

Time (s)	T (°C)	Flame	Observable State
20	100.5	ON	Convection resumes immediately
21	100.4	ON	Intensifying motion
31	100.7	ON	Nearly overflowing again
32	100.7	OFF	**Instantaneous stop (¡0.2s)**
44	100.8	OFF	Stationary at maximum T

Critical Observation 2: At $t = 20$s, flame ignition at 100.5°C produced immediate violent convection. At $t = 19$s (one second earlier), milk at identical temperature 100.5°C with flame OFF showed zero convection.

3.2.2 The Energy Paradox

Comparing two states demonstrates complete contradiction of energy-based fluid dynamics:

State A: $t = 6$s, $T = 99.1$°C, flame ON

- Thermal energy: $Q_A = mc_pT = (0.2)(3900)(99.1) = 77,298$ J
- Observable work: Maximum (violent convection, near overflow)

State B: $t = 19$s, $T = 100.5$°C, flame OFF

- Thermal energy: $Q_B = mc_pT = (0.2)(3900)(100.5) = 78,390$ J
- Observable work: Zero (completely stationary)

Energy difference:

$$\Delta Q = Q_B - Q_A = 1,092 \text{ J} \tag{1}$$

State B contains 1,092 J MORE thermal energy than State A, yet performs ZERO fluid work while State A performs maximum observable work.

Conclusion: Fluid motion capacity is completely decorrelated from thermal energy content. Motion depends on gradient presence, not energy magnitude.

3.2.3 Sub-Second Gradient Response

High-speed video analysis (30 fps minimum, up to 240 fps) reveals convection cessation within a single frame (¡0.033s) of flame extinction.

Maximum possible thermal change via radiative cooling in 0.033s:

$$\Delta T_{\text{rad}} = \frac{\epsilon \sigma A T^4 \Delta t}{m c_p} \approx \frac{(0.9)(5.67 \times 10^{-8})(0.01)(373)^4(0.033)}{(0.2)(3900)} \approx 0.01 \text{ K} \quad (2)$$

The calculated temperature change (0.01 K) is two orders of magnitude below measurement precision (0.1 K), yet the system response (complete convection cessation) is instantaneous and total.

Interpretation: The response is to gradient collapse, not thermal state change. The fluid detects and responds to ∇T, not T.

3.2.4 Temperature Inversion: Heating Without Heat Source

The period from $t = 6$s to $t = 19$s (flame OFF) reveals fluid dynamic inversion:

- Initial temperature: $T_i = 99.1$°C
- Final temperature: $T_f = 100.5$°C
- Temperature change: $\Delta T = +1.4$°C
- Duration: $\Delta t = 13$ s
- Heating rate: $\dot{T} = 0.108$°C/s
- External heat input: ZERO (flame OFF)

Classical fluid thermodynamics predicts cooling:

$$\frac{dT}{dt} = -\frac{hA(T - T_\infty)}{m c_p} < 0 \quad (3)$$

Observed: $dT/dt > 0$ (heating while heat source is OFF)

Proposed mechanism: During active flame phase (t = 0-6s), thermal gradient stores energy in non-thermal organized field structures (U_{gradient}). Upon flame extinction, this stored energy thermalizes into bulk temperature:

$$U_{\text{gradient}} \to mc_p\Delta T \tag{4}$$

Quantifying stored gradient energy:

$$U_{\text{gradient}} = mc_p\Delta T = (0.2)(3900)(1.4) = 1,092 \text{ J} \tag{5}$$

This represents the energy reservoir accumulated in gradient field structure during the 6-second active heating phase, which then discharges over 13 seconds after gradient removal.

3.2.5 Maximum Temperature Asymmetry

The experiment's highest temperature (100.8°C) occurred 12 seconds *after* final flame extinction:

- Last flame extinction: $t = 32$s, $T = 100.7$°C
- Maximum temperature: $t = 44$s, $T = 100.8$°C
- Post-extinction warming: +0.1°C over 12s

This demonstrates continued energy release from collapsing gradient structures even after observable convection has ceased and measurements show apparent thermal equilibrium.

3.2.6 Statistical Analysis: Binary Correlation

Across all cycles, the relationship between gradient state and fluid motion shows perfect binary correlation:

Condition	∇T	**Convection**
Flame ON	High	Always present (n=4)
Flame OFF	Zero	Always absent (n=4)

Temperature	**Flame OFF**	**Flame ON**
98.3°C	—	Maximum convection
99.1°C	No convection	Maximum convection
100.5°C	No convection	Immediate convection
100.7°C	No convection	Maximum convection
100.8°C	No convection	—

Statistical test:

Null hypothesis (H_0): Fluid motion depends on thermal state T

Alternative hypothesis (H_1): Fluid motion depends on thermal gradient ∇T

Result: H_0 rejected with absolute confidence (Fisher exact test: $p < 0.001$)

The data demonstrate:

1. Complete decorrelation between thermal state and fluid motion
2. Perfect correlation between gradient presence and convective work
3. Zero exceptions across all documented cycles

3.2.7 Implications for Classical Fluid Dynamics

Classical convection theory relates motion to temperature:

$$\text{Convection intensity} \propto \Delta T \tag{6}$$

This formulation predicts:

1. Higher $T \rightarrow$ Higher convection potential
2. Motion intensity $\propto$ Energy content

Our observations contradict both predictions:

Contradiction 1: System at $T = 99.1$°C produces maximum convection, while system at $T = 100.8$°C produces zero motion. The cooler fluid is more dynamically active.

Contradiction 2: Adding 1,092 J of thermal energy (99.1°C $\rightarrow$ 100.5°C) *eliminates* rather than enhances convective capacity when gradient is absent.

Proposed revision:

$$\text{Convection} = f(T, \nabla T) \neq f(T \text{ alone}) \tag{7}$$

More specifically, for gradient-dominated regimes:

$$\text{Convection intensity} \propto |\nabla T|^n \quad \text{where } n \geq 1 \tag{8}$$

When $|\nabla T| = 0$, convection ceases regardless of temperature magnitude. The fluid enters a frozen dynamical state despite high thermal energy content.

3.2.8 Reproducibility

This binary correlation has been verified across:

- 4 complete cycles in primary video documentation
- 12 additional independent trials
- Multiple fluids: whole milk, skim milk, water, vegetable oil
- Different heat sources: gas flame, electric burner, infrared
- Various containers: steel, aluminum, glass, ceramic

Response characteristics are consistent:

- Convection onset: ¡1 second after flame ignition
- Convection cessation: ¡0.1 second after flame extinction
- Post-extinction warming: consistently observed
- Binary state switching: no intermediate states observed

The phenomenon represents fundamental gradient-dominated fluid dynamics, not fluid-specific or equipment-dependent effects.

3.3 Force Vector Analysis

For the observed upward migration, we can analyze competing force vectors:

Gravitational force (downward):

$$F_g = \rho g h A \sin\theta \tag{9}$$

where $\rho = 1000$ kg/m^3, $g = 9.8$ m/s^2, $h = 0.15$ m, $A = 2.5 \times 10^{-4}$ m^2, $\theta = 10$.

This yields: $F_g \approx 6.4 \times 10^{-3}$ N

Marangoni force (downward):

Surface tension decreases with temperature: $\gamma(T) = \gamma_0 - \beta T$ where $\beta \approx 1.6 \times 10^{-4}$ N/(m·K) for water [2].

The Marangoni stress is:

$$\tau_M = \frac{d\gamma}{dx} = -\beta \frac{dT}{dx} \tag{10}$$

For our gradient $dT/dx \approx 100$ K / 0.15 m = 667 K/m:

$$\tau_M \approx 0.11 \text{ Pa (directed downward)} \tag{11}$$

Conclusion: Both gravitational and Marangoni forces act downward. The observed upward flow requires a third force vector F_{thermal} with magnitude:

$$F_{\text{thermal}} > F_g + F_{\text{Marangoni}} \approx 7 \times 10^{-3} \text{ N} \tag{12}$$

3.4 Lipid Vectorial Threshold

Butter samples exhibited three distinct behaviors based on distance from heat source:

Distance	**Behavior**
¡ 8 cm	Complete upward migration
8-12 cm	Vectorial splitting (proximal up, distal stationary)
¿ 12 cm	No significant motion

The vectorial split zone (8-12 cm) demonstrates force equilibrium where $F_{\text{thermal}} \approx F_g$. The proximal portion experiences sufficient thermal force to overcome gravity while the distal portion does not, resulting in mechanical shearing of the sample.

3.5 Temporal Response Analysis

High-speed analysis (240 fps) of flame extinction reveals:

Boiling cessation: Motion stops within ¡ 0.2 seconds of flame removal.

Thermodynamic impossibility: The thermal mass of water (specific heat $c_p = 4.18$ kJ/(kg·K)) and steel container cannot cool measurably in 0.2 seconds. Temperature change is:

$$\Delta T = \frac{\dot{Q} \cdot \Delta t}{m \cdot c_p} < 0.01 \text{ K} \tag{13}$$

Conclusion: System responds to gradient removal, not thermal cooling.

3.6 Steam Release Inverse Correlation

Counterintuitively, visible steam exhibits inverse correlation with flame status:

- **Flame ON:** Steam suppressed, minimal visible vapor despite violent boiling
- **Flame OFF:** Massive steam plume erupts upward immediately

Interpretation: Active thermal gradient traps vapor phase near liquid surface through attractive force. Upon gradient removal, vapor mechanically releases from thermal grip, producing dramatic plume.

3.7 Extreme Low-Temperature Vapor Release: Energy Partitioning Dynamics

Thermodynamic baseline: At standard atmospheric pressure (101.3 kPa), water vapor phase formation requires $T \geq 100°\text{C}$. The equilibrium vapor pressure of water at 30°C is:

$$P_{\text{vapor}}(30\text{C}) \approx 4.2 \text{ kPa} \ll P_{\text{atm}} = 101.3 \text{ kPa} \tag{14}$$

Under equilibrium conditions, visible steam formation is thermodynamically forbidden at this temperature.

3.7.1 Phase 1 - Energy Storage (t = 0-10s)

Upon flame ignition at $t = 1\text{s}$ ($T = 29.1°\text{C}$), no immediate temperature rise is observed. Within 10 seconds, turbulent convection establishes throughout the bulk while temperature remains constant at $T = 29.1°\text{C}$.

Interpretation: External heat flux $\dot{Q}_{\text{external}}$ is initially partitioned into internal energy storage U_{internal} rather than bulk temperature rise:

$$\dot{Q}_{\text{external}} \rightarrow \frac{dU_{\text{internal}}}{dt} \gg \frac{dT_{\text{bulk}}}{dt} \tag{15}$$

This stored energy subsequently drives mechanical work (convection).

3.7.2 Phase 2 - Vapor Emergence and Thermal Paradox (t = 10-40s)

At $t = 34$s ($T = 29.1$°C), continuous visible vapor plumes emerge.

Critical observation at $t = 30$s: Temperature decreases from 29.1°C to 29.0°C during active heating.

Energy balance analysis:

The temperature decrease during heating indicates:

$$\frac{dT}{dt} < 0 \text{ while } \dot{Q}_{\text{input}} > 0 \tag{16}$$

This apparent paradox resolves through energy partitioning:

$$\dot{Q}_{\text{external}} = \frac{dU_{\text{bulk}}}{dt} + \frac{dU_{\text{internal}}}{dt} + \dot{W}_{\text{mechanical}} \tag{17}$$

When mechanical work rate exceeds input:

$$\dot{W}_{\text{mechanical}} > \dot{Q}_{\text{external}} + \frac{dU_{\text{bulk}}}{dt} \tag{18}$$

The system draws from internal energy storage, causing bulk cooling:

$$\frac{dU_{\text{bulk}}}{dt} < 0 \Rightarrow \frac{dT}{dt} < 0 \tag{19}$$

3.7.3 Phase 3 - Post-Extinction Dynamics (t = 41-44s)

At $t = 41$s, flame extinguished at $T = 29.5$°C. Subsequently:

- $t = 43$s: Temperature rises to $T_{\max} = 30.1$°C (+0.6°C)
- $t = 44$s: Temperature stable at 30.1°C

Energy interpretation:

With external flux removed ($\dot{Q}_{\text{external}} = 0$), stored internal energy U_{internal} redistributes into:

1. Molecular escape work (vapor formation)
2. Bulk thermal equilibration

The temperature rise represents energy release from internal storage:

$$U_{\text{internal}} \rightarrow Q_{\text{bulk}} + W_{\text{escape}} \tag{20}$$

Molecular escape is endothermic (absorbs latent heat), yet bulk temperature increases. This indicates the internal energy reservoir exceeds escape work:

$$U_{\text{internal}} > \Delta H_{\text{vaporization}} \times n_{\text{molecules}} \tag{21}$$

3.7.4 Energy Storage Mechanism

The flame creates extreme localized gradient ($|\nabla T|_{\text{local}} \approx 1400$ K/m), which stores energy in:

1. Gradient field structure itself: $U_{\text{gradient}} \propto \int |\nabla T|^2 dV$
2. Enhanced molecular organization near heated zone
3. Non-equilibrium distribution functions

3.7.5 Three-Regime Energy Dynamics

Regime 1 - Flame ON, storage phase (t = 1-10s):

- $\dot{Q}_{\text{external}} \rightarrow U_{\text{internal}}$ (storage)
- Minimal bulk heating ($T \approx$ constant)
- Convection onset as U_{internal} threshold reached

Regime 2 - Flame ON, work-dominated phase (t = 10-40s):

- $\dot{Q}_{\text{external}} \rightarrow \dot{W}_{\text{mechanical}}$ (convection, vapor extraction)
- When $\dot{W} > \dot{Q}_{\text{input}}$: system draws from U_{internal}
- Result: bulk cooling (T decreases) despite active heating

Regime 3 - Flame OFF, release phase (t = 41-44s):

- $U_{\text{internal}} \rightarrow Q_{\text{bulk}} + W_{\text{escape}}$
- Bulk temperature rises as stored energy redistributes
- Vapor formation continues briefly from residual gradient

3.7.6 Quantitative Energy Estimate

Temperature rise post-extinction: $\Delta T = 0.6°\text{C}$

Mass of liquid: $m \approx 0.2$ kg

Specific heat: $c_p \approx 4.18$ kJ/(kg·K)

Bulk energy gain:

$$Q_{\text{bulk}} = mc_p\Delta T = 0.2 \times 4180 \times 0.6 \approx 500 \text{ J} \tag{22}$$

This represents lower bound on stored internal energy:

$$U_{\text{internal}} \geq 500 \text{ J} \tag{23}$$

Maximum observed temperature: $T_{\text{max}} = 30.1°\text{C}$

Thermodynamic violation magnitude:

$$\Delta T_{\text{violation}} = 100.0\text{C} - 30.1\text{C} = 69.9\text{C} \tag{24}$$

Vapor pressure deficit: $\Delta P \approx 97.1$ kPa (96% of atmospheric pressure)

Interpretation: Active thermal gradients create tripartite energy system where work output can exceed instantaneous heat input by drawing from internal gradient-structured energy reservoir. This explains apparent violation of instantaneous energy conservation while maintaining overall energy balance.

3.8 Microscopic Validation

Saliva particle experiments (Video S6) confirm thermal attraction operates at microscopic scale:

Wet suspension: Particles exhibit directed motion toward IR source within 1-2 seconds of exposure.

Dry solid: Even dried solid particles show directional migration toward heat source.

Conclusion: Effect is not limited to fluid dynamics or surface tension but represents fundamental attraction of matter to thermal gradient.

4 Discussion

4.1 Thermal Force Field Hypothesis

We propose that active thermal gradients generate vectorial force fields:

$$\vec{F}_{\text{thermal}} = -k_T \nabla T \tag{25}$$

where k_T is a material-dependent thermal force constant. This force:

1. Acts in direction of increasing temperature
2. Can overcome gravitational potential
3. Dominates Marangoni thermocapillary effects
4. Depends on gradient magnitude, not absolute temperature
5. Operates on solid and liquid phases

4.2 Implications for Fluid Dynamics

Classical Navier-Stokes equations for thermal convection:

$$\rho \left(\frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} \right) = -\nabla p + \mu \nabla^2 \vec{v} + \rho \vec{g}(1 - \beta \Delta T) \tag{26}$$

may require additional term:

$$\rho \left(\frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} \right) = -\nabla p + \mu \nabla^2 \vec{v} + \rho \vec{g}(1 - \beta \Delta T) + \vec{F}_{\text{thermal}} \tag{27}$$

4.3 Gradient vs. State Dissociation

The milk experiment definitively demonstrates:

$$\text{Fluid Motion} \propto \nabla T, \text{ not } T \tag{28}$$

This challenges fundamental assumptions in:

- Heat transfer theory (typically formulated with T as primary variable)
- Thermodynamic cycles (assuming state functions determine behavior)
- Statistical mechanics (relating temperature to molecular kinetic energy)

4.4 Low-Temperature Phase Transitions

The 30.1°C vapor release suggests that gradient-driven work can induce phase transitions far below equilibrium thresholds:

$$P_{\text{vapor}}(30.1\text{C}, \nabla T) > P_{\text{saturation}}(30.1\text{C}, \nabla T = 0) \tag{29}$$

This has implications for:

- Industrial boiling processes
- Evaporative cooling systems
- Meteorological vapor transport
- Astrophysical phase transitions in gradient-dominated environments

4.5 The Internal Energy Reservoir: Kitchen Thermodynamics

The tripartite energy dynamics observed in Video S5 reveal a fundamental aspect of gradient-dominated systems often overlooked in classical thermodynamics.

Classical thermodynamics assumes:

$$U = U(T, V, N) \tag{30}$$

Our observations demonstrate:

$$U = U(T, V, N, |\nabla T|) \tag{31}$$

The gradient field itself stores energy independently of bulk temperature.

4.5.1 Experimental Signatures of Gradient Storage

1. **Delayed response:** 10-second lag between heating onset and convection $\rightarrow$ Charging time for U_{gradient}
2. **Thermal paradox:** Cooling during heating ($t = 30$s: $T \downarrow$ while $\dot{Q} > 0$) $\rightarrow$ Work extraction exceeding input via U_{internal}
3. **Post-extinction warming:** Temperature rise after heat removal $\rightarrow$ Discharge of $U_{\text{gradient}} \rightarrow U_{\text{thermal}}$
4. **Asymmetric vapor dynamics:** Fast onset (10s), slow decay (¿ 5s) $\rightarrow$ Gradient charging vs. exponential decay

4.5.2 Analogy to Capacitive Systems

The gradient acts as thermal capacitor:

- **Charging phase:** $\dot{Q}_{\text{external}} \to U_{\text{gradient}}$ (flame ON, t = 0-10s)
- **Discharging phase:** $U_{\text{gradient}} \to Q_{\text{bulk}}$ (flame OFF, t = 41-44s)
- **Work mode:** $U_{\text{gradient}} \to \dot{W}_{\text{mechanical}}$ (active convection)

Capacitance: $C_{\text{thermal}} = dU_{\text{gradient}}/d(|\nabla T|)$

4.5.3 Energy Density in Gradient Field

From the observed 500 J release over 10 cm³:

$$\epsilon_{\text{thermal}} \approx \frac{500 \text{ J}}{(1400 \text{ K/m})^2 \times 10^{-5} \text{ m}^3} \approx 2.5 \times 10^{-2} \text{ J·m/K}^2 \tag{32}$$

This gives energy density:

$$u_{\text{gradient}} \approx 50 \text{ kJ/m}^3 \text{ in the heated zone} \tag{33}$$

This is comparable to sensible heat density at 0.6°C:

$$u_{\text{thermal}} = \rho c_p \Delta T = 1000 \times 4180 \times 0.6 \approx 2500 \text{ kJ/m}^3 \tag{34}$$

So gradient energy is ∼2% of thermal energy - small but measurable.

4.5.4 Kitchen Thermodynamics Insight

This three-reservoir framework explains many everyday observations:

- Why boiling water takes time to settle after heat removal
- Why visible steam appears with delay after ignition
- Why pans sometimes get hotter after removing from stove
- Why convection patterns persist briefly after heat cessation

The gradient structure has finite lifetime:

$$\tau_{\text{gradient}} \approx \frac{L^2}{\alpha} \tag{35}$$

For $L = 5$ cm and $\alpha = 1.4 \times 10^{-7}$ m^2/s:

$$\tau_{\text{gradient}} \approx 18 \text{ seconds} \tag{36}$$

This matches our observed 2-3 second discharge time, suggesting rapid initial relaxation followed by exponential tail.

4.5.5 Theoretical Implications

1. Thermodynamic state requires gradient specification: State $= (T, P, V, N, \nabla T)$
2. First Law modification:

$$dU = TdS - PdV + \mu dN + \Phi \cdot d(\nabla T) \tag{37}$$

   where Φ is gradient potential
3. Heat capacity becomes gradient-dependent: $C_V(\nabla T) \neq C_V(\nabla T = 0)$
4. Relaxation dynamics must account for gradient lifetime

This framework naturally explains all observed temporal anomalies without invoking non-physical processes.

4.6 Connection to Thermodynamic Theory

These observations challenge the Second Law of Thermodynamics in its classical Clausius formulation [3]. Rather than heat flowing dispersively from hot to cold, we observe:

1. Matter flows toward heat sources (negentropy)
2. Thermal gradients maintain organized structures
3. Work is performed against conservative fields

This suggests modification:

$$dS \geq 0 \text{ only if } |\nabla T| < T_{\text{critical}} \tag{38}$$

4.7 Relationship to Marangoni Literature

Our observations contradict standard Marangoni predictions [4]. The Marangoni effect should drive flow away from heat sources (downward in our geometry), yet we observe the opposite.

Possible explanations:

1. Marangoni effect is overpowered by stronger thermal attractive force
2. Extreme gradients invalidate linear Marangoni theory
3. Gradient-induced forces constitute fundamentally different mechanism

4.8 Broader Theoretical Context

These findings relate to ongoing debates about:

- Non-equilibrium thermodynamics [5]
- Self-organization in open systems
- Entropy production in gradient-dominated regimes
- Fundamental nature of heat as energy vs. force

4.9 The Thermal Convenience Paradox: Logical Impossibility of Conventional Explanations

Anticipated conventional objections to our fluid dynamic observations contain an irreconcilable logical contradiction that we term the "Thermal Convenience Paradox." Critics attempting to preserve classical fluid dynamics must simultaneously invoke two mutually exclusive thermal states for the same system at the same moment.

4.9.1 The Dual Objection Structure

Objection A - Against stationary milk at high temperature:

"The milk temperature increases from 99.1°C to 100.5°C with flame OFF because the hot pan retains thermal mass and continues transferring residual heat to the liquid."

Implication: The pan's internal environment is a thermal furnace. Metal walls radiate intense heat, warming the bulk liquid and, by thermal necessity, the confined air above it even more intensely (lower thermal mass, faster equilibration).

Objection B - Against explosive vapor release:

"Visible steam suddenly appears upon flame extinction because the air above the liquid rapidly cools, causing vapor condensation."

Implication: The pan's internal environment undergoes instantaneous thermal collapse. The confined air experiences rapid cooling sufficient to shift phase equilibrium within 0.2 seconds.

4.9.2 The Logical Contradiction

These objections cannot coexist. The pan cannot simultaneously be:

1. **Hot enough** to elevate 200g of liquid by 1.4°C over 13 seconds (Objection A)

2. **Cold enough** to thermally collapse the air space in 0.2 seconds (Objection B)

The mathematical impossibility becomes clear when we consider the thermal hierarchy:

If Objection A is true:

The pan retains sufficient thermal energy to heat the bulk liquid:

$$Q_{\text{residual}} = mc_p\Delta T = (0.2)(3900)(1.4) = 1092 \text{ J over 13 seconds} \tag{39}$$

The confined air, having vastly lower thermal mass than the liquid, and being in direct contact with both the hot liquid surface ($T \approx 100$°C) and hot metal walls, should equilibrate to an even higher temperature, not cool. With convection ceased, this air space becomes a stagnant thermal trap, accumulating heat from both boundaries.

The claim that this same air cools in 0.2 seconds contradicts the premise that the pan retains heating capacity.

If Objection B is true:

The air cools rapidly enough to cause condensation. But the pan still retains the metal walls at ≈100°C (as evidenced by continued liquid heating). Air trapped between a 100°C liquid surface below and 100°C metal above cannot cool—it is thermally imprisoned between two heat sources.

The cooling air hypothesis requires the pan to have lost its thermal capacity, contradicting Objection A.

4.9.3 Experimental Evidence of the Contradiction

The chalk + detergent experiment (Video S4) definitively refutes the air cooling hypothesis:

- Response time: $\Delta t < 0.2$ seconds
- Phenomenon occurs at multiple vertical positions simultaneously
- Effect persists across varied atmospheric conditions

If air cooling were responsible, we should observe:

1. Gradient response: top surface affected first, propagating downward
2. Time delay: several seconds for thermal diffusion through air layer
3. Sensitivity to ambient humidity and pressure

None are observed. The response is instantaneous, uniform, and gradient-dependent.

4.9.4 Resolution Through Gradient Field Theory

Our framework requires no contradictory assumptions. A single mechanistic principle explains all observations:

During gradient presence (flame ON):

- Gradient field ∇T stores energy in organized structure
- Field performs mechanical work (convection, vapor retention)
- Energy partitioned: $\dot{Q}_{\text{input}} = \dot{W}_{\text{mechanical}} + \dot{W}_{\text{storage}}$
- Bulk temperature remains lower as energy diverted to work

Upon gradient collapse (flame OFF):

- Mechanical work ceases instantaneously (convection stops)

- Stored gradient energy thermalizes (temperature rises)
- Vapor trap releases (explosive plume)
- All phenomena derive from single cause: field collapse

The framework is internally consistent:

1. Pan thermal mass explains continued heating: consistent with gradient discharge
2. Instantaneous response explains vapor release: consistent with field collapse
3. No thermal paradox: same mechanism explains both observations

4.9.5 Theoretical Implications

This logical contradiction reveals a deeper issue with classical fluid dynamic formulation. The conventional framework lacks the conceptual vocabulary to describe gradient-dominated flow regimes. Attempting to force-fit gradient phenomena into equilibrium convection theory generates logical contradictions.

The resolution requires acknowledging that thermal gradients constitute a distinct fluid dynamic variable:

$$\text{Flow State} = (\vec{v}, T, P, \rho, \nabla T) \tag{40}$$

Classical fluid dynamics represents the limiting case where $|\nabla T| \to 0$. Our observations probe the opposite limit where $|\nabla T| \gg T/L_{\text{system}}$.

The Thermal Convenience Paradox demonstrates that conventional explanations fail not merely quantitatively, but logically. Any theoretical framework that must invoke contradictory thermal states to explain sequential observations is internally inconsistent and must be revised.

5 Alternative Explanations Considered

5.1 Capillary Action

Cannot explain 15 cm vertical rise in millimeter-scale surface irregularities. Capillary height:

$$h_{\text{cap}} = \frac{2\gamma \cos\theta}{\rho g r} \approx 1 \text{ cm} \tag{41}$$

5.2 Vapor Pressure Pumping

Would cause downward drainage as expanding vapor escapes upward, opposite to observed continuous upward liquid flow.

5.3 Thermal Expansion

Insufficient to drive continuous flow against gravity. Expansion coefficient:

$$\alpha = \frac{1}{V}\frac{\partial V}{\partial T} \approx 2 \times 10^{-4} \text{ K}^{-1} \tag{42}$$

5.4 Electromagnetic Fields

No external fields applied. Flame produces negligible EM radiation at frequencies relevant to bulk matter motion.

5.5 Experimental Artifacts

- Pan slope verified mechanically pre/post-heating (rolling sphere test)
- Multiple trials with consistent results ($n = 15$)
- Multiple fluid types tested (water, milk, herbal infusions, lipids)
- Effect persists across different heat source types

6 Future Experimental Directions

6.1 Quantitative Force Measurements

- Direct measurement of thermal force magnitude vs. gradient

- Force spectroscopy at various distances from heat source
- Material-dependent thermal force constants k_T

6.2 Microgravity Experiments

Eliminate gravitational component to isolate pure thermal force:

$$\vec{F}_{\text{total}} = \vec{F}_{\text{thermal}} \quad (F_g = 0) \tag{43}$$

6.3 Controlled Gradient Geometries

- Radial gradients (point source)
- Linear gradients (planar heating)
- Complex gradient topologies

6.4 Molecular Dynamics Simulations

Computational investigation of molecular-scale mechanisms underlying macroscopic thermal attraction.

7 Conclusion

We have presented systematic experimental evidence that active thermal gradients produce fluid behavior fundamentally different from classical predictions. Matter flows upward against gravity toward heat sources, convection depends on gradient rather than temperature, and vapor release occurs far below standard boiling points at 30.1°C—69.9°C below the equilibrium boiling point.

The milk convection experiments provide definitive evidence that fluid motion depends on gradient magnitude $|\nabla T|$ rather than thermal state T, with sub-second response times precluding energy-based explanations. The complete decorrelation between thermal energy content (maximum at 100.8°C with zero motion) and convective work (maximum at 99.1°C) directly contradicts classical formulations.

These observations suggest that thermal gradients generate directional attractive forces not accounted for in standard fluid dynamics or thermodynamic theory. The dissociation between thermal state (T) and thermal flux (∇T) challenges foundational assumptions about heat transfer mechanisms.

The tripartite energy framework ($\dot{Q}_{\text{external}}$, U_{internal}, U_{bulk}) explains apparent violations of instantaneous energy conservation through gradient-mediated internal storage, providing a kitchen thermodynamics understanding of everyday thermal phenomena.

While these results have profound implications for thermodynamic theory [6, 7], they represent empirical observations that demand explanation regardless of theoretical preferences. We invite the fluid dynamics and thermodynamics communities to attempt replication and develop theoretical frameworks accounting for these phenomena.

Supplementary Materials

- Video S1: Iceberg experiment - upward water migration (25 min)
- Video S2: Lipid climbing and vectorial split (25 min time-lapse)
- Video S3: Milk boiling cycles - flame on/off (4 cycles) - **Complete temporal documentation**
- Video S4: High-speed flame extinction - steam release (240 fps)
- Video S5: Extreme low-temperature vapor release at 30.1°C
- Video S6: Microscopic particle thermal attraction

All supplementary materials available at: [Zenodo DOI to be assigned]

Data Availability

All experimental protocols, high-speed video recordings, temperature logs, and raw data will be made available upon publication.

Acknowledgments

The author thanks the fluid dynamics and thermodynamics communities for consideration of these experimental findings.

References

[1] De Gennes, P.G., Brochard-Wyart, F., & Quéré, D. (2013). *Capillarity and Wetting Phenomena: Drops, Bubbles, Pearls, Waves*. Springer.

[2] Marangoni, C. (1865). Sull'espansione delle gocce liquide. *Nuovo Cimento*, 2(56), 239-273.

[3] Clausius, R. (1865). *The Mechanical Theory of Heat*. London: John van Voorst.

[4] Scriven, L.E., & Sternling, C.V. (1960). The Marangoni Effects. *Nature*, 187, 186-188.

[5] Prigogine, I. (1978). Time, Structure, and Fluctuations. *Science*, 201(4358), 777-785.

[6] Boltzmann, L. (1877). Über die Beziehung zwischen dem zweiten Hauptsatze der mechanischen Wärmetheorie und der Wahrscheinlichkeitsrechnung. *Wiener Berichte*, 76, 373-435.

[7] Planck, M. (1926). *Treatise on Thermodynamics*. Dover Publications.

[8] Maxwell, J.C. (1871). *Theory of Heat*. Longmans, Green, and Co.