The seventh ternary density Hales–Jewett number

P. Graham
https://orcid.org/0009-0007-8493-1505

October 19, 2026

Abstract

For a word $w \in \{0,1,2,*\}^n$ with at least one wildcard, its combinatorial line is $\{w(0), w(1), w(2)\}$, where $w(a)$ replaces every wildcard by a. Let $c_{n,3}$ be the maximum size of a subset of $\{0,1,2\}^n$ containing no line. We give a computer-assisted proof that $c_{7,3} = 1302$, confirming D.H.J. Polymath's tentative 2010 conjecture. Polymath's construction supplies the lower bound; the contribution is the unrestricted upper bound. Bounds on lower-dimensional slices and one alphabet permutation acting in every coordinate reduce the exclusion of a 1303-point set to nineteen propositional unsatisfiability obligations: three auxiliary bounds and sixteen slice cases. Executable audits check every encoding clause under explicit counting interpretations. All nineteen LRAT refutations were checked with FRAT-rs; eleven also with the formally verified checker cake_lpr. The original computation additionally checked all nineteen with drat-trim. The reduction and encoding soundness remain written arguments, rather than a formalization of the whole theorem. The accompanying programs regenerate the refutations through bounded pipes.

The proofs, the search code, the certificate checkers and the text were produced by AI models (from OpenAI and Anthropic), working under the author's direction. The result has not been reviewed by human experts.

1 Introduction

The density Hales–Jewett theorem of Furstenberg and Katznelson [1], with a later combinatorial proof by Polymath [2], concerns large subsets of word spaces and the combinatorial lines they must contain. Its finite extremal values are not determined by the asymptotic theorem. For a fixed dimension n, write $c_{n,3}$ for the maximum size of a line-free subset of $\{0,1,2\}^n$. D.H.J. Polymath determined these numbers through dimension six and supplied a 1302-point construction in dimension seven. Section 2 of its 2010 paper calls this the bound "which we tentatively conjecture to be the correct bound" [3]. OEIS A156989 currently lists the established terms through dimension six, ending at 450 [4]. Splitting into slices immediately gives the elementary upper bound $3 \cdot 450 = 1350$. We use this as a benchmark, without asserting that it was the strongest previously recorded bound: some historical tables could not be recovered. A search of the primary paper, OEIS, recent citing works and current postings through October 20, 2026 found no earlier unrestricted settlement. Unrecovered historical wiki, table, spreadsheet and blog sources leave this a likely-new assessment, not an exhaustive priority claim.

Theorem 1. *Every subset of $\{0,1,2\}^7$ with no combinatorial line has at most 1302 points, and a line-free subset of size 1302 exists. Equivalently, $c_{7,3} = 1302$.*

The contribution claimed here is the upper bound for arbitrary subsets. The matching construction and the conjectured value belong to Polymath. The upper bound applies to every subset, without a symmetry assumption on that subset. The argument uses satisfiability computations after elementary counting reductions. The complete theorem is computer-assisted, with no whole-theorem Lean verification. FRAT-rs checked all nineteen propositional refutations, cake_lpr checked eleven, and the original computation used drat-trim on all nineteen; the passage from sets to formulas remains a written argument, backed by independent semantic audits.

The trusted base consists of the written cap induction and global symmetry argument; soundness of the set-to-formula counting interpretation, checked exhaustively clause by clause; the proof checkers within the coverage just stated; and the exhaustive lower-bound witness checks. In the public streaming replay, the very same audited CNF file is read by the solver and all checkers. No receipt or solver verdict substitutes for a refutation check. The small-dimensional upper caps are proved anew by the three auxiliary refutations, not imported as unchecked premises from Polymath.

Recent work addresses adjacent questions: Mouhib's symmetric and one-weight colouring bounds [5], Conlon–Farnsworth–Robertson's colouring questions [6], Shelah's general bounds [7], and Dahia's formalization of the general density theorem [8]. These are different endpoints and are not premises of this proof. This result determines one finite extremal value, not a new proof or improved general bound for the density theorem.

2 Lines, caps and symmetry

Any line-free set of at least 1303 points has a line-free 1303-point subset. Our formulas directly exclude the stronger condition of size at least 1303.

A wildcard pattern is a word in $\{0, 1, 2, *\}^n$ with at least one wildcard. Its line consists of $w(0), w(1), w(2)$, replacing all wildcards simultaneously. There are $4^n - 3^n$ lines; thus the seven-dimensional audit checks 14,197 lines. A coordinate subcube is obtained by fixing some coordinates. A line-free set meets a d-dimensional subcube in at most $c_{d,3}$ points. Splitting on one coordinate gives

$$c_{d,3} \leq 3c_{d-1,3}. \tag{1}$$

For brevity put $c_d = c_{d,3}$ and $c_0 = 1$.

Lemma 1. *A single permutation π of the alphabet, applied simultaneously in every coordinate, preserves lines, line-freeness, cardinality and coordinate-subcube bounds. It permutes the three first-coordinate slice sizes arbitrarily.*

Proof. Apply π to each constant of a wildcard pattern, leaving its wildcard positions unchanged. The image of $w(a)$ is the resulting pattern evaluated at $\pi(a)$. Since π is bijective, the image of the three-point line is exactly a line. The same map is bijective on all words and carries coordinate subcubes to subcubes of the same dimension. A first-coordinate slice labelled a becomes the slice labelled $\pi(a)$. Choosing π to send the three slice labels to their order by size proves the last assertion, including ties. □

Permuting symbols in the first coordinate alone is not the symmetry used here. For example, swapping 0 and 1 in that coordinate sends the line $\{00, 11, 22\}$ to $\{10, 01, 22\}$, which is not a combinatorial line. Earlier proof descriptions used this invalid local explanation. Lemma 1 corrects it; the case formulas and the case cover are unchanged.

3 The finite reduction

The three auxiliary obligations are denoted L_1, L_4, L_5. They exclude line-free sets of sizes at least 3, 53 and 151 in dimensions 1, 4 and 5, respectively. Their formula inputs are:

Obligation	Dimension	Required size	Proper-subcube caps used
L_1	1	3	none
L_4	4	53	$2, 6, 18$
L_5	5	151	$2, 6, 18, 52$

The refutations establish $c_1 \le 2$, then $c_4 \le 52$, then $c_5 \le 150$, in that order. Between them, (1) establishes $c_2 \le 6$, $c_3 \le 18$ and $c_6 \le 450$. There is no use of a cap of the dimension currently being proved. Thus no published small-dimensional optimality result is needed as an unchecked premise.

Suppose a line-free $A \subseteq \{0,1,2\}^7$ has $|A| \ge 1303$. Write $a_v = |A \cap \{x_1 = v\}|$. Lemma 1 permits $a_0 \ge a_1 \ge a_2$. Since each slice is six-dimensional,

$$435 = \lceil 1303/3 \rceil \le a_0 \le 450.$$

For each integer s in this interval the main formula encodes line-freeness, size at least 1303, the proper-subcube caps $2, 6, 18, 52, 150, 450$, and

$$a_0 = s \ge a_1 \ge a_2.$$

All sixteen formulas are unsatisfiable. Every counterexample to the upper bound would, after the global relabelling, satisfy one of them. Their refutations therefore imply $c_7 \le 1302$.

4 Encoding and independent semantic audit

Point $x = (x_1, \dots, x_n)$ has variable index $1 + \sum_i x_i 3^{n-i}$, interpreted as $x \in A$. Every line contributes the clause consisting of its three negated point variables. Hierarchical unary totalizers count points in coordinate subcubes and in intermediate unions of two child subcubes. Auxiliary variable (S, j) means $|A \cap S| \ge j$, where S is a product of nonempty subsets of the alphabet. Thresholds are truncated only beyond an available cap; the first forbidden threshold is explicitly false. A root threshold forces $|A| \ge K$. Units force $a_0 = s$ and threshold implications force the slice order. The encoder and independent auditor are supplied as `checks/encode.py` and `checks/reduction.py`.

For clarity, the audit proves the direction required for an upper bound: every hypothetical counterexample extends to a satisfying assignment by the stated threshold interpretation. It does not rely on a solver's interpretation of auxiliary variables. For a box $S = M_1 \times \cdots \times M_n$, let d be its number of unrestricted coordinates. If $d < n$, disjoint coordinate subcubes give the valid bound

$$U(S) = c_d \prod_{i:\,|M_i|<3} |M_i|.$$

For the entire ambient cube use the trivial bound 3^n, so the current claim is never its own premise. The empty set has bound zero.

The auditor recognizes every clause as one of the following valid statements under that interpretation.

1. A line clause forbids its three points together.

2. An upward counting clause has the form $(S,k) \vee \bigvee_r \neg(P_r, j_r)$, where the P_r are disjoint subsets of S and $\sum_r j_r \geq k$. Its validity follows by adding counts.

3. A downward counting clause has the form $\neg(S,k) \vee \bigvee_r (P_r, j_r)$. The P_r are disjoint subsets of S and the remainder is empty or a box. When all positive literals are false,

$$|A \cap S| \leq \sum_r \min(j_r - 1, U(P_r)) + U(S \setminus \bigcup_r P_r) \leq k - 1,$$

which proves the clause.

4. A cap unit $\neg(S,k)$ has $U(S) \leq k-1$.

5. A premise clause follows from $|A| \geq K$, $a_0 = s \geq a_1 \geq a_2$: these include applicable positive root and slice thresholds, negative slice thresholds beyond s, and order implications $\neg(a_b \geq t) \vee (a_a \geq t')$ for $a < b$ and $t' \leq t$.

The product sets and thresholds are read from an encoder-generated semantic map, but the auditor independently checks point indexing, set inclusion, disjointness, bounds and every clause. The map supplies meanings, not evidence of their validity. The audit also checks all line clauses are present, the size and profile premises are present, header counts agree, and the exact integer case cover is complete. For the main formulas the rule counts are 14,197 line clauses, 940,716 upward clauses, 940,644 downward clauses, 765 cap clauses and 903 premise clauses (902 at $s = 450$, where a cap already supplies one unit).

The logical dependency is therefore explicit: the written validity of these rules and the cap induction, the exhaustive clause audits, and checked unsatisfiability together rule out each set-theoretic counterexample. These steps have not been formalized together in a proof assistant.

5 Refutations and the lower bound

The exact tool revisions and build commands are in `checks/README.md`. CaDiCaL generates text LRAT proofs. The accepted computation regenerated all 19 refutations and checked every one with FRAT-rs's `lratchk`; cake_lpr additionally checked the three auxiliary obligations and the eight main cases $s = 443, \ldots, 450$. The original computation used drat-trim's `lrat-check` on all 19, together with cake_lpr on those 11 or a separate streaming checker on the other eight. This is not a claim that cake_lpr checked all 19.

The eight obligations not replayed with cake_lpr are the largest main cases, $s = 435, \ldots, 442$. Its heap usage grows with proof additions; the fixed-heap runs do not cover these proofs.

The public replay sends the generated stream to FRAT-rs and the supplied drat-trim checker, with cake_lpr also enabled for those eleven obligations. An UNSAT exit code alone never passes: each applicable checker must accept an empty-clause derivation. The large proof streams are not distributed. Full regeneration produces about 1.56 TB of text, consumed through pipes; the accepted independent computation used about 64 solver CPU-hours, 9.7 FRAT-rs CPU-hours and 0.9 cake_lpr CPU-hours, plus encoding and audits. A fresh public run also pays for drat-trim checking. The slowest case took about 21.4 hours elapsed. These are measured costs, not a promise for other hardware. The quick check audits the reduction and the witness and runs a small proof with corruption controls; it does not replay the sixteen main refutations.

For the lower bound, the supplied 1302 words (written over $\{1,2,3\}$) are a realization of Polymath's construction [3, Section 2]. They consist of all words with letter-count triples

$$034, 043, 124, 142, 223, 232, 250, 304,$$
$$313, 331, 340, 412, 421, 502.$$

Their multinomial weights sum to 1302. Two independently written witness checkers verify distinctness, cardinality and absence of all 14,197 combinatorial lines. The data is a transcription of the published construction, not a new lower-bound construction. Together with the upper bound this proves Theorem 1.

Availability and licence. The accompanying bundle contains the manuscript source, the explicit witness, the formula specification, the encoder, semantic and symmetry audits, witness checkers, a standard LRAT checker and instructions for pinned external tools. Large proofs are regenerated. The work is licensed under Apache-2.0; the bundled drat-trim checker retains its MIT notice. See `THIRD_PARTY.md` for dependencies and attribution.

References

[1] H. Furstenberg and Y. Katznelson, *A density version of the Hales–Jewett theorem*, J. Anal. Math. 57 (1991), 64–119.

[2] D.H.J. Polymath, *A new proof of the density Hales–Jewett theorem*, Ann. of Math. 175 (2012), 1283–1327, doi:10.4007/annals.2012.175.3.6.

[3] D.H.J. Polymath, *Density Hales–Jewett and Moser numbers*, in An Irregular Mind, Bolyai Society Mathematical Studies 21, Springer, 2010, arXiv:1002.0374; doi:10.1007/978-3-642-14444-8_22.

[4] OEIS Foundation, A156989: density Hales–Jewett numbers for an alphabet of size 3.

[5] Y. Mouhib, *Lower Bounds for the Hales–Jewett Numbers via Symmetric and One-Weight Colorings*, arXiv:2606.22155v2, 2026.

[6] N. Conlon, N. Farnsworth and A. Robertson, *On Hales–Jewett and Related Numbers*, arXiv:2607.18111, 2026.

[7] S. Shelah, *HJ numbers revisited*, arXiv:2607.14732v3, 2026.

[8] G. Dahia, *DensityHalesJewett*, formalization of the Dodos–Kanellopoulos–Tyros proof, `https://github.com/gdahia/DensityHalesJewett`, 2026.