The Thomson problem for N = 10

P. Graham
https://orcid.org/0009-0007-8493-1505

October 22, 2026

Abstract

Thomson's problem, originating in his 1904 atomic model, asks for the least Coulomb energy of points on a sphere. For $N = 10$ distinct points on the unit sphere in $\mathbb{R}^3$, we prove that the energy $\sum_{i<j} \|x_i - x_j\|^{-1}$ is minimized uniquely, up to orthogonal maps and relabeling, by the D_{4d} bicapped square antiprism. Its height parameter is the distance of either square from the equatorial plane. The chosen height is the unique critical point of the energy in a specified exact interval. Adapting the nine-point proof design, the proof first confines the inner products of every competitor of no greater energy to seven small intervals. Classifying the possible interval patterns and aligning the configurations then reduces the problem to a quantitative local energy inequality that forces equality with the candidate. The computer-assisted proof is formally verified in Lean.

The proofs, the Lean formalization, the certificate checkers and the text were produced by AI models (from OpenAI and Anthropic), working under the author's direction.

1 Introduction

Thomson's problem asks for the least Coulomb energy of N distinct unit vectors in three dimensions. The problem originates in Thomson's 1904 atomic model [1]. For $N = 10$ charges, the familiar candidate has two poles and two squares at opposite heights, rotated by $\pi/4$ relative to one another (Figure 1). Their height is approximately 0.4226874 and the energy is approximately 32.71694946. We prove that this D_{4d} bicapped square antiprism is the unique global minimizer, up to orthogonal maps and relabeling; Theorem 1 gives the exact statement. No symmetry is assumed of competitors.

Optimizing within this one-parameter family does not settle the global problem: competitors may have no symmetry. The main task is to localize all such competitors. The two-point bound used here is only about 32.66696, below the candidate energy, so it cannot establish optimality. Its role is to separate points. A three-point bound then puts every competitor's energy above the candidate energy minus 10^{-30} whenever all distinct pair inner products are at most 13/20. The hard step is to turn this small energy error into control of every competitor's geometry. The window classification does this; the local energy inequality removes the error entirely.

The proof has four steps. First, a two-point bound separates distinct points, after which a polynomial lower bound for pair energy and a three-point energy estimate force all pairwise inner products of a competitor of no greater energy into seven small intervals around the candidate's inner products. Second, a finite classification shows that these intervals can occur on the pairs in only the candidate's pattern, up to relabeling. Third, an orthogonal alignment converts this control of inner products into a bound on the point displacements. Finally, the height equation and symmetry give full spherical stationarity, and a positive second variation, together with a bound

on the higher-order error, makes every sufficiently small nonzero aligned displacement increase the energy. This last step turns approximate localization into exact equality.

The argument builds directly on the nine-point proof published by Aravind Karthigeyan, which Anthropic's Opus 5.5 and OpenAI's GPT-6 Astra generated under his direction and which, unreviewed, he released requesting external review [14], including substantial reuse of its MIT-licensed Lean code. For the ten-point geometry we supply new energy, window-classification and Hessian certificates. The formal proof covers optimality, every equality case and the fact that the candidate consists of distinct unit vectors. To our knowledge no earlier rigorous proof for $N = 10$ exists, and we make no priority claim. Concurrent ten-point Lean proofs establish local minimality and a long-chord condition [8]; neither asserts global optimality. The prior exact cases and these concurrent results are described in Section 5. Neither the present work nor the nine-point work is yet peer-reviewed. The theorem includes uniqueness, and its Lean verification is a formal proof. Being the first rigorous proof or the first formal proof is a historical claim, subject to the qualification just stated.

What is new beyond nine points. The proof follows the design of the nine-point project step by step and reuses most of its Lean code: the form of the statement, the three-point bound with its certificate checks, the Bernstein checks of one-variable inequalities, the interval tests of the window classification, the orthogonal alignment, the higher-order error bound and the interval method for the Hessian. As it stood, that machinery did not reach ten points in three places. First, a three-point certificate belongs to one configuration, and we needed one that is sharp at the antiprism, so that it misses the candidate energy by less than 10^{-30}. Second, the nine-point classification excludes inner products near -1 by a Gram-determinant estimate and works with six windows. Here the poles are antipodal, so -1 is a genuine seventh window, and the two poles with two opposite points of a square lie in one plane through the origin. Every triangle among those four points is then degenerate, so the nine-point proof that four points always contain an anchor triple fails; a counting argument over five points replaces it (Section 3). Third, the local bound certifies one matrix. The ten-point Hessian is 30×30, and its smallest eigenvalue transverse to rotations is about 0.096, against about 0.144 for nine points, so the shift, the radius and the interval data are new, as is the stationarity calculation for the D_{4d} symmetry. The first of the three was the hardest. A numerical semidefinite solution is far less accurate than 10^{-30}; our certificate rounds, to rationals with denominator 2^{200}, an exact three-point certificate that is sharp for ten points, whose one-variable polynomial touches the pair potential at -1 and at the six other candidate inner products.

Section 2 defines the candidate and states the theorem. Section 3 proves the two global localization steps; Section 4 completes the argument by alignment and local rigidity. The separate numerical energy enclosure follows the theorem, and Section 5 records the prior and concurrent work. Appendix A supplies the exact parameter interval, formal declarations, certificate sizes and checking requirements. The text explains the mathematical reductions; the accompanying Lean proof checks the rational inequalities and finite classifications they use.

2 Statement and candidate

This section specifies the configuration and the exact optimization problem. Every competitor consists of ten distinct unit vectors. For an injective map $x : \{0, \dots, 9\} \to S^2 \subset \mathbb{R}^3$, put

$$\mathcal{E}(x) = \sum_{0 \le i < j \le 9} \frac{1}{\|x_i - x_j\|}.$$




Figure 1: The candidate $p(h_*)$. Left: a side view of the ten points and the edges of their convex hull, hidden edges dashed. The poles p_0, p_1 are red, the upper square $p_2, \dots, p_5$ at height $h = h_*$ is blue, and the lower square $p_6, \dots, p_9$ at height $-h$ is orange; the dashed ellipses are the equator and the two circles of latitude. Right: the view from above, in which the lower square (dashed) is the upper square rotated by $\pi/4$. A label i marks p_i. The figure is drawn from the Lean definition of the candidate by `checks/draw_antiprism.py`, which checks that the drawn points are the candidate's; it is not part of the proof.

We call two configurations congruent when one is obtained from the other by an element of $O(3)$ and a permutation of the ten indices; reflections are allowed.

For $0 < h < 1$, write $r = \sqrt{1 - h^2}$ and $q = r/\sqrt{2}$. Define $p(h)$ by

$$
\begin{aligned}
&p_0 = (0, 0, 1), && p_1 = (0, 0, -1), && && \\
&p_2 = (r, 0, h), && p_3 = (0, r, h), && p_4 = (-r, 0, h), && p_5 = (0, -r, h), \\
&p_6 = (q, q, -h), && p_7 = (-q, q, -h), && p_8 = (-q, -q, -h), && p_9 = (q, -q, -h).
\end{aligned}
$$

These coordinates are preserved by rotation through $\pi/4$ about the polar axis followed by reflection in the equatorial plane, and by reflection in the vertical plane through p_2 and the poles. The 16-element group generated by these maps is conventionally denoted D_{4d}.

The 45 unordered pairs consist of one pole pair, eight neighboring and four opposite pairs within the squares, two classes of eight pairs between the squares, and sixteen pole–square pairs. Grouping them gives

$$
\begin{aligned}
F(h) = \frac{1}{2} + \frac{4\sqrt{2} + 2}{\sqrt{1 - h^2}} + \frac{8}{\sqrt{2 - \sqrt{2} + (2 + \sqrt{2})h^2}} + \frac{8}{\sqrt{2 + \sqrt{2} + (2 - \sqrt{2})h^2}} \\
+ \frac{8}{\sqrt{2 - 2h}} + \frac{8}{\sqrt{2 + 2h}}.
\end{aligned} \tag{1}
$$

The terms appear in that order: pole pair, pairs within the squares, the two between-square classes, and the two pole–square classes. Write $S(h) = F'(h)$ for the derivative of this energy with respect

to height. Its explicit formula is given in Appendix A.1. Let $[a, b]$ be the exact rational interval specified in Appendix A.1. It satisfies $0 < a < b < 1$ and $b - a = 1/(2D) < 2^{-300}$, where the integer denominator D is given there. For orientation, this narrow interval lies inside

$$[0.42268742404398, 0.42268742404400].$$

The derivative S is continuous and strictly increasing on $[a, b]$, and its endpoint signs satisfy $S(a) < 0 < S(b)$. The intermediate value theorem and strict monotonicity therefore give a unique zero in this interval. Define h_* by $S(h_*) = 0$ and $h_* \in [a, b]$. Its decimal begins $0.422687424043986\ldots$. The root existence and uniqueness in this specified interval are proved in Lean. This interval selects the candidate; the global theorem still compares it with every admissible configuration, including all other heights.

Theorem 1. *The parameter h_* exists and is unique in $[a, b]$. The configuration $p = p(h_*)$ consists of ten distinct points on S^2. For every injective $x : \{0, \ldots, 9\} \to S^2$,*

$$\mathcal{E}(p) \le \mathcal{E}(x), \qquad \mathcal{E}(x) = \mathcal{E}(p) \quad \Longleftrightarrow \quad \exists Q \in O(3),\ \exists \sigma \in S_{10},\ x_i = Q p_{\sigma(i)} \text{ for all } i.$$

Here and below, admissibility means precisely unit norm and injectivity. The candidate is admissible: in the coordinates above, unit norm follows from $r^2 = 1 - h_*^2$ and $2q^2 = r^2$. The two poles and the two squares have distinct heights; within each square the four planar coordinates are distinct because $r, q > 0$. Thus the ten points are distinct. Appendix A.2 identifies the formal statements of both the optimization theorem and this candidate-validity assertion.

The seven distinct inner products of the candidate are

$$-1, \quad h_*, \quad -h_*, \quad h_*^2, \quad 2h_*^2 - 1, \quad \frac{1 - h_*^2}{\sqrt{2}} - h_*^2, \quad -\frac{1 - h_*^2}{\sqrt{2}} - h_*^2.$$

Remark (energy enclosure). A separate rigorous evaluation of the candidate energy gives

$$32.716949460147 < \mathcal{E}(p) < 32.716949460149,$$

so the energy starts $32.71694946\ldots$. This rigorous rational-interval computation is separate from the formal theorem. Appendix A.8 gives the independent Python calculation and its exact inputs.

3 From energy bounds to a common interval pattern

We now show that every admissible competitor of energy at most the candidate's has nearly the same matrix of pairwise inner products, after relabeling. Let $f(t) = (2 - 2t)^{-1/2}$. For unit vectors, the pair energy is $f(\langle x_i, x_j \rangle)$. Throughout this section, $p = p(h_*)$ and x is an arbitrary admissible configuration satisfying $\mathcal{E}(x) \le \mathcal{E}(p)$.

3.1 Energy bounds force seven windows

We bound the energy lost by replacing the pair potential with a polynomial; this loss will force every pair into one of seven short intervals.

Pair separation. Rational interval evaluation gives $\mathcal{E}(p) \le 32717/1000$ (Appendix A.5), so this is also an energy cap for the competitor. Let g be the quartic polynomial specified in Appendix A.3. It lies below f on the whole interval $[-1, 1)$ and has nonnegative coefficients in the Legendre basis. The spherical-harmonic addition identity therefore expresses the sum of $g(\langle x_i, x_j\rangle)$ over all ordered pairs, including the diagonal, as a constant plus nonnegative weighted squares. Each square is the square of the sum, over the ten points, of a fixed spherical harmonic of degree one through four. Subtracting the diagonal contribution and dividing by two gives a lower bound of $13111005315721/401353681050 > 32.666$ for the sum over unordered distinct pairs. If any pair had inner product at least $13/20$, its energy minus g would exceed $51/1000$. All other pair differences are nonnegative. Adding $51/1000$ to the exact polynomial lower bound exceeds $32717/1000$ by more than 0.00096, a contradiction. Thus $\langle x_i, x_j\rangle < 13/20$ for every $i \ne j$.

Three-point bound. The next bound uses a fixed rational polynomial and a finite set of rational matrices; together these data form the *three-point certificate*. Define $m(t) = \sum_{j=0}^{16} c_j t^j - 2^{-130}$, where the coefficients c_j are `hCoeff` in `lean/ThomsonN10/Data.lean`. The downward shift ensures that m lies below f throughout $[-1, 13/20]$. Let U be the fixed rational `certU` in the same file. Interval evaluation of the candidate gives $\mathcal{E}(p) \le U$, where $U \approx 32.716949460147588154555705751969994$. Appendix A.5 gives its exact value.

For three unit vectors x_i, x_j, x_k, use the argument convention $u = \langle x_i, x_j\rangle$, $v = \langle x_i, x_k\rangle$ and $t = \langle x_j, x_k\rangle$. The matrices specify two polynomials K and W. Define the residual polynomial $\mathcal{R}$ by

$$\frac{m(u) + m(v) + m(t)}{3} - \frac{U - 45 \cdot 2^{-130}}{45} = K(u, v, t) + W(u, v, t) + \mathcal{R}(u, v, t).$$

The content of the certificate is threefold. First, the sum of K over all ordered triples of distinct indices is nonnegative for every configuration of ten unit vectors. This is the three-point positivity of Bachoc–Vallentin and Cohn–Woo [17, 18], with the repeated-index corrections specified in Appendix A.4. Second, W is a sum of weighted polynomial squares, and is nonnegative on the *capped Gram domain* $u, v, t \in [-1, 13/20]$, $1 + 2uvt - u^2 - v^2 - t^2 \ge 0$. The six weights are listed in Appendix A.5. Third, the sum $\|\mathcal{R}\|_1$ of the absolute values of the residual's rational coefficients is tiny: $\|\mathcal{R}\|_1 \approx 1.1673222012517146 \cdot 10^{-56}$. Every residual monomial has absolute value at most one on this domain, so this coefficient sum bounds the residual pointwise.

Put $L = 45(2^{-130} + \|\mathcal{R}\|_1)$; numerically, $L \approx 3.306077861687684 \cdot 10^{-38} < 10^{-30}$. These decimals describe the exact rational data, and are not used as proof inputs. Averaging the defining identity over distinct triples and using the two positivity properties gives the following bound.

Lemma 1 (Certified energy bound)**.** *For the fixed polynomial m and constants U, L just specified, every configuration of ten unit vectors with $\langle x_i, x_j\rangle \le 13/20$ for all $i \ne j$ satisfies*

$$\mathcal{E}(p) \le U \le 32717/1000, \qquad U - L \le \sum_{i<j} m(\langle x_i, x_j\rangle), \qquad L < 10^{-30}.$$

The candidate inequality in the display is the separately established interval bound above. For the pair-sum inequality, the unit-vector hypotheses put every triple in the Gram domain, while the pair cap gives $u, v, t \le 13/20$, where all weights of W are nonnegative. Appendix A.5 gives the triple-to-pair averaging and explains the factor 45 in L. The much weaker separation bound used Delsarte–Yudin linear programming; here the error is small enough to force every pair into a candidate window.

For each pair, call the nonnegative difference between its true energy and its polynomial lower bound its *slack*. Their total is at most L, because the energy is at most that of the candidate:

$$\sum_{i<j}(f-m)(\langle x_i,x_j\rangle)=\mathcal{E}(x)-\sum_{i<j}m(\langle x_i,x_j\rangle)\le\mathcal{E}(p)-(U-L)\le L.$$

Slack outside the windows. A *window* is an open interval of radius $\delta=10^{-10}$ centered at one of the seven candidate inner products in Section 2. The slack satisfies $f(t)-m(t)\ge 10^{-30}$ for every $t\in[-1,13/20]$ outside all seven windows. Since $L<10^{-30}$, even one pair outside the windows would exceed the total available slack. Every pair therefore belongs to a window.

3.2 The window pattern is the candidate's pattern

We now classify which assignments of these seven windows to the pairs can arise from ten unit vectors in three dimensions. Number the seven candidate inner products by the colors zero through six, as in the table below. Assign a color to each pair according to the window containing its inner product. For example, the candidate's pole pair receives the color of the window centered at -1. The windows are disjoint, so each pair has just one color.

Color	Candidate inner product	Number of pairs
0	-1	1
1	h_*	8
2	$-h_*$	8
3	h_*^2	8
4	$2h_*^2-1$	4
5	$(1-h_*^2)/\sqrt{2}-h_*^2$	8
6	$-(1-h_*^2)/\sqrt{2}-h_*^2$	8

An *anchor triple* consists of three points whose three-by-three Gram matrix has determinant greater than $1/100000$. Such a triple is linearly independent. For each sorted triple of pair colors, evaluate the three-by-three Gram determinant over rational outer enclosures of the three windows. The resulting *determinant interval* contains every determinant that those colors can realize. An anchor *type* is a sorted triple of pair colors. A type is *eligible* when this interval's upper endpoint is greater than the threshold; every actual anchor triple has an eligible type.

An anchor triple must exist. The determinant intervals show that a triangle with determinant at most the threshold can have only a color permutation of $(0,1,2)$, $(1,1,4)$ or $(2,2,4)$. Each such triangle has exactly one edge of color zero or four. If every triangle among the five configuration points with indices zero through four had one of these types, counting these marked edges over the ten triangles would give ten. But each edge belongs to three triangles, so the same count must be divisible by three. This contradiction supplies an anchor triple. The subsequent enumeration treats all 43 eligible anchor types. Reorder the three anchor points so that the colors of their pairs (first, second), (first, third), (second, third) are nondecreasing. Permuting three vertices permits every permutation of their three edges, so such an order always exists. The node coordinates below use this fixed order. For example, the candidate triple p_2,p_3,p_6 has ordered pair colors $(3,5,5)$.

Each of the remaining seven points determines a *node*: the three colors of its pairs with the three anchor points. The four-by-four Gram determinant of this point and the anchor points vanishes because four vectors lie in three dimensions; interval evaluation removes nodes for which this is impossible.

For two remaining points, form a four-by-four matrix with rows indexed by the three anchor points followed by the first point, and columns indexed by the same three anchor points followed by the second point. Each entry is the inner product of its row-index and column-index vectors. This is their *cross-Gram matrix*. It has rank at most three, so its determinant vanishes. Two nodes are *compatible* with a proposed color between them when interval evaluation of this determinant contains zero. The nodes specify the windows for inner products with the anchor points, and the proposed color specifies the window for the one remaining entry.

Join two nodes when at least one color survives the pair test, retaining the list of all surviving colors on that edge. A *clique* is a set of seven nodes with an edge between every two. The seven remaining points of an actual configuration give such a clique. They give distinct nodes because the pair test excludes all seven colors when the two node indices are equal; that test still allows the two points' inner products with the anchors to vary independently within the same windows.

The enumeration checks every clique in every eligible anchor type. For each clique, its witness is a bijection from the three anchor points and seven nodes to the candidate's ten indices. The witness preserves the anchor and node colors and checks *every surviving color* on each clique edge against the candidate color. Thus no separate choice of edge colors is needed: all colors that an actual configuration could use are forced to agree with the same candidate labeling. The nine witnesses correspond to nine enumerated node subsets with their anchor types; all give the same candidate pattern up to relabeling.

Classification. The finite classification proves that there exists a permutation τ for which

$$|\langle x_{\tau(i)}, x_{\tau(j)}\rangle - \langle p_i, p_j\rangle| \le \delta \quad \text{for all } i, j. \tag{2}$$

Thus each pair has been assigned the same window as the corresponding candidate pair. The permutation τ labels the competitor by the candidate indices. The conclusion fixes an interval pattern and bounds every Gram-matrix entry; it does not yet say that the inner products are equal. The certificate counts and its formal soundness statement are given in Appendix A.5.

4 Alignment and exact local rigidity

This section proves that a competitor of no greater energy satisfying (2) is congruent to the candidate. We first align it with the candidate and then show that any remaining displacement raises energy.

4.1 From inner products to point displacements

We convert the bounds on inner products into a small displacement from the candidate. Among orthogonal maps, choose Q minimizing the sum of squared displacements from p, and put

$$y_i = Q x_{\tau(i)}, \qquad \eta_i = y_i - p_i, \qquad \rho^2 = \sum_i \|\eta_i\|^2.$$

Write $\eta \in \mathbb{R}^{30}$ for the concatenation of the ten displacement vectors η_i. The quantity ρ is their total displacement. For alignment we choose the fixed candidate triple p_2, p_3, p_6 and the corresponding relabeled competitor triple. These need not be the anchor triple used during classification: (2) controls every pair, so it also controls these chosen triples. Aligning the anchor frames—the orthonormal bases obtained by Gram–Schmidt from these triples—controls their coordinates; the inner products with the anchor points then control the other points. The estimates in Appendix A.6

construct an alignment with total displacement at most 10000δ. A minimizing alignment is at least as good, so

$$\rho \le 10000\delta = 10^{-6}.$$

The global and local constants serve different steps. The windows have radius 10^{-10}, and the coordinate estimate loses a factor of 10^4. The resulting radius 10^{-6} is one thousand times smaller than the local energy estimate's permitted radius 10^{-3}.

Minimality of this orthogonal Procrustes alignment—least squares over orthogonal maps—also gives the skew conditions below. Here k, l index the three coordinates, and $\eta_{i,k}$ is coordinate k of the displacement η_i:

$$\sum_i (\eta_{i,k} p_{i,l} - \eta_{i,l} p_{i,k}) = 0 \quad (k, l \in \{1, 2, 3\}). \tag{3}$$

These equations make the displacement orthogonal to infinitesimal rotations. Each displacement also has the radial component $\langle p_i, \eta_i \rangle = -\|\eta_i\|^2/2$, by the exact unit-sphere identity. Thus the displacement itself need not lie in the tangent plane. Both properties enter the energy estimate below.

4.2 Stationarity and the full second variation

We establish stationarity and a positive quadratic bound for every perturbation after removing infinitesimal rotations. The candidate is stationary with respect to arbitrary tangent motions of all ten points. Write ∇_i for the gradient with respect to the three coordinates of point i. Since the energy is invariant under orthogonal maps and relabeling, its gradient transforms under the candidate's symmetries in the same way as the points do. A quarter-turn fixes each pole, so its gradient is vertical. The reflection in the vertical plane through p_2 fixes p_2, so its gradient lies in that plane. After removing the radial direction, the only possible gradient component there is along the motion of p_2 as the height changes.

All eight square points form one orbit under the fixed D_{4d} group, and changing height respects these same symmetries. Their contributions to the height derivative are therefore equal; the poles do not move. Thus $S(h_*)$ is eight times the contribution of p_2. Its vanishing makes that tangential component zero, and symmetry does the same at all eight square points. Every gradient is consequently radial.

More precisely, $\nabla_i \mathcal{E}(p) = \lambda_i p_i$, where $\lambda_i = -\frac{1}{2} \sum_{j \ne i} \|p_i - p_j\|^{-1}$. We call λ_i the radial coefficient; it is the spherical Lagrange multiplier. The gradient is the negative of the physical force. The corresponding formal lemmas are listed in Appendix A.7.

Let $H(h)$ be the 30×30 matrix given by the following blocks at $p(h)$, and abbreviate $H = H(h_*)$. At the stationary candidate, this is the constrained second variation. In the block formula at a general height, the radial coefficient means $\lambda_i(h) = -\frac{1}{2} \sum_{j \ne i} \|p_i(h) - p_j(h)\|^{-1}$; stationarity is asserted only at h_*. For a nonzero pair-difference vector d, the pair Hessian is

$$B(d) = \frac{3dd^{\mathsf{T}}}{\|d\|^5} - \frac{I_3}{\|d\|^3},$$

where the off-diagonal block of H for two different point indices is the negative of $B(p_i - p_j)$. The diagonal block at point i is the sum of $B(p_i - p_j)$ over the other nine points, minus the radial coefficient at that point times the three-by-three identity. These blocks specify the sphere-constrained second variation, including the radial correction. Similarly, let $R(h)$ have as its three columns the standard coordinate-axis infinitesimal rotations of $p(h)$, each with unit angular speed, and abbreviate $R = R(h_*)$.

Matrix bound. The exact matrix bound is

$$H + RR^{\mathsf{T}} - \frac{1}{20} I_{30} \succeq 0. \tag{4}$$

Rational interval arithmetic verifies the same inequality with $H(h)$ and $R(h)$ throughout $[a, b]$, hence at h_*. An invertible change of basis, applied by congruence to the matrix on the left of (4), transforms it into one with strictly positive diagonal entries, each greater than the sum of the absolute values of the other entries in its row. This strict diagonal dominance proves positive definiteness; the inverse change of basis gives (4). Appendix A.7 identifies the checked matrix calculation. For the concatenated displacement η, equation (3) gives $R^{\mathsf{T}}\eta = 0$, and (4) implies $\eta^{\mathsf{T}} H \eta \geq \rho^2/20$. The alignment allows all orthogonal maps; R removes their continuous rotation directions. The estimate covers every nearby point of $(S^2)^{10}$ after orthogonal alignment, including perturbations that break D_{4d} symmetry.

4.3 A finite-radius energy gap

To pass from the second variation to an energy inequality, we bound the higher-order error. For a nonzero pair-difference vector d and its perturbation w, put $\ell = \|d\| > 0$.

Pair remainder bound. For $\|w\| \leq \ell/200$, the algebraic pair estimate is

$$\frac{1}{\|d + w\|} \geq \frac{1}{\ell} - \frac{\langle d, w\rangle}{\ell^3} + \frac{3\langle d, w\rangle^2 - \ell^2\|w\|^2}{2\ell^5} - \frac{51}{50}\frac{\|w\|^3}{\ell^4}. \tag{5}$$

For this second-order Taylor bound with cubic error, put $z = (2\langle d, w\rangle + \|w\|^2)/\ell^2$. The proof bounds $(1 + z)^{-1/2}$ below by $1 - z/2 + 3z^2/8 - 5z^3/16$ on $-1 < z \leq 1$, then applies this bound to the normalized pair displacement. Every candidate pair distance is at least one. For $d = p_i - p_j$, $w = \eta_i - \eta_j$, the geometric bounds are

$$\|\eta_i - \eta_j\| \leq \tfrac{3}{2}\rho, \qquad \sum_{i<j} \|\eta_i - \eta_j\|^3 \leq 15\rho^3.$$

The first follows from $\|\eta_i - \eta_j\| \leq \sqrt{2}\rho \leq \frac{3}{2}\rho$. For the second, the identity for the sum of squared pair differences gives $\sum_{i<j} \|\eta_i - \eta_j\|^2 \leq 10\rho^2$. Multiplying this by the largest pair displacement gives the cubic bound. For $\rho \leq 1/1000$, these satisfy the hypothesis of (5). Summing the error over all 45 pairs bounds it by $(51/50)15\rho^3 = (153/10)\rho^3$.

The linear terms also have a quadratic expression on the sphere. By radial stationarity, their sum is the sum, over points, of the radial coefficient times $\langle p_i, \eta_i\rangle$. The exact sphere identity $\langle p_i, \eta_i\rangle = -\|\eta_i\|^2/2$ makes this minus one half of the sum of the radial coefficients times the squared displacements. Adding the quadratic terms from the pair Hessians therefore gives one half of the constrained quadratic form $\eta^{\mathsf{T}} H \eta$: the radial correction is exactly the one subtracted in the diagonal blocks of H. Consequently, throughout $\rho \leq 1/1000$, one obtains

$$\mathcal{E}(y) - \mathcal{E}(p) \;\geq\; \frac{1}{2}\eta^{\mathsf{T}} H \eta - \frac{153}{10}\rho^3 \;\geq\; \frac{\rho^2}{40} - \frac{153}{10}\rho^3 \;\geq\; \frac{\rho^2}{200}.$$

For every nonzero aligned displacement with $\rho \leq 1/1000$, the final quantity is positive. Since $\mathcal{E}(y) = \mathcal{E}(x) \leq \mathcal{E}(p)$, the inequality forces $\rho = 0$, hence $y = p$ exactly. Every competitor of energy at most $\mathcal{E}(p)$ is therefore congruent to p; all remaining competitors have strictly larger energy. Conversely, orthogonal transformations and permutations preserve energy. This proves both optimality and the equality case in Theorem 1. The candidate itself attains the minimum.

5 Prior and concurrent work

The theorem fits a sequence of exact results for small numbers of points. We record that context, the code and proof design used here, and the concurrent partial results for ten points.

The numerical candidate predates these formal proofs. Erber and Hockney's 1991 numerical study [3] reports ten-point energy 32.71694946. The Cambridge Cluster Database [4] lists D_{4d} symmetry and energy 32.7169494. The separate rigorous enclosure remark in Section 2 agrees with the Erber–Hockney value and refines the database's shorter numerical approximation.

The immediate precedent is the MIT-licensed `thomson-n9-lean` project [14], directed and published by Aravind Karthigeyan. Its README calls it AI-generated, by Anthropic's Opus 5.5 and OpenAI's GPT-6 Astra, with the mathematics, formal code, checks and exposition "developed through AI agents under human direction", and describes it as "a research release requesting external review and fresh-machine reproduction, not a claim of completed peer review or priority". We adapt its global localization and local rigidity argument, including substantial reuse of its formal code. Appendix A.9 identifies the revision, its nine-point theorem and our ten-point changes.

The classical configurations include the antipodal pair, equilateral triangle, regular tetrahedron and regular octahedron. The tetrahedral four-electron configuration appears in Föppl's 1912 stability analysis of electron arrangements [2]; this historical stability result should be distinguished from a global minimum theorem. Yudin's energy bounds, published in Russian in 1992 and in English in 1993, prove the octahedral case [16]. For twelve points, Andreev's 1996 paper proves that the icosahedron maximizes the product of pairwise distances and, as its zbMATH review records, claims the corresponding results for the Coulomb energy [6]; Cohn and Kumar credit him with the icosahedron as the unique minimum [7], and their 2007 universal-optimality theorem, which covers the Coulomb energy, includes the tetrahedron, octahedron and icosahedron. Schwartz proved the five-point triangular bipyramid optimal in a 2010 preprint, published in 2013 [5].

Recent work includes Hung Tran's AI-assisted seven-point Lean proof, published on the Vals AI blog in September 2026: he fixed the two Lean statements and deployed ten agents running Anthropic's Sonnet 5.5, which did the proof work in about 15 hours, and he ran five checks on the result, including a second kernel [9]. Kryvonos, Liehr and Taylor's September 18, 2026 paper [10] proves that the square antiprism is the unique global minimizer for eight points for both Coulomb and logarithmic energy, with a computer-assisted proof verified in Lean. Joseph Tooby-Smith and Alex Zughaid also released a Lean-checked warrant in September 2026 for the eight-point Coulomb problem [11]. Their README describes it as a formal check that the square antiprism solves the eight-point case uniquely, and calls it a complete warrant, not a proof, that is still a work in progress; it does not state uniqueness up to orthogonal maps and relabeling. Alex Zarco (`alejandrozarco`) released AI-produced, unreviewed Lean warrants for two smaller cases in September–October 2026. For five points, a single sharp three-point semidefinite certificate, with a finite enumeration for uniqueness, shows that the triangular bipyramid uniquely minimizes the Coulomb energy, a theorem already known from Schwartz; the Lean formalization states uniqueness up to orthogonal maps and relabeling [12]. His seven-point repository, prepared in late September with its Lean part added on October 1 and chiefly on the logarithmic energy, added on October 8 a second check of Tran's Coulomb theorem with its own certificates and a Lean development that uses Tran's local lemma [13]. We use no code from either. The general methodology belongs to the Delsarte–Yudin theory of energy bounds [15, 16]. The specific three-point lineage is the semidefinite-programming framework of Bachoc and Vallentin [17] and its energy-minimization extension by Cohn and Woo [18]. The immediate design and formal implementation precedent here is the nine-point project.

Concurrent ten-point formalizations maintained by `saberwatchmanga` (formerly `terry2418`), in

the `-lean-finite-certificates` project [8], give two further partial results. Its October 7, 2026 posting proves full-space local minimality for an exactly specified height, with strict increase on a 17-dimensional slice removing rotations. Its October 11, 2026 posting proves that any configuration of energy at most 32.717 has a chord longer than 1.935. These are rigorous partial results; that project expressly makes no claim of global optimality. The present proof uses neither result as a premise and uses no code from that project. Both use the cutoff 32.717. We cannot establish whether our choice was independent or taken from that work; Appendix A.9 explains its mathematical role.

A Formal proof and certificates

This appendix identifies the formal statements, certificates and independent computations behind the preceding argument. The distribution contains a normal Lake project, rational certificates and a checker using Python's standard library. The Lean theorem begins with the definitions of unit sphere, injectivity, Coulomb energy and congruence; no numerical optimization output is treated as an axiom. The final theorem combines the root proof, pair separation, three-point bound, polynomial lower bound, the slack bound outside the seven windows, classification of the interval patterns and quantitative local energy inequality described above. The formal trust base is the Lean kernel and the three axioms printed below, together with the faithful match between its statement and Theorem 1. Paths beginning with `lean/` or `checks/` are relative to the bundle root. Other Lean source paths below are relative to `lean/ThomsonN10/`.

Checking artifacts. The project uses Lean 4.34.1. Its Mathlib revision is

`d13f23b723b8a846827a245b89c10fc7d3f11612`.

Both dependencies are pinned by the included toolchain and manifest. A resumed build using two compiler threads and a partially populated build/dependency cache took about 33 minutes. A full build from an empty cache has not been timed. The build checks the theorem statements and the following `#print axioms` outputs. Any change to an axiom list makes the build fail:

```
'ThomsonN10.thomson_ten_complete' depends on axioms:
[propext, Classical.choice, Quot.sound]
'ThomsonN10.candidate_admissible' depends on axioms:
[propext, Classical.choice, Quot.sound]
'ThomsonN10.Cert.L1.Data.residual_agrees_JSON' depends on axioms:
[propext, Classical.choice, Quot.sound]
```

Neither `native_decide` nor `ofReduceBool` is used in the included Lean proof, including the decompressed certificate source. In particular, no native-computation axiom occurs in the printed transitive dependencies of the final theorem. These are the standard Lean axioms for propositional extensionality, classical choice and quotients.

The standard-library Python checker is an independent implementation that checks the same rational three-point certificate used by Lean. It expands the kernel and weighted-square polynomials from their definitions, checks the factor matrices and all 274 residual coefficients, bounds their total absolute error, and checks the candidate-energy upper bound over the entire height interval. The command `python3 -I checks/check.py` also runs eight corruption tests, all of which are rejected: it changes entries in each kind of factor matrix, a minorant coefficient, the candidate upper bound, row exponents, row exponents together with the residual, a residual entry, and the residual norm. The Python implementation does not check the scalar minorant, the slack bound

outside the windows, exhaustive passage from spherical configurations to interval patterns, window classification, full stationarity, orthogonal alignment or local Hessian argument; those obligations are proved by Lean. Its precise inputs and obligations are documented in `checks/README.md`. Independence here concerns the implementation of the shared certificate checks, not a second independent proof of every step of the global theorem.

A.1 The exact height interval

This subsection supplies the exact root interval and derivative formula. For the rational height interval used in Section 2, let A and D be the integers specified below, with D the denominator. Its endpoints a, b are defined by

$$a = A/D, \qquad b = (2A+1)/(2D),$$

$A = 513213568640798381215999065069135718731748351905698829816800875926548421101580455514066781,$
$D = 2^{283} \cdot 5^7.$

For the explicit stationarity equation, set

$$u_-(h) = 2 - \sqrt{2} + (2+\sqrt{2})h^2, \qquad u_+(h) = 2 + \sqrt{2} + (2-\sqrt{2})h^2.$$

The derivative of the family energy is

$$\begin{aligned} S(h) = {} & \frac{(4\sqrt{2}+2)h}{(1-h^2)\sqrt{1-h^2}} - \frac{8(2+\sqrt{2})h}{u_-(h)\sqrt{u_-(h)}} - \frac{8(2-\sqrt{2})h}{u_+(h)\sqrt{u_+(h)}} \\ & + \frac{8}{(2-2h)\sqrt{2-2h}} - \frac{8}{(2+2h)\sqrt{2+2h}} = F'(h). \end{aligned} \tag{6}$$

The root argument in Section 2 is formalized in `Param/Root.lean`. `Param/Final.lean` identifies those functions and endpoints with `stationarity`, `heightLower` and `heightUpper`. It proves `ParameterExistsUnique_holds`.

The type `CandidateParameter` consists exactly of reals in this interval satisfying $S(h) = 0$. Root uniqueness is asserted in this specified interval, not for all real heights.

A.2 Formal theorem statements

This subsection identifies the declarations that implement the theorem and their precise class of admissible competitors. The optimality and uniqueness assertions, together with parameter existence and uniqueness, form the declaration `ThomsonN10.thomson_ten_complete`. Its type is `ThomsonN10.Challenge.CompleteTarget`.

The other declaration `ThomsonN10.candidate_admissible` proves unit norm and injectivity for every `CandidateParameter`. Theorem 1 combines these two declarations. Each certificate assertion used in the proof is itself proved in Lean. In Lean, a configuration is a map from `Fin 10` to Euclidean three-space, and `IsSphereConfiguration` means exactly that each vector has norm one and the map is injective. The total Lean function uses the field convention $0^{-1} = 0$, so a collision is excluded by the injectivity hypothesis; it is not assigned infinite energy.

The following are the definitions in `Basic.lean`, with the mathematical glyphs typeset:

```
abbrev R3 := EuclideanSpace ℝ (Fin 3)
abbrev Configuration := Fin 10 → R3

def IsSphereConfiguration (x : Configuration) : Prop :=
  (∀ i, ‖x i‖ = 1) ∧ Function.Injective x

noncomputable def coulombEnergy (x : Configuration) : ℝ :=
  ∑ i : Fin 10, ∑ j ∈ Finset.Ioi i, (‖x i - x j‖)^-1

def Congruent (x y : Configuration) : Prop :=
  ∃ (g : R3 ≃ₗᵢ[ℝ] R3) (p : Equiv.Perm (Fin 10)), ∀ i, x i = g (y (p i))
```

Here the isometry-equivalence notation means a bijective real linear isometry, hence an element of $O(3)$. The challenge declarations in `lean/Challenge.lean` are:

```
def ParameterExistsUnique : Prop :=
  ∃! h : ℝ, (heightLower : ℝ) ≤ h ∧ h ≤ (heightUpper : ℝ) ∧ stationarity h = 0

def GlobalOptimality : Prop :=
  ∀ h : CandidateParameter, ∀ x : Configuration,
    IsSphereConfiguration x → coulombEnergy (candidate h) ≤ coulombEnergy x

def GlobalUniqueness : Prop :=
  ∀ h : CandidateParameter, ∀ x : Configuration,
    IsSphereConfiguration x →
    (coulombEnergy x = coulombEnergy (candidate h) ↔ Congruent x (candidate h))

def CompleteTarget : Prop := ParameterExistsUnique ∧ GlobalOptimality ∧ GlobalUniqueness
```

The first component specifies the height; the second is the lower bound; the third gives both directions of the equality case. Candidate validity is supplied by the separate theorem already identified above.

A.3 The separation polynomial

This records the quartic used to exclude too-close pairs before applying the three-point bound. Write P_k for the Legendre polynomial of degree k, normalized by $P_k(1) = 1$. The polynomial is

$$g(t) = \frac{17\,366\,680\,031}{21\,123\,877\,950} + \frac{833\,419\,493\,024}{1\,672\,307\,004\,375}P_1(t) + \frac{1\,760\,733\,078\,976}{7\,023\,689\,418\,375}P_2(t) + \frac{17\,766\,191\,104}{185\,811\,889\,375}P_3(t) + \frac{5\,504\,237\,568}{260\,136\,645\,125}P_4(t).$$

The nonnegative coefficients give the spherical-harmonic sum identity. The inequalities $g(t) \le f(t)$ on $[-1, 1)$ and $f(t) - g(t) > 51/1000$ on $[13/20, 1)$ follow from exact polynomial identities with nonnegative Bernstein coefficients. The identities and harmonic expansion are in `Cert/LP/Data.lean`; the scalar inequalities are in `Cert/LP/Scalar.lean`. The sum bound and separation conclusion are in `Cert/LP/Proof.lean`. The letter g here denotes the polynomial called `h` in those files.

A.4 The three-point kernel polynomials

This subsection specifies K and explains its nonnegative triple sum. Define polynomials $C_n(\alpha, \beta)$ by

$$C_0 = 1, \qquad C_1 = \alpha, \qquad C_{n+2} = 2\alpha C_{n+1} - \beta C_n, \qquad T_k(u, v, t) = C_k(t - uv, (1 - u^2)(1 - v^2)).$$

Here u and v are the inner products with the first, distinguished point, and t is the inner product of the other two points. Fixing that first point and projecting the others onto its perpendicular plane identifies the projected vectors with complex numbers. The recurrence expresses real parts of their powers. Therefore the quadratic sums of T_k, with weights given by products of real point weights, are squared absolute values of complex sums and are nonnegative. This is the spherical three-point positivity used in the proof.

For block k, let $J^{(k)}$ be its supplied factor matrix (the integer entries divided by 2^{200}, with unused entries set to zero in an eleven-by-eleven matrix). Its column polynomials and the polynomial with this distinguished first point are

$$\phi_{kr}(u) = \sum_{i=0}^{10} J^{(k)}_{ir} u^i, \qquad \mathcal{A}_k(u,v,t) = \sum_{r=0}^{10} \phi_{kr}(u)\phi_{kr}(v)T_k(u,v,t).$$

Let $\overline{\mathcal{A}}_k$ be the average of $\mathcal{A}_k$ over all six permutations of its three arguments. The repeated-index correction and full kernel are

$$K_k(u,v,t) = 8\overline{\mathcal{A}}_k(u,v,t) + \overline{\mathcal{A}}_k(u,u,1) + \overline{\mathcal{A}}_k(v,v,1) + \overline{\mathcal{A}}_k(t,t,1) + \tfrac{1}{9}\overline{\mathcal{A}}_k(1,1,1),$$
$$K(u,v,t) = \sum_{k=0}^{9} K_k(u,v,t).$$

The coefficients account for repeated indices in the full triple sum. There are $10 \cdot 9 \cdot 8 = 720$ ordered triples with distinct indices, $10 \cdot 9 = 90$ for each of the three ways exactly two indices coincide, and ten with all indices equal. Divide the full sum by 90: the distinct triple average has coefficient $720/90 = 8$, each repeated-pair average has coefficient one, and the constant has coefficient $10/90 = 1/9$. Averaging K_k over distinct triples gives exactly this normalized full sum. Its nonnegativity therefore gives the required distinct-triple positivity of K. The recurrence and projection argument are in `Cert/SDP/CircleKernel.lean` and `Cert/SDP/ProjectionComplex.lean`; the symmetrization and correction are in `Cert/SDP/FactoredKernels.lean`.

A.5 Certificate data and finite classification

This subsection specifies the rational data and weights used in Lemma 1, then describes the finite classification. The coefficients c_j are `hCoeff` in `lean/ThomsonN10/Data.lean`; `certU` there is the exact rational

$$U = \frac{52\,574\,110\,779\,609\,793\,015\,977\,559\,478\,279\,245\,057\,879\,406\,410\,836\,764\,596\,571\,436}{2^{200}}.$$

These are fixed data, not optimization variables in the theorem. The candidate upper bound is verified by `Cert/CandidateUpper.lean`.

Write $\Delta = 1+2uvt-u^2-v^2-t^2$ for the Gram determinant. Before averaging over permutations of u, v, t, the six weights multiplying polynomial squares are 1, $1+u$, $13/20-u$, Δ, $(1+u)(13/20-u)$ and $\Delta(1+u)(13/20-u)$. Each is used for two blocks, one symmetric and one antisymmetric under interchange of v, t; after squaring and permutation averaging all twelve blocks are nonnegative on the capped Gram domain. These weights are `domainWeight` in `Cert/SDP/CertificateBound.lean`.

Averaging the residual identity over the $10 \cdot 9 \cdot 8$ ordered triples of distinct indices turns its pair term into $\sum_{i<j} m(\langle x_i, x_j\rangle)/45$. This is because each of the 45 unordered pairs occurs equally often in the three positions. The polynomial shift 2^{-130} is included in m, so the reference constant is $(U - 45 \cdot 2^{-130})/45$. Bounding the averaged residual by $\|\mathcal{R}\|_1$ and multiplying by 45 gives

$U - L$. In particular, the shift is included once in the defined loss L. The displayed 10^{-30} bound is conservative: most of L comes from the downward minorant shift, while the residual norm is about 10^{-56}. No exact-sharpness assertion is inferred from these small numbers.

The certificate stores 20 matrices of factors: eight nonzero matrices for the three-point kernels (called moment blocks in the data), and twelve for the weighted squares (called sum-of-squares blocks). It has 42,950 integer entries and 274 residual coefficients; two zero moment blocks are omitted. Its compressed JSON file is 1,246,399 bytes and expands to 2,670,249 bytes. Expanding these factors and bounding the coefficient sum are exact rational calculations. The identity is `Cert.L1.l1_hidentity`; its summation is `Cert.SDP.factored_certificate_pair_bound`.

In the data, a *norm test* is the four-by-four Gram-determinant interval test for one proposed node; an *extension record* stores one retained node. A *pair-color test* checks the cross-Gram determinant for two nodes and a proposed pair color. A *seven-clique witness* is the color-preserving candidate labeling for seven compatible nodes and the three anchor points.

The finite certificate treats 84 sorted anchor-color triples, of which 43 are eligible. It contains 14,749 norm tests, 79 extension records and 1,960 pair-color tests. Its nine seven-clique witnesses each give a bijective relabeling to the same reference interval pattern. The count is of witnesses in the enumeration, not of incongruent ten-point configurations.

The interval-pattern conclusion is the theorem `Windows.WindowClassification_holds`.

Scalar checks and data formats. The minorant and the slack bound outside the windows are certified by subdividing rational intervals and checking nonnegative Bernstein coefficients of integer polynomials. The substitution $t = 1 - 2y^2$, with $y = \sqrt{(1-t)/2} > 0$, turns the reciprocal-square-root inequalities into polynomial inequalities after multiplication by positive denominators. The soundness proof is in `Scalar/Checker.lean`; `Scalar/Bridge.lean` carries out the substitution and `Scalar/Exterior.lean` proves the slack bound outside the windows.

The compressed JSON file is input to the independent Python checker. Lean uses the corresponding coefficient tables as literal Lean data; it does not parse that JSON file during proof checking. The theorem `residual_agrees_JSON` equates the residual computed from those factors with the literal polynomial `JSONResidual` exported from the supplied table. The Python checker separately verifies the external JSON residual and its coefficient norm. The compressed file `AmbientKernelData.lean.gz` contains Lean source for the Hessian certificate, not JSON. The build expands that source before importing it; all its assertions are then kernel-checked. No proof generator is run as a trusted build step.

A.6 Conditioning of the anchor alignment

A quantitative Gram-stability argument turns (2) into an orthogonal alignment. It uses the candidate anchor points p_2, p_3, p_6: the absolute values of the inner products of pairs $(2,3)$, $(2,6)$ and $(3,6)$ are at most $19/100$, $41/100$ and $41/100$, respectively. The corresponding pairs of relabeled competitor anchor points consequently satisfy the absolute inner-product bounds $1/5, 5/12, 5/12$. In Gram–Schmidt, each residual is the component of an anchor point orthogonal to the span of the preceding anchor points. The lengths of the two residuals are at least $2/3$ and $1/2$. Thus the divisions used to recover coordinates from inner products have denominators bounded away from zero.

In the resulting orthonormal frames, the coordinate differences for each point are at most δ, 22δ and 646δ, respectively. These scalar estimates follow by applying the product, square-root and quotient difference bounds to the Gram–Schmidt formulas; they are proved in `Local/GramStability.lean`. Relaxing each to 650δ and summing the 30 coordinate squares gives $\rho^2 \leq$

$30(650\delta)^2 \le (10000\delta)^2$ for the constructed alignment. `Local/GramAlignment.lean` transfers this bound to the minimizing alignment.

A.7 Local inequalities in Lean

This subsection identifies the formal inequalities used for full stationarity and the finite-radius energy gap. The interval Hessian calculation encloses the arithmetic and square-root operations over the full height interval. Its change-of-basis matrix is upper triangular with all diagonal entries equal to one, hence invertible. Interval bounds on the transformed matrix certify strict diagonal dominance. In `Local/Certificates/AmbientKernelBounds.lean`, the theorem `flattened_posDef` proves positive definiteness.

The theorem that proves (4) for the explicitly defined Hessian is listed below its source file. The last line is the Lean namespace prefix for both this theorem and `flattened_posDef`:

```
Local/Certificates/AmbientHessian.lean
candidate_augmented_hessian_posSemidef
ThomsonN10.Local.Certificates.AmbientKernel
```

The formal coordinate calculation rewrites the height equation as a relation among the seven inverse-cube distance weights, one for each candidate inner-product class, and checks the gradient at every point. The stationarity calculation in Section 4 has the following formal correspondence. In `Local/LocalStationarity.lean`, `candidate_force_relation` converts that derivative equation to a linear relation among the seven inverse-cube distance weights. The identity `orbit_force_balance` checks all force components and uses this relation to make each point's force radial; `candidate_force_eq_radialCoefficient` identifies the multiplier with the exact spherical radial coefficient. This proves stationarity for arbitrary tangent perturbations at all ten points.

The pair remainder inequality (5) is `pair_potential_algebraic_lower`. The local estimate uses the exact unit-sphere identity $\langle p_i, y_i - p_i\rangle = -\|y_i - p_i\|^2/2$. Its final conclusion is $\mathcal{E}(y) - \mathcal{E}(p) \ge \rho^2/200$ for aligned displacement norm $\rho \le 1/1000$.

A.8 The separate energy calculation

This subsection explains how to check the separate energy enclosure in Section 2. This calculation is checked only by Python. It is computed by rational interval evaluation of the candidate's 45 pair energies over $[a, b]$. Square roots are enclosed using integer square roots on a dyadic grid of spacing 2^{-420}, rounding outward. Reciprocal and summation operations preserve the rational bounds. The script `checks/energy_enclosure.py` performs this computation and prints the two endpoints in the energy remark in Section 2. Lean proves that the height interval $[a, b]$ contains its unique stationary root; Python uses the interval as input and does not prove root existence or uniqueness.

A.9 Code and cutoff provenance

This subsection records the source of the reused formal code and the relation between the two ten-point energy cutoffs.

The MIT-licensed Lean code of the `thomson-n9-lean` project [14] supplies the immediate precedent. Its AI-produced October 6, 2026 revision `168ff0c` proves that the D_{3h} tricapped trigonal prism is the unique Coulomb minimizer for nine distinct unit-sphere points, up to orthogonal maps and permutations, with its parameter defined by a unique root in a specified interval. Our Lean files adapt its theorem organization, energy bound with a small certified error, classification of inner-product intervals, orthogonal alignment and local energy argument. The adaptations supply

the ten-point geometry, candidate inner products, certificate data and their connection to the final theorem. A file-level attribution and account of changes is provided in `THIRD_PARTY.md`. The prose is new; the reused material is formal code.

Our preliminary energy cutoff 32717/1000 is exactly the 32.717 used in the October 11, 2026 chord-bound result of `saberwatchmanga` [8]: it is the known candidate energy rounded upward to three decimal places. Under this cap, our two-point certificate proves that every distinct pair has inner product below 13/20. That assertion is not deduced from their existence of a chord longer than 1.935. We cannot establish whether 32.717 was chosen independently or taken from that work; no step of this proof uses the earlier result.

References

[1] J. J. Thomson, On the structure of the atom: an investigation of the stability and periods of oscillation of a number of corpuscles arranged at equal intervals around the circumference of a circle; with application of the results to the theory of atomic structure, *Philosophical Magazine* 7 (1904), 237–265. doi:10.1080/14786440409463107.

[2] L. Föppl, Stabile Anordnungen von Elektronen im Atom, *Journal für die reine und angewandte Mathematik* 141 (1912), 251–302. doi:10.1515/crll.1912.141.251. See also the author's ICM 1912 report, vol. II, p. 190, https://www.mathunion.org/fileadmin/ICM/Proceedings/ICM1912.2/ICM1912.2.ocr.pdf.

[3] T. Erber and G. M. Hockney, Equilibrium configurations of N equal charges on a sphere, *Journal of Physics A: Mathematical and General* 24 (1991), L1369–L1377, Table 1. doi:10.1088/0305-4470/24/23/008. https://lss.fnal.gov/archive/1991/pub/Pub-91-222-T.pdf.

[4] D. J. Wales and S. Ulker, Cambridge Cluster Database, Thomson problem, ten-point entry. https://www-wales.ch.cam.ac.uk/~wales/CCD/Thomson/table.html.

[5] R. E. Schwartz, The Five-Electron Case of Thomson's Problem, *Experimental Mathematics* 22 (2013), 157–186. doi:10.1080/10586458.2013.766570. Preprint: The 5 Electron Case of Thomson's Problem, January 21, 2010, arXiv:1001.3702.

[6] N. N. Andreev, An extremal property of the icosahedron, *East Journal on Approximations* 2 (1996), no. 4, 459–462. Author's publication list. Review: Zbl 0877.51021.

[7] H. Cohn and A. Kumar, Universally optimal distribution of points on spheres, *Journal of the American Mathematical Society* 20 (2007), 99–148. arXiv:math/0607446.

[8] `saberwatchmanga` (formerly `terry2418`), Lean-verified local minimality for ten charges on a sphere (October 7, 2026), and A Lean-checked distance restriction for ten charges on a sphere (October 11, 2026). https://github.com/saberwatchmanga/-lean-finite-certificates.

[9] H. Tran, A Lean Proof of the Thomson Problem for Seven Electrons, Vals AI blog, September 28, 2026. https://www.vals.ai/blogs/thomson-n7-lean-proof.

[10] L. Kryvonos, L. Liehr and M. A. Taylor, Energy minimization for eight points on the sphere, September 18, 2026. arXiv:2609.22077.

[11] J. Tooby-Smith and A. Zughaid, `Thomson-N-8-Warrant`, Lean-checked warrant for the eight-point Coulomb Thomson problem, September 2026. https://github.com/jstoobysmith/Thomson-N-8-Warrant.

[12] A. Zarco (`alejandrozarco`), *Five points on the sphere: exact three-point certificates for the Riesz energies $s = 1, \dots, 6$, and Lean formalisations for $s = 1$ (Coulomb) and $s = 2$*, October 2026. doi:10.5281/zenodo.23125753. https://github.com/alejandrozarco/five-points-riesz.

[13] A. Zarco (`alejandrozarco`), *The pentagonal bipyramid as the unique logarithmic-energy minimiser for seven points on the sphere: an unreviewed Lean 4 warrant*, September–October 2026, with a Coulomb second check added October 8, 2026. doi:10.5281/zenodo.23086382. `https://github.com/alejandrozarco/thomson-n7-log`.

[14] A. Karthigeyan (director and publisher), `thomson-n9-lean`, MIT-licensed Lean code, AI-generated by Anthropic's Opus 5.5 and OpenAI's GPT-6 Astra, unreviewed, revision `168ff0c`, October 6, 2026. `https://github.com/aravindkarthigeyan/thomson-n9-lean`.

[15] P. Delsarte, J. M. Goethals and J. J. Seidel, Spherical codes and designs, *Geometriae Dedicata* 6 (1977), 363–388. doi:10.1007/BF03187604.

[16] V. A. Yudin, Minimum potential energy of a point system of charges, *Diskretnaya Matematika* 4 (1992), no. 2, 115–121; English translation in *Discrete Mathematics and Applications* 3 (1993), no. 1, 75–81. doi:10.1515/dma.1993.3.1.75.

[17] C. Bachoc and F. Vallentin, New upper bounds for kissing numbers from semidefinite programming, *Journal of the American Mathematical Society* 21 (2008), 909–924. doi:10.1090/S0894-0347-07-00589-9.

[18] H. Cohn and J. Woo, Three-point bounds for energy minimization, *Journal of the American Mathematical Society* 25 (2012), 929–958. arXiv:1103.0485.