

Brindlewythe Biosciences Ltd

QP07 metabolic-signal kinase potency and selectivity

Final signed study report — revision B

Study BW01_0059

Accession B-ACC-2001-01010

Client: Quorenthel Pharma GmbH

Date: 18 May 2001

Author: Dr B. Rowcroft, Screening Operations Director

1 Summary and control acceptance

The Primary potency assay and Selectivity sweep were accepted for the QP07 aminopyrazine screening set. The best primary IC50 was 4.4 nM. The primary ranking and the fixed-concentration off-target profile answer separate screening questions.

Control acceptance preceded interpretation. Ines Alvarado checked the acceptance record and its link to the sample cards before the screening conclusions were released. Censored estimates and individual qualifications remain part of the accepted endpoint record; study acceptance does not remove them.

Table 1. Study summary and benchmarks

Metric	Value
Project	BW01_0059
Client	Quorenthel Pharma GmbH
Compounds tested	24
Series / chemotype	Quorenthel aminopyrazine series 07 (Aminopyrazine)
Assays	Primary kinase IC50, Kinase selectivity panel
Turnaround (calendar days)	42
Price	GBP 10,700

2 Study objective and chronology

Determine potency in the metabolic-signal kinase assay and examine the off-target fingerprint of representatives from the QP07 screening set.

The study covered 24 compounds for Quorentel Pharma GmbH under BW01_0059, accession B-ACC-2001-01010. Work began on 06 Apr 2001. The interim position was recorded on 27 Apr 2001 and the completed assessment was issued on 18 May 2001. Revision B supersedes the interim position.

The screen gate concerned the next chemistry decision. It was not a clinical assessment or an assertion that the intended target mechanism had been demonstrated. Dr B. Rowcroft was responsible for the scientific interpretation and final release.

3 Test articles and accountability

Full QP07 identifiers linked the sample cards to the primary dilution map. The Selectivity sweep used a representative subset, identified by the panel column headings. Only those representatives have both primary and panel evidence; membership of the primary screening set does not imply that a compound was tested in the sweep.

Test lot identity was preserved from receipt through preparation and measurement. The reportable compound identifier belongs to the material submitted for this study. A replacement lot, new solid form or formulation is not covered merely because it has the same chemistry designation.

The sample card provides the connection between retained material and the assay record. Follow-up work should carry that connection forward while recording any change of lot. Result transfer must use full identifiers rather than rack position or abbreviated suffix.

4 Experimental procedure

The Primary potency assay used recombinant enzyme with ATP at its assay K_m . Test articles were prepared as a ten-point, three-fold dilution from 10 μM with duplicate wells. Vehicle and reference inhibitor wells accompanied the dilution series. Residual activity was expressed relative to vehicle after subtraction of assay background.

The Selectivity sweep tested representatives at 1 μM against the kinases listed in its result table. Each kinase had its own vehicle and background measurements. Raw signals from different enzymes were not compared directly. The screen returned percent inhibition at a stated concentration rather than off-target concentration-response estimates.

Preparation and read order were recorded on the run sheets. Primary and panel measurements were reduced separately. Reference response and background separation established plate acceptance before compound ranking was considered. A satisfactory control does not make an unusual compound curve mechanistically interpretable.

5 Plate acceptance and calculation

Plate acceptance required separation of background and vehicle activity and reference-inhibitor response within the established assay range. Ines Alvarado checked the acceptance record before release. Numerical per-plate controls remain in the run sheets; no control values have been inferred from the compound result table.

Primary concentration responses were fitted with a sigmoidal model and released as IC₅₀ in nM with Hill slope. Panel inhibition was normalised to the corresponding kinase vehicle control. Neither a low fixed-concentration response nor a shallow fitted slope was converted into a claim about binding mode.

Ines Alvarado checked units, column mapping and retention of qualifiers during report preparation. Bethany Rowcroft then considered the accepted endpoints against the study objective. A reference control demonstrates assay responsiveness under the test conditions; it does not establish specificity or suitability of a test compound outside those conditions.

6 Results: numerical record

Table 2 gives the full primary screening set. Table 3 is the Selectivity sweep at 1 μ M; its column headings define the compounds actually counterscreened. The best primary IC₅₀, 4.4 nM, is an assay endpoint rather than a direct affinity measurement.

Table 2. Primary potency assay: IC₅₀ (nM) and Hill slope

Compound	IC ₅₀ (nM)	Hill slope
QP07-08506	20	0.91
QP07-08518	17	0.94
QP07-08522	5.0	1.20
QP07-08530	51	0.99
QP07-08536	1260	1.04
QP07-08542	19	0.94
QP07-08558	51	0.88
QP07-08566	162	1.20
QP07-08578	18	1.01
QP07-08588	3902	1.05
QP07-08590	112	1.18
QP07-08594	85	1.04
QP07-08608	15	1.13
QP07-08620	271	0.91
QP07-08630	4.4	1.18
QP07-08632	23	1.14
QP07-08636	50	1.16
QP07-08642	12	0.85
QP07-08660	34	0.87
QP07-08662	111	1.24
QP07-08668	13	0.91
QP07-08672	19	1.20
QP07-08686	24	0.96
QP07-08690	33	1.21

ATP at Km; 10-point, 3-fold; staurosporine control

Table 3. Selectivity sweep: inhibition (%) at 1 μ M

Kinase	QP07-08506	QP07-08518	QP07-08522	QP07-08530	QP07-08536	QP07-08542
ABL1	21	32	0	22	0	0
AURKA	36	12	28	0	22	12
CDK2	21	26	72	0	47	34
CHEK1	40	0	48	33	15	14
EGFR	10	0	33	9	24	15
FLT3	33	18	30	0	2	4
GSK3B	54	70	62	23	35	0
JAK2	8	5	21	34	35	22
KDR	45	49	36	48	0	0
LCK	21	4	1	12	47	20
MAPK14	41	42	18	20	10	47
PIK3CA	55	0	36	49	7	3
SRC	3	64	14	23	24	18
SYK	43	51	39	16	42	4
BTk	35	0	35	41	16	43
ROCK1	0	0	37	0	60	12

The released tables are the primary numerical record. A blank cell is not a zero. A greater-than value is not an exact estimate. Concentration, matrix and endpoint units must remain attached to each result when it is copied from this report.

7 Comparison within the screening set

The panel did not overturn use of the primary ranking for the present screen gate. The compact heteroaryl subset remains the preferred branch, but the panel columns delimit that evidence. The same off-target profile cannot be assigned to unscreened neighbours solely because they share the screening-set designation.

Comparison within the present study is preferred to an unqualified comparison with historical screening sets. Test lot, preparation and assay conditions can change the apparent separation between candidates. A useful chemistry association remains a screening observation unless a further experiment distinguishes its possible causes.

The assay profile should be read in full before selecting a favourable isolated result. An optimisation preference does not change the endpoint, and a less favoured branch does not become an assay failure because the client chooses another direction.

8 Qualifications and interpretation limits

A fixed-concentration off-target result cannot establish an IC₅₀ or a selectivity ratio. The panel is restricted to its listed kinases and representatives; it cannot exclude activity at untested enzymes or other protein families. The primary assay does not establish cellular activity or an in vivo mechanism.

Hill slope is retained as a curve descriptor, not evidence of cooperativity or binding stoichiometry. Closely spaced potency values should be considered with the underlying curves. Negative or low panel inhibition must not be reinterpreted as stimulation or proof of absence of binding.

No exposure margin has been calculated. Such a calculation requires an agreed exposure measure and a justified connection between that measure and this assay endpoint. The report establishes neither a dose nor an in vivo response. These limits define the intended use of the screen.

9 Screen gate and client decision

The primary potency ranking survived the selectivity counterscreen, and Quorenthel Pharma GmbH progressed the compact heteroaryl subset.

Progression of the compact heteroaryl subset is a screening preference. Less favoured compounds remain in the archive as comparators; their accepted measurements are not assay failures.

Quorenthel Pharma GmbH owns the chemistry decision. Brindlewythe Biosciences Ltd supplies the accepted experimental profile and its interpretation limits. The decision is recorded separately so that a progression preference is not mistaken for an additional experimental observation.

10 Follow-up work specification

Concentration responses on the principal off-target signals of a retained candidate would resolve whether its fixed-concentration fingerprint gives useful potency separation. A repeat primary assay using the same test lot would permit comparison under a common material identity. Cellular target activity and exposure require separate studies.

Any further study should state its question before allocating samples. Retain the existing compound identifier and test lot, and use a new accession for newly submitted material. A repeat intended to resolve a qualification should retain that purpose in its run sheet rather than being merged silently with the present result.

Proposed follow-up does not imply that the additional experiment has already been completed. Only the assays and endpoints in this report are released under the current study.

11 Study records and data transfer

The primary and Selectivity sweep run sheets form distinct record groups. Preserve the complete representative identifiers in the panel headings when transferring the tables to the client chemistry register. Do not expand the panel scope by joining it to all rows in the primary table.

The study file contains the sample-card link, preparation record, run sheets and released endpoint tables. Bethany Rowcroft prepared the interpretation and Ines Alvarado checked it against the accepted assay record. The final report should be retained with those records rather than used as a replacement for them.

When quoting the assessment, retain BW01_0059, revision B, the report date and endpoint units. A copied ranking without its qualifications loses information needed at the screen gate. This report is confidential to Quorenthel Pharma GmbH and Brindlewythe Biosciences Ltd.

12 Conclusion and release

The primary potency ranking survived the selectivity counterscreen, and Quorentel Pharma GmbH progressed the compact heteroaryl subset.

The accepted experimental record supports that screening decision within the conditions specified. It does not extend the scope to untested compounds, matrices or biological endpoints. The numerical tables and qualifications remain the basis for any subsequent use of the results.

Revision B is the final report issued on 18 May 2001. The signatures record scientific release and the check of sample mapping, units and result qualifications.

Name	Role	Signature	Date
Bethany Rowcroft	Screening Operations Director — scientific release	<u>B Rowcroft</u>	18 May 2001
Ines Alvarado	Assay Quality Coordinator — report check	<u>I Alvarado</u>	18 May 2001