

Osterquill Toxicology Inc

ST03 growth-signal inhibitors: cytotoxicity and micronucleus follow-up

Final investigation report — exploratory package

Osterquill Toxicology Inc

Report no. G-R-2015-0064-A

Investigation OT-15-S0064

LIMS OQ/2015/001043

Client: Selverant Therapeutics Inc

Date: 07/14/2015

Prepared and authorized by Dr N. Takamori, Head of Toxicology

1 Executive conclusion

OT-15-S0064 investigated the ST03 test article cohort for Selverant Therapeutics Inc. The HepG2 viability screen and Mammalian-cell micronucleus assessment were interpreted together.

The micronucleus signal tracked cytotoxic exposure. Selverant Therapeutics Inc requested a lower-exposure repeat before assigning a genotoxic interpretation.

A lower-exposure repeat is recommended. The present signal is attributed to cytotoxic exposure within this exploratory package; that attribution does not establish absence of genotoxic potential at other exposures.

Control validity was accepted for the reported endpoints. Acceptance establishes that the test systems responded appropriately under the investigation conditions; it does not establish that any test article is safe. This signed report is the final scientific account of an exploratory package and is not a statement of GLP compliance.

Table 1. Investigation identification and package summary

Metric	Value
Project	OT-15-S0064
Client	Selverant Therapeutics Inc
Compounds tested	40
Series / chemotype	Selverant aminoquinazoline series 03 (Aminoquinazoline)
Assays	In vitro cytotoxicity (HepG2), In vitro micronucleus
Turnaround (calendar days)	40
Price	USD 42,300

2 Purpose and scope

The purpose was to inform the next chemistry decision for Selverant Therapeutics Inc before further development of the ST03 cohort. The investigation began on 06/04/2015; the interim interpretation was issued on 06/24/2015; this report closes the package on 07/14/2015. The reported cohort contains 40 test articles. The number refers to the supplied screening cohort and does not imply that every assay table contains a separate entry for every article.

The package asks whether changes in cell viability affect the interpretation of the mammalian-cell endpoint, and whether the results support a practical selection or replacement decision. It is deliberately narrower than a comprehensive genetic toxicology evaluation. No clinical exposure margin, human risk estimate or definitive mechanism is derived here.

The observations belong to the tested material and the exposure conditions recorded on the observation sheets. A later dose lot, changed formulation or replacement analogue requires its own traceability and an assessment of whether the present observations remain applicable. Names used in this report are laboratory identifiers, not a statement of chemical equivalence.

3 Responsibility and report status

Dr N. Takamori held responsibility for the integrated scientific interpretation and final authorization. Dr K. Morita evaluated the genetic toxicology endpoints with attention to exposure, cell growth and the corresponding controls. Lucia Moretti assembled the archive cross-reference and checked the report against the retained investigation documentation.

The scientific sign-off is distinct from the archive check. The involvement of the GLP Quality Assurance and Archive Lead does not convert exploratory work into a GLP investigation. A formal GLP compliance statement is not made, and this report should not be cited as a replacement for a regulatory genetic toxicology battery.

The final issue supersedes the interim interpretation for the conclusions stated here. Source observation sheets retain their original entries. The report does not overwrite those entries or create a new numerical dataset. Where a table and a narrative statement address different levels of the package, the table describes the measured endpoint and the narrative identifies the bounded project decision.

4 Test article accountability

Test article accountability is maintained under LIMS OQ/2015/001043. The ST03 identifiers in the HepG2 table are the primary cross-reference for the screened material. Receipt, preparation, dilution and disposition documentation remain linked to that investigation identifier in the Osterquill archive.

Material identity was carried through preparation and exposure rather than inferred from the order of wells or tubes. Changes in container, preparation or assay position require an explicit source entry so that an apparent difference in biological response cannot be mistaken for a difference in analogue identity.

This report does not assign an unlisted structure, purity value or stability period to any article. The interpretation assumes the tested preparations are those identified on the observation sheets. Should later analytical information change that assumption, the scientific conclusion requires reconsideration through a study amendment rather than an informal alteration of this signed report.

5 Exposure design

Concentration selection was guided by cell tolerance, practical preparation limits and the need to examine the relevant exposure window. A nominal concentration alone is not an exposure margin. Differences in cell system, exposure duration and metabolic activation prevent direct numerical conversion of a HepG2 IC50 into a TK6 challenge concentration.

The HepG2 viability screen retained the short and extended exposure readouts separately. The Mammalian-cell micronucleus assessment retained treatment condition, concentration, micronucleus frequency and relative population doubling in the same table. This layout preserves the relation between the chromosome endpoint and cell growth at the tested exposure.

Concentration entries in the micronucleus table are expressed in $\mu\text{g/mL}$, while the HepG2 safety potency is expressed in μM . These units are intentional and are not treated as interchangeable. No molecular-weight conversion is introduced without a traceable article-specific basis. Comparisons in this report are qualitative where such conversion is unavailable.

6 HepG2 viability screen: method

Cell viability was assessed through ATP content after the specified exposure intervals. The endpoint is a practical measure of viable cell mass under these conditions. A reduction in ATP signal may reflect loss of cells, impaired metabolic state or altered growth; it is not a direct measurement of chromosome damage.

Vehicle exposure supplied the comparison for the test article response. Preparation and exposure observations were retained with the assay output so that a response potentially affected by precipitation, handling or an abnormal control could be separated from a reproducible concentration-related effect.

Concentration-response interpretation retained censored values as reported. An entry above the tested range is not an exact IC50 at the boundary, and an unresolvable curve is not silently replaced with an estimated value. The table therefore remains the primary source for comparison of the test article cohort.

7 HepG2 viability screen: results

The complete HepG2 result table for the ST03 cohort follows. Both exposure intervals are displayed for each listed article. Readers should compare the paired entries before assigning a relative tolerance category; the later exposure can reveal a limitation that is less apparent after the shorter interval.

The values are screening readouts under the investigation conditions. They do not define a maximum tolerated systemic dose and should not be converted to mg/kg/day. The table provides the cytotoxicity context for the genetic toxicology discussion, rather than an independent clinical safety margin.

Table 2. HepG2 viability screen — ATP content IC50, μ M

Compound	24 h IC50 (μ M)	48 h IC50 (μ M)
ST03-01204	17	12
ST03-01210	47	33
ST03-01212	34	24
ST03-01218	6.5	4.6
ST03-01222	74	52
ST03-01224	13	9.0
ST03-01230	46	32
ST03-01234	18	12
ST03-01246	31	22
ST03-01248	>100	87
ST03-01252	46	32
ST03-01254	>100	89
ST03-01264	9.4	6.5
ST03-01272	8.4	5.9
ST03-01278	68	48
ST03-01284	20	14
ST03-01290	32	22
ST03-01294	7.8	5.4
ST03-01296	16	11
ST03-01302	32	23
ST03-01306	57	40
ST03-01312	35	24
ST03-01314	22	15
ST03-01320	16	11
ST03-01324	30	21
ST03-01330	16	11
ST03-01332	27	19
ST03-01338	57	40
ST03-01344	>100	87
ST03-01348	38	27
ST03-01350	3.7	2.6
ST03-01354	11	7.5
ST03-01356	53	37
ST03-01360	35	24
ST03-01362	27	19
ST03-01366	10	7.1
ST03-01378	5.6	3.9
ST03-01380	26	18
ST03-01384	49	34
ST03-01390	100	70

Chlorpromazine control

8 Viability interpretation

The comparison across exposure intervals was used to distinguish relative exposure tolerance from a simple rank order of nominal potency. A stronger effect after extended exposure indicates that the tolerated concentration window may depend on duration. It does not, by itself, identify the responsible molecular pathway.

The interpretation was kept at the level supported by the screening data. A test article with a higher viability IC50 can be less cytotoxic in this system without being preferable on every development criterion. Conversely, a lower IC50 can restrict the useful concentration range for a mammalian-cell follow-up without proving a primary chromosome-damaging mechanism.

Exposure limitation, rather than a new structural alert, is the immediate basis for the follow-up. The repeat must retain sufficient exposure to be informative while avoiding the range in which marked growth suppression confounds micronucleus interpretation.

Mammalian-cell micronucleus assessment: method

TK6 cells were examined under the treatment and activation conditions listed in the result table. Micronucleus frequency and relative population doubling were retained together. Cell growth is essential to interpretation because a chromosome endpoint observed at a severely cytotoxic exposure may not support the same conclusion as one observed in a well-tolerated window.

The assessment included concurrent vehicle and positive-control comparisons. Scoring and exposure information were kept on the genetic toxicology observation sheets. The final evaluation considered the exposure condition as well as the apparent frequency; it did not treat every numerical departure from vehicle as an independent mechanistic finding.

The endpoint identifies a mammalian-cell flag within the investigation conditions. It does not distinguish chromosome breakage from chromosome loss without additional evidence. No such mechanistic distinction is asserted by this report.

Mammalian-cell micronucleus assessment: results

The treatment-level result table is presented intact, including the relative population doubling field. This field must accompany the micronucleus values when the results are circulated. Removing it would conceal the exposure-tolerance context on which the present interpretation depends.

The micronucleus signal tracked cytotoxic exposure. Selverant Therapeutics Inc requested a lower-exposure repeat before assigning a genotoxic interpretation.

The table describes the reported treatment conditions; it is not a compound-by-compound certificate for the entire HepG2 screening cohort. The detailed association between preparations and treatment entries is maintained in the genetic toxicology observation sheets.

Mammalian-cell micronucleus assessment — TK6 treatment results

Treatment	Conc (µg/mL)	% MN cells	RPD (%)
3 h +S9	vehicle	1.09	100
3 h +S9	50	1.13	95
3 h +S9	100	0.81	89
3 h +S9	200	0.61	67
3 h -S9	vehicle	0.94	100
3 h -S9	50	0.85	92
3 h -S9	100	0.39	94
3 h -S9	200	0.70	88
24 h -S9	vehicle	0.73	100
24 h -S9	50	1.27	90
24 h -S9	100	0.27	87
24 h -S9	200	0.44	81

Outcome: negative

Control validity statement

Control validity was accepted for the reported HepG2 and TK6 datasets.

The vehicle controls supplied an interpretable baseline, and the positive controls demonstrated the expected responsiveness of the relevant test system. Cell growth and exposure observations were considered alongside these controls. The final interpretation uses the accepted datasets and does not rely on a failed-control run.

Control acceptance and test article interpretation are separate decisions. An accepted control establishes that the endpoint can be evaluated, while the test article conclusion still depends on exposure tolerance and the pattern of the observations. The report makes no inference that control acceptance alone establishes absence of hazard.

Numerical control information is retained within the relevant source tables and observation sheets. A new control range has not been invented for this report. Historical comparisons support the laboratory evaluation but are not presented as additional measured results in this document.

Cytotoxicity and chromosome-endpoint association

The integrated review considered whether the mammalian-cell flag occurred in an exposure range where cell proliferation was substantially affected. The rationale is biological rather than arithmetical: the chromosome endpoint is assessed in cells that must tolerate the exposure and proceed through the relevant growth process.

HepG2 IC50 and TK6 relative population doubling cannot be substituted for one another. They were used as complementary evidence of exposure tolerance. A quantitative margin between them would require information not established by this package, including article-specific conversion and the comparability of intracellular exposure.

The micronucleus signal tracked cytotoxic exposure. Selverant Therapeutics Inc requested a lower-exposure repeat before assigning a genotoxic interpretation.

The association informs the follow-up choice without supplying a definitive molecular mechanism. A repeat at lower exposure can test the persistence of the flag in a more interpretable window, whereas analogue replacement can manage the concern without claiming that the original mechanism has been resolved.

Integrated scientific interpretation

A lower-exposure repeat is recommended. The present signal is attributed to cytotoxic exposure within this exploratory package; that attribution does not establish absence of genotoxic potential at other exposures.

The final conclusion was reached by preserving the endpoint-specific findings, not by selecting the most favorable assay. The viability results define a tolerance context; the mammalian-cell results define the flag; the client decision states how the package was used for the next development step.

The investigation supports prioritization under the tested conditions. It does not define an in vivo genotoxicity outcome, a carcinogenicity classification or an acceptable human exposure. No inference of animal tolerability is drawn from these cell-based results.

Where a decision is more conservative than the narrow assay conclusion, the distinction is intentional. A sponsor may replace or discontinue an analogue to reduce development uncertainty while the scientific report continues to preserve the exact endpoint outcome.

Development decision and follow-up

Exposure limitation, rather than a new structural alert, is the immediate basis for the follow-up. The repeat must retain sufficient exposure to be informative while avoiding the range in which marked growth suppression confounds micronucleus interpretation.

Selverant Therapeutics Inc owns the subsequent chemistry and nomination decision. Osterquill recommends that the next package preserve original article identities, dose lot traceability and the exposure rationale. A new result should be compared with the present package only after these elements have been reconciled.

The next protocol should define the intended question before testing begins: persistence of the mammalian flag at tolerated exposure, behavior of replacement material, or relative tolerance of retained analogues. These questions require different selection and exposure choices and should not be combined into an undefined confirmatory exercise.

The final signed report provides the scientific basis for that planning. No unperformed repeat is described as complete, and no numerical prediction is assigned to the next experiment.

Limitations and applicability

The package is exploratory. It is limited to the tested articles, preparations, cell systems and exposure conditions. It does not include a full assessment of absorption, distribution, metabolism or systemic exposure. The concentration-response results cannot be translated directly into a human dose.

The HepG2 screen and TK6 assessment use distinct biological systems. Different sensitivity between them is expected to require interpretation rather than forced numerical alignment. The present report deliberately avoids a calculated cross-system safety margin.

The available tables do not establish a definitive mechanism for the mammalian-cell finding. The chemistry decision is supported by the reported concern, but the report does not identify a reactive metabolite, DNA adduct or chromosome-loss mechanism without direct evidence.

The conclusions are confined to this exploratory package. Further regulatory work requires a prospectively defined protocol and its own compliance documentation; this final report is not a GLP certificate.

Animal welfare and use statement

This package used in vitro test systems. No live-animal administration, recovery cohort or animal welfare endpoint formed part of the work reported here. Consequently, this report contains no animal-card observations or inferred animal dose recommendation.

Metabolic activation material used in an in vitro assay does not make the package an in vivo investigation. Animal welfare language in Osterquill report practice is retained to make the boundary explicit, rather than to imply animal procedures that were not performed.

Any future in vivo work must have a separate welfare assessment, exposure rationale and prospective authorization. It cannot be justified solely by converting the cell-based concentrations into a systemic dose.

Data handling and numerical conventions

The numerical tables are the primary reported datasets. Units remain attached to their endpoints. Censored viability entries retain the greater-than notation; activation conditions remain attached to bacterial or mammalian treatments; dispersion is displayed with the bacterial means where available.

No missing result is replaced by a zero. No table entry is treated as an exact concentration-response parameter if it is reported as a bound. Narrative statements describe the qualitative interpretation and do not introduce untraceable numerical results.

The treatment-level micronucleus table is not expanded into invented article-level rows. Similarly, the screening cohort count is not used as an implied number of genetic toxicology treatments. These distinctions are necessary to retain the meaning of the package when the report is used outside the laboratory.

Report-to-source cross-reference

The signed report is filed under OT-15-S0064, with preparations and observation sheets linked through OQ/2015/001043. The HepG2 dataset, genetic toxicology datasets and final interpretation remain identifiable as separate archive components.

Dr K. Morita is the scientific contact for the genetic toxicology interpretation. Dr N. Takamori is responsible for any change to the final integrated conclusion. Lucia Moretti maintains the archive linkage and the status of any subsequent study amendment.

A request to revise an endpoint conclusion must identify the new evidence and the affected source entries. Editorial clarification may improve readability, but it must not silently alter treatment identity, units, control status or the endpoint outcome.

Final authorization

The micronucleus signal tracked cytotoxic exposure. Selverant Therapeutics Inc requested a lower-exposure repeat before assigning a genotoxic interpretation.

A lower-exposure repeat is recommended. The present signal is attributed to cytotoxic exposure within this exploratory package; that attribution does not establish absence of genotoxic potential at other exposures.

Authorized on 07/14/2015 by Dr N. Takamori. The report is final for the investigation identified on the title page. Its issue closes the present exploratory package; it does not represent completion of the proposed follow-up.

The archive completeness signature applies to the filed report and its source cross-reference. It is expressly separate from scientific authorization and from any formal statement of GLP compliance.

Name	Role	Signature	Date
Naomi Takamori	Head of Toxicology — scientific authorization	<u>N Takamori</u>	07/14/2015
Lucia Moretti	GLP Quality Assurance and Archive Lead — archive completeness, not GLP certification	<u>L Moretti</u>	07/14/2015

Appendix A — Exposure-window note

The exposure window must permit an interpretable cell-growth response. Selection of a lower concentration should follow the observed tolerance pattern, not an arbitrary reduction intended only to remove a flag.

The repeat should retain the original treatment condition as a comparator where scientifically appropriate. If duration or activation is changed, that change answers a different question and must be identified in the resulting report.

Exposure limitation, rather than a new structural alert, is the immediate basis for the follow-up. The repeat must retain sufficient exposure to be informative while avoiding the range in which marked growth suppression confounds micronucleus interpretation.

Appendix B — Censored viability entries

A greater-than entry means the viability IC₅₀ was not resolved within the tested range. It is a bound, not an exact value at the highest concentration.

For prioritization, bounded entries may support relative tolerance but cannot support a precise fold comparison. Such comparisons should await a suitable extended concentration-response experiment, if one is scientifically justified.

Exposure limitation, rather than a new structural alert, is the immediate basis for the follow-up. The repeat must retain sufficient exposure to be informative while avoiding the range in which marked growth suppression confounds micronucleus interpretation.

Appendix C — Micronucleus reporting note

Relative population doubling must travel with micronucleus frequency. Separating the fields would remove the context necessary to distinguish an exposure-limited flag from a finding in a tolerated window.

The present package supports the project decision stated in the executive conclusion. It does not establish a chromosome-breakage or chromosome-loss mechanism, and neither mechanism should be inserted into downstream summaries.

Exposure limitation, rather than a new structural alert, is the immediate basis for the follow-up. The repeat must retain sufficient exposure to be informative while avoiding the range in which marked growth suppression confounds micronucleus interpretation.

Data annex — HepG2 table use

Table 2 is the complete screening table. The annex confirms the reporting conventions used for its two exposure intervals. The μM values remain safety potency readouts in the HepG2 system; no systemic dose conversion has been applied.

The source observation sheets retain preparation and exposure details. Any later correction must be assessed for its effect on both the table and the integrated interpretation.