# Four-zero tail-palindromic polynomials over $\mathbb{F}_2$: reciprocal obstructions and quadratic-transform intersections

Baohua Li

29 October 2026

**Abstract**

We study monic polynomials of even degree over $\mathbb{F}_2$ whose nonleading coefficients form a palindrome with constant term one and exactly four zeros. We determine their reciprocal common divisor and count the position pairs passing this obstruction. We classify their intersections with full-degree quadratic rational transformations: the translated $Q$ branch comprises two infinite shape families and three degree-14 exceptions, while the translation-invariant branch comprises four reducible polynomials at degrees 14 and 30. More generally, a translation-invariant tail palindrome has $n+2$ a power of two and, for $n \geq 6$, is reducible if its number of missing tail coefficients is divisible by four. The family-specific content is the compatibility of classical invariant representations and irreducibility criteria with the prescribed support, yielding restrictions on quadratic degree-doubling constructions. Independent arithmetic checks recover 2663 primitive polynomials at all even degrees from 6 to 200 and verify the first solutions through degree 300. General primitive existence remains unproved.

## 1 The family and the question

Put $n = 2k \geq 6$ and $S_n(x) = 1 + x + \cdots + x^n$. We consider

$$f_{n,i,j}(x) = S_n(x) + x^i + x^j + x^{n-1-i} + x^{n-1-j}, \qquad 1 \leq i < j \leq k-1, \tag{1}$$

over $\mathbb{F}_2$. These are exactly the monic degree-$n$ polynomials with constant term one whose nonleading coefficient vector is a palindrome with four zero entries. The restrictions on $i, j$ exclude the constant position. There are $\binom{k-1}{2}$ candidates. Let $N_4(n)$ denote the number of primitive polynomials in this family. Here primitive means irreducible with a root of multiplicative order $2^n - 1$; standard background is in [2].

**Conjecture 1.1** ($C_4$)**.** For every even $n \geq 6$, one has $N_4(n) > 0$.

Primitive polynomials with prescribed coefficients form a classical subject; see Cohen [1]. In the present question the field is fixed and all but four tail coefficients are one, with the four exceptions coupled by reversal. Results prescribing a small number of coefficients do not directly settle this growing collection of constraints. A tail palindrome can also be encoded by the self-reciprocal polynomial $xf(x) + 1$ of odd degree $n + 1$. This change of polynomial preserves the coefficient description, but does not preserve irreducibility or primitivity.

The main results are the reciprocal obstruction and surviving position count in Section 2, and the complete quadratic support intersections in Section 3. Their invariant representations and irreducibility criteria are classical; the additional arguments concern compatibility with the prescribed support. Section 4 records independently verified finite data. Appendix A collects the auxiliary classical coefficient and field descriptions. No general irreducibility criterion or proof of Conjecture 1.1 is asserted.

## 2 The reciprocal obstruction

This section isolates a part of the reducibility problem that admits an exact description. Its tools are classical: polynomial reciprocity, the binomial gcd identity [2], and unit counts related to the Schemmel functions [5]. The contribution here is their explicit specialization to the four-zero family, including a complete reciprocal common divisor and an exact count of the surviving position pairs. These are auxiliary necessary conditions; no new general irreducibility principle is asserted.

**Lemma 2.1** (Reciprocity and multiplicities)**.** *Let $f = x^n + \sum_{r=0}^{n-1} c_r x^r$, where $n$ is even, $c_0 = 1$, and $c_r = c_{n-1-r}$. Write $f^*(x) = x^n f(1/x)$. Then*

$$f^* = xf + x^{n+1} + 1, \qquad \gcd(f, f^*) = \gcd(f, x^{n+1} + 1).$$

*This gcd is squarefree. In particular, every self-reciprocal irreducible factor of $f$ occurs with multiplicity one. If a non-self-reciprocal irreducible $q$ and $q^*$ both divide $f$, the smaller of their multiplicities is one.*

*Proof.* The coefficient symmetry gives the displayed identity, hence the gcd identity by subtraction. The derivative of $x^{n+1} + 1$ is $x^n$, which is coprime to it; thus its divisors are squarefree. A self-reciprocal factor has the same multiplicity in $f$ and $f^*$. For a reciprocal pair the two multiplicities are interchanged, so their minimum is the multiplicity in the gcd. This proves the claims. □

**Proposition 2.2** (Exact reciprocal common divisor)**.** *For $f = f_{n,i,j}$ from (1), put*

$$N = n + 1, \quad \Delta = j - i, \quad R = n - 1 - i - j, \quad a = \gcd(N, \Delta), \quad b = \gcd(N, R), \quad h = \gcd(a, b).$$

*Then, as an exact polynomial quotient in $\mathbb{F}_2[x]$,*

$$\gcd(f, f^*) = \frac{(x^a + 1)(x^b + 1)}{(x + 1)(x^h + 1)}, \qquad \deg \gcd(f, f^*) = a + b - h - 1. \tag{2}$$

*Consequently, irreducibility of $f$ requires $a = b = 1$.*

*Proof.* The nonleading terms of $f$ occur in pairs, so $f(1) = 1$. By Lemma 2.1, with $S_n = (x^N + 1)/(x + 1)$,

$$\gcd(f, f^*) = \gcd(f, S_n) = \gcd(S_n, P), \qquad P = x^i(1 + x^\Delta)(1 + x^R).$$

The oddness of $N$ makes $S_n$ squarefree. The classical binomial gcd formula gives

$$\gcd(S_n, 1 + x^\Delta) = \frac{x^a + 1}{x + 1}, \qquad \gcd(S_n, 1 + x^R) = \frac{x^b + 1}{x + 1}.$$

Since $S_n$ is squarefree, its gcd with their product is their least common multiple. Their intersection is $(x^h + 1)/(x + 1)$, proving (2). If this gcd were all of $f$, monicity and equal degrees would give $f = f^*$, hence $(x + 1)f = x^N + 1$ and $f = S_n$, contrary to its four zero coefficients. Thus a nonconstant gcd is a proper divisor. The right side is constant exactly when $a = b = 1$. □

**Corollary 2.3** (An explicit infinite reducible family)**.** *For every odd $m \geq 3$, the valid position pair $(n, i, j) = (3m - 1, m - 2, m)$ satisfies*

$$\begin{aligned} f_{3m-1,m-2,m}(x) = (1 + x + \cdots + x^{m-1}) \\ \cdot (x^{2m} + x^{m+1} + x^{m-1} + x^{m-2} + 1). \end{aligned} \tag{3}$$

*In particular it is reducible, and the displayed forced divisor has unbounded degree $m - 1$.*

*Proof.* The inequalities $1 \le m-2 < m \le (3m-3)/2$ make the positions valid. Here $N = 3m$, $\Delta = 2$, and $R = m$, so $a = 1$, $b = m$ in Proposition 2.2. For the explicit quotient, group the geometric sum as $S_{3m-1} = S_{m-1}(1 + x^m + x^{2m})$. The four deleted terms sum to

$$x^{m-2}(1+x^2)(1+x^m) = S_{m-1}x^{m-2}(x+1)^3.$$

Adding these expressions and cancelling the two $x^m$ terms inside the second factor proves (3). □

**Proposition 2.4** (Number of positions surviving the screen)**.** *For odd $N = n+1$, let $\varphi$ be Euler's totient and put*

$$\mathfrak{S}_2(N) = N \prod_{p \mid N} \left(1 - \frac{2}{p}\right).$$

*The number of four-zero position pairs with* $\gcd(f, f^*) = 1$ *is*

$$E_N = \binom{\varphi(N)/2}{2} - \frac{\mathfrak{S}_2(N) - 1}{2}. \tag{4}$$

*Thus $N_4(n) \le E_{n+1}$. This is only an upper bound; $E_{n+1} > 0$ does not imply $N_4(n) > 0$.*

*Proof.* Set $u = i+1$, $v = j+1$. The two coprimality conditions in Proposition 2.2 are precisely that $u - v$ and $u + v$ be units modulo $N$. Since 2 is invertible, the map $(u, v) \mapsto (u - v, u + v)$ is a bijection on $(\mathbb{Z}/N\mathbb{Z})^2$, giving $\varphi(N)^2$ ordered pairs before excluding zero coordinates. The cases $u = 0$ and $v = 0$ contribute $\varphi(N)$ each and are disjoint. The independent sign changes of $u, v$ and their interchange act freely on the remaining pairs: admissibility excludes $u = \pm v$, and a nonzero coordinate cannot equal its negative. Dividing by eight gives $\varphi(N)(\varphi(N) - 2)/8$ unordered pairs of distinct nonzero sign classes.

Choose representatives $1, \dots, k$ of those classes. The allowed positions use $2, \dots, k$, so the class represented by 1 must be removed. The Chinese remainder theorem counts $\mathfrak{S}_2(N)$ residues $v$ for which both $v - 1$ and $v + 1$ are units: modulo each prime $p \mid N$ two residues are forbidden. One residue is $v = 0$; the others form pairs $\{v, -v\}$. Hence the removed class has $(\mathfrak{S}_2(N) - 1)/2$ neighbors. Subtraction proves (4). □

*Remark* 2.5 (Scope and classical predecessors). These results describe the reciprocal common part, not the complete factorization. For example,

$$f_{8,1,2} = (x^3 + x^2 + 1)(x^5 + x^2 + 1)$$

has trivial reciprocal gcd but is reducible. Nor are candidates always squarefree:

$$f_{14,2,5} = (x^2 + x + 1)(x^6 + x^4 + x^3 + x + 1)^2.$$

The squarefreeness hypothesis in the classical Stickelberger–Swan factor-parity criterion [3] must therefore be checked. An odd number of factors still need not mean irreducibility: $f_{12,4,5}$ has three distinct factors of degrees $3, 4, 5$.

For fixed $i, j$, one also has

$$(x+1)f_{n,i,j} = x^{n-1-j}A_{i,j} + B_{i,j},$$

where $A_{i,j} = x^{j+2} + (x+1)(x^{j-i} + 1)$ and $B_{i,j} = 1 + (x+1)(x^i + x^j)$ are fixed polynomials. This places fixed-position factor-parity questions in the classical framework of Fredricksen, Hales and Sweet [4]. Neither eventual periodicity in that framework nor the elementary formulas above are claimed as a new general factorization method.

# 3 Quadratic-transform intersections

## 3.1 The translated $Q$ branch

For a degree-$m$ polynomial $h$, define

$$\mathcal{T}(h)(x) = (x+1)^m h\left(\frac{x^2}{x+1}\right), \qquad \mathcal{M}_n(f)(x) = (x+1)^n f\left(\frac{x}{x+1}\right). \tag{5}$$

The first polynomial is the classical $Q$-transform of $h$, followed by $x \mapsto x+1$. The equivalence between $\mathcal{M}_{2m}(f) = f$ and $f = \mathcal{T}(h)$ is a special case of Mattarei and Pizzato [8, Lemma 9]: take $a = b = 1, c = 0$, numerator $x^2$ and denominator $x+1$ in their notation. The nondegeneracy conditions hold, and the representation is unique; indeed the summands $x^{2j}(x+1)^{m-j}$ have distinct degrees $m+j$. More general invariant representations are given by Reis [9, Theorem 1.3]. These representations and the transform are classical. Our task here is to determine their intersection with the four-zero coefficient support, including reducible polynomials.

Write $\Delta a = a(x+1) + a(x)$, and use reversal with respect to a specified formal degree when the leading coefficient may vanish.

**Lemma 3.1** (A descent for tail palindromes)**.** *Let $f = x^{2k} + \sum_{j=0}^{2k-1} c_j x^j$, with $c_0 = 1$ and $c_j = c_{2k-1-j}$. Set $d = k-1$ and*

$$e = \sum_{a=0}^{k-1} c_{2a} x^a, \qquad o = x^d e(1/x).$$

*Then*

$$\mathcal{M}_{2k}(f) = f \quad \Longleftrightarrow \quad \Delta e = 0 \text{ and } e = \Delta(xo). \tag{6}$$

*If the tail of $f$ has $2s$ zeros, the formal coefficient vector of $e$ has $s$ zeros.*

*Proof.* Tail symmetry gives $f = x^{2k} + e^2 + xo^2$. Put $p = xf + x^{2k+1} = xe^2 + x^2o^2$, and homogenize $p$ to formal degree $2k+1$, obtaining $P(X,Y)$. The tail condition is exactly $P(X,Y) = P(Y,X)$. Moreover, $\mathcal{M}_{2k}(f) = f$ is equivalent to $P(X, X+Y) = P(X,Y)$. Conjugating this substitution by the interchange of $X, Y$ gives $(X,Y) \mapsto (X+Y, Y)$. Hence, for this symmetric $P$, the condition is equivalent to $\Delta p = 0$. Direct expansion gives

$$\Delta p = x(\Delta e)^2 + \big(e(x+1) + \Delta(xo)\big)^2.$$

The two summands have disjoint odd and even supports. Thus both vanish exactly under the conditions in (6). Each tail mirror pair contains one even and one odd position, which proves the assertion about zeros. □

**Lemma 3.2** (Two missing coefficients and translation)**.** *Let $d$ be positive and even, and $1 \le a < b \le d-1$. Then $e = S_d + x^a + x^b$ satisfies $\Delta e = 0$ exactly in the following cases:*

| | $d$ | $\{a, b\}$ |
|---|---|---|
| I | $2^r - 2,\ r \ge 3$ | $\{2^u, 2^v\},\quad 0 \le u < v \le r-1$ |
| II | $2^r,\ r \ge 2$ | $\{2^r - 1, 2^t\},\quad 0 \le t \le r-1$ |
| III | $2^r + 2^s - 2,\ r > s \ge 2$ | $\{2^r - 1, 2^s - 1\}$. |

(7)

*Proof.* The binary expansion identity $(x+1)^v = \prod_{j:\, v_j = 1} (x^{2^j} + 1)$ shows that $\Delta x^v$, for $v > 0$, consists of the proper binary submasks of $v$, and has $2^{\mathrm{wt}_2(v)} - 1$ terms. Consequently $\Delta x^v = 1$

exactly for powers of two. For distinct positive $v, w$, equality $\Delta x^v = \Delta x^w$ also occurs exactly when both are powers of two: with at least two binary digits, the singleton submasks recover all set digits. We will also use

$$x(x+1)\Delta S_t = x^{t+2} + (x+1)^{t+2} + 1. \tag{8}$$

It follows that $\Delta S_t = 0$ exactly when $t+2$ is a power of two.

For $0 \le b \le t$, the one-missing-coefficient polynomial $S_t + x^b$ is translation invariant exactly when

$$(t, b) = (2^s - 2, 0) \quad \text{or} \quad (2^s - 1, 2^s - 1), \qquad s \ge 1.$$

Indeed equality of its values at $0, 1$ forces $b = 0$ if $t$ is even. If $t$ is odd, translation invariance forces its leading term to cancel, since a nonzero invariant polynomial has even degree, so $b = t$. Equation (8) completes both cases, including the zero polynomial at $t = b = 0$.

Put $d = 2u$ and write $e = E^2 + xO^2$. Odd and even coefficient comparison gives $\Delta e = 0 \iff \Delta O = 0,\ \Delta E = O$. The useful residual is

$$B_u = S_{u-1} + \Delta S_u = \frac{x^{u+1} + (x+1)^{u+1} + 1}{x}.$$

If $a, b$ are even, then $O = S_{u-1}$. Thus $u+1$ is a power of two and $B_u = 0$. The remaining condition is $\Delta(x^{a/2} + x^{b/2}) = 0$, so $a/2, b/2$ are powers of two. These are precisely the even-even pairs in type I.

If $a, b$ are odd, then $E = S_u$ and $B_u = x^{(a-1)/2} + x^{(b-1)/2}$. Its two terms force $\mathrm{wt}_2(u+1) = 2$. Writing $u + 1 = 2^v + 2^w$, $v > w \ge 0$, gives $B_u = x^{2^v - 1} + x^{2^w - 1}$. This yields type III when $w > 0$; $w = 0$ yields type II with $t = 0$.

Finally suppose the missing positions are $2A$ and $2B+1$. Then $O = S_{u-1} + x^B$ and $B_u = x^B + \Delta x^A$. The one-missing-coefficient classification gives either $u = 2^s - 1, B = 0$, or $u = 2^s, B = u - 1$. In the first case $B_u = 0$; in the second $B_u = x^{u-1} + 1$. Both force $\Delta x^A = 1$, so $A$ is a power of two. They give respectively type I with one missing position 1, and type II with $t > 0$. These cases exhaust the parities. All displayed pairs satisfy the required equations on substitution, so the classification is sufficient as well as necessary. □

**Theorem 3.3** (Complete four-zero intersection)**.** *For the candidates* (1)*, $\mathcal{M}_n(f_{n,i,j}) = f_{n,i,j}$ if and only if one of the following holds:*

$$n = 2^R + 2^S - 2, \quad R > S \ge 2, \qquad (i, j) = (2^S - 2, 2^S - 1); \tag{9}$$

$$n = 2^R + 2, \quad R \ge 3, \qquad (i, j) = (3, 2^{R-1}); \tag{10}$$

$$n = 14, \qquad (i, j) \in \{(2, 4), (2, 5), (4, 5)\}. \tag{11}$$

*Equivalently, these are exactly the four-zero polynomials in the image of $\mathcal{T}$.*

*Proof.* Write $n = 2k$. The coefficient of $x^{n-1}$ in $\mathcal{M}_n(f)$ is $f'(1) = k - 2$ in $\mathbb{F}_2$, since the four missing tail positions include two odd positions. Its value in $f$ is 1, so $k$ is odd. Comparison of the $x^2$ coefficients gives $[x]f = \binom{2k}{2} = k = 1$. Thus positions $1, n-2$ are not missing. In Lemma 3.1, $e = S_d + x^a + x^b$ is therefore monic of even degree $d = k - 1$, with $1 \le a < b \le d - 1$.

By Lemma 3.2, $e$ has one of the three types in (7). The second descent condition is equivalent to

$$x^a + x^b + \Delta x^{d+1-a} + \Delta x^{d+1-b} = \frac{x^{d+2} + (x+1)^{d+2} + 1}{x}. \tag{12}$$

For type I the right side is zero. The binary masks $d+1-a$ and $d+1-b$ have all $r$ digits set except the distinct digits specified by $a, b$. Their proper-submask supports have symmetric difference of size $2^{r-1}-2$: among all submasks the difference has size $2^{r-1}$, and deleting the two top masks removes two terms. Equation (12) requires that difference to be the two terms $x^a + x^b$, so $r = 3$. For $r = 3$ it is exactly that pair, giving $d = 6$ and the three cases (11).

For type II put $\{a, b\} = \{d-1, 2^t\}$. The equation becomes

$$\Delta x^{d+1-2^t} = x^{2^t} + x + 1.$$

For $t = 0$ both sides are 1. For $t \geq 1$ the right side has three terms, while $d+1-2^t$ has $r-t+1$ set digits. Thus $2^{r-t+1}-1 = 3$, forcing $t = r-1$, which also suffices. The two choices yield respectively (9) with $S = 2$, and (10).

For type III the two $\Delta$ terms are both 1 and cancel. The right side is $x^a + x^b$, so every such pair satisfies the condition. Restoring the even zero positions $2a, 2b$ and their tail mirrors gives (9) with $R = r+1, S = s+1 \geq 3$. All these steps are reversible, proving the assertion in every degree. The equivalent image statement follows from the classical representation preceding Lemma 3.1. □

**Corollary 3.4** (Reducibility and failure of iteration)**.** *Every member of* (9) *is reducible. If $h$ is itself a four-zero polynomial of the form* (1)*, then $\mathcal{T}(h)$ is never a member of that family. Among the three polynomials in* (11)*, only $f_{14,2,4}$ is irreducible.*

*Proof.* For $f = \mathcal{T}(h)$ let $m = n/2$; its constant term gives $h(0) = 1$, and the $x^2$ coefficient gives

$$[x]h = [x^2]f + \binom{m}{2}. \tag{13}$$

In (9), $S = 2$ gives $m \equiv 1 \pmod 4$ and $[x^2]f = 0$, while $S \geq 3$ gives $m \equiv 3 \pmod 4$ and $[x^2]f = 1$. Hence $[x]h = 0$ in either case. If $h$ is reducible, multiplicativity of the transform makes $f$ reducible. If $h$ is irreducible, the classical binary $Q$ criterion [13, Theorem 1.1] says that $h^Q$, and hence $f = h^Q(x+1)$, is irreducible exactly when $[x]h = 1$. This proves the first assertion.

The theorem, or its coefficient proof, gives $n \equiv 2 \pmod 4$ for every four-zero output. A four-zero input has even degree, so its transformed degree is divisible by four. This proves the iteration obstruction for this particular transform.

For the exceptional pairs $(2,4), (2,5), (4,5)$ the corresponding $h$ are, respectively,

$$x^7 + x^6 + x^3 + x + 1, \quad x^7 + x^6 + x^3 + x^2 + x + 1, \quad x^7 + x^6 + x^4 + x^2 + 1.$$

The first satisfies $x^{128} \equiv x \pmod h$ and $\gcd(h, x^2 + x) = 1$, so is irreducible by the prime-degree Rabin test; its linear coefficient is 1. The second has root 1, and the third has zero linear coefficient. The same $Q$ criterion finishes the proof. □

**Explicit descent for the remaining family.** For (10), put $q = 2^{R-1}$. The unique polynomial with $\mathcal{T}(h_R) = f_{2q+2,3,q}$ is

$$h_R(x) = 1 + x + x^2 + (1+x)\left(x^q + \sum_{t=2}^{R-2} x^{q-2^t+1}\right), \qquad R \geq 3, \tag{14}$$

where the sum is empty for $R = 3$. To verify the formula uniformly, set $L(x) = \sum_{a=0}^{R-2} x^{2^a}$ and

$$F(x) = x^{q+2} h_R(1/x) = (x^2 + x + 1)x^q + (x+1)(L(x) + x^2).$$

For $s = X^{-1}$ and $w = s + s^2$, telescoping gives $L(w) = s + s^q$; substitution and collection of terms give

$$s^{-2q-4}F(w) = (X+1)f_{2q+2,3,q}(X).$$

The left side is $(X+1)\mathcal{T}(h_R)(X)$ by definition, proving (14). The distinct leading degrees of the transform basis give uniqueness. In particular, $\deg h_R = q+1$ and $[x]h_R = 1$, so the classical $Q$ criterion makes its irreducibility equivalent to that of the output. An all-parameter irreducibility classification remains unresolved.

### 3.2 The translation branch

We next impose $f(x+1) = f(x)$. The invariant representation $f = H(x^2+x)$ and its Artin–Schreier irreducibility criterion are classical [12, Lemma 2.1 and Theorem 2.2]. The additional constraints below come from the tail palindrome and its missing coefficients.

**Lemma 3.5** (Degree compatibility)**.** *If a monic tail-palindromic polynomial $f$ of even degree $n \geq 6$ satisfies $f(x+1) = f(x)$, then $n+2$ is a power of two.*

*Proof.* Let $F(X,Y) = Y^n f(X/Y)$ and $N = n+1$. Tail symmetry gives

$$XF(X,Y) + YF(Y,X) = X^N + Y^N.$$

By translation invariance, put $A = F(X,Y) = F(X+Y,Y)$, $B = F(Y,X) = F(X+Y,X)$ and $C = F(X,X+Y) = F(Y,X+Y)$. Applying the displayed identity to three pairs of variables yields

$$\begin{aligned} XA + YB &= X^N + Y^N, \\ (X+Y)A + YC &= (X+Y)^N + Y^N, \\ XC + (X+Y)B &= X^N + (X+Y)^N. \end{aligned}$$

Multiply these equations by $X+Y$, $X$ and $Y$, respectively, and add. All terms on the left cancel, leaving $(X+Y)^{n+2} = X^{n+2} + Y^{n+2}$. In characteristic two this forces $n+2$ to be a power of two: if its binary expansion has at least two nonzero digits, a singleton proper submask gives a nonzero mixed binomial term. □

**Proposition 3.6** (A missing-coefficient obstruction)**.** *Let $f$ be monic of even degree $n = 2k \geq 6$, with constant term one and a tail palindrome containing $2s$ zeros. If $f(x+1) = f(x)$, write $f = H(x^2+x)$ with $H$ monic of degree $k$. Then*

$$[x^{k-1}]H = s \pmod 2. \tag{15}$$

*In particular, if the number of tail zeros is divisible by four, then $f$ is reducible.*

*Proof.* With $e, o$ as in Lemma 3.1, tail symmetry gives $f = (x^k+e)^2 + xo^2$. Odd and even coefficient comparison gives

$$\Delta f = 0 \quad \Longleftrightarrow \quad \Delta o = 0, \qquad \Delta(x^k + e) = o. \tag{16}$$

Indeed $\Delta f = x(\Delta o)^2 + [\Delta(x^k+e) + o(x+1)]^2$. The formal coefficient vector of $o$ has length $k$ and $s$ zeros, so $[x^{n-2}]f = o(0) = o(1) = k - s$ in $\mathbb{F}_2$. Lemma 3.5 gives $k = 2^{R-1} - 1 \equiv 3 \pmod 4$, where $R \geq 3$. The classical coefficient comparison in $H(x^2+x)$ therefore gives

$$[x^{k-1}]H = [x^{2k-2}]f + \binom{k}{2} = s \pmod 2.$$

Now suppose that $s$ is even. If $H$ is reducible, so is its composition. Otherwise let $\beta$ be a root of the irreducible $H$ in $\mathbb{F}_{2^k}$. Its absolute trace is $[x^{k-1}]H = 0$, so $z^2 + z = \beta$ has a solution in that same field. Thus $f$ has a root of degree at most $k$ and is reducible. The last step is classical: specialize [11, Theorem 2.2(2)] to $p = 2$, $n = r = 1$ and $a = 1$ in Agou's notation, with irreducible outer polynomial $H$; see also [12, Theorem 2.2]. The same binary specialization follows from Long [10, Theorems 4.1–4.2, pp. 69 and 71]; the basic Artin–Schreier trace criterion is classical, as noted in [10, p. 63]. The associated high-coefficient restriction appears in [12, Theorem 2.8(3)]. □

The restriction $n \geq 6$ is essential: $x^2+x+1$ is a translation-invariant irreducible tail palindrome with no zeros. For odd $s$, (15) only removes this trace obstruction; it does not establish irreducibility of $H$.

**Theorem 3.7** (Complete translation intersection)**.** *The polynomial $f_{n,i,j}$ in* (1) *satisfies $f_{n,i,j}(x+1) = f_{n,i,j}(x)$ if and only if*

$$(n, i, j) \in \{(14, 3, 4), (30, 5, 9), (30, 5, 12), (30, 9, 12)\}. \tag{17}$$

*All four polynomials are reducible.*

*Proof.* In (16), put $d = k - 1$. The polynomial $o$ is monic of degree $d$ and has two missing coefficients. Since $k$ is odd, $o(1) = k - 2 = 1$; invariance gives $o(0) = 1$. Thus $o = S_d + x^a + x^b$ with $1 \leq a < b \leq d - 1$, and $e = S_d + x^{d-a} + x^{d-b}$. By Lemma 3.5, $d = 2^r - 2$, $r \geq 2$. The case $r = 2$ has only one interior position and is impossible. For $r \geq 3$, Equation (8) gives $\Delta S_d = 0$. Hence $\Delta x^a = \Delta x^b$, which, by the binary-support observation in the proof of Lemma 3.2, forces $a, b$ to be distinct powers of two. The second condition in (16), using $\Delta x^{d+1} = S_d$, becomes

$$\Delta x^{d-a} + \Delta x^{d-b} = x^a + x^b. \tag{18}$$

If $a, b$ are both even, take square roots in this equation. The two exponents on the left are $(r-1)$-digit all-one binary masks with different singleton digits removed. Their full submask supports have symmetric difference of size $2^{r-2}$; removing the two top masks leaves $2^{r-2} - 2$ terms. The right side has two terms, so $r = 4$. For three-digit masks the remaining difference is precisely the two removed singleton digits; thus all pairs from $\{2, 4, 8\}$ satisfy (18).

Otherwise $a = 1$ and $b$ is an even power of two. The term $\Delta x^{d-b}$ has only even exponents. The mask $d - 1 = 2^r - 3$ has all digits except digit one set, so $\Delta x^{d-1}$ has $2^{r-2} - 1$ odd terms. The right side has only the odd term $x$, forcing $r = 3$. Here $\Delta x^5 = x^4 + x + 1$ and $\Delta x^4 = \Delta x^2 = 1$; of $b = 2, 4$, exactly $b = 4$ satisfies the equation.

A missing position $a$ in $o$ corresponds to the two positions $2a + 1$ and $n - 2 - 2a$ in $f$. Thus $d = 6$, $\{a, b\} = \{1, 4\}$ gives $(14, 3, 4)$; $d = 14$ and the three pairs from $\{2, 4, 8\}$ give the degree-30 cases in (17). The substitutions above also prove sufficiency. Reducibility follows uniformly from Proposition 3.6 with $s = 2$. □

### 3.3 Scope of the quadratic construction

Here a full-degree quadratic pullback over $\mathbb{F}_2$ means

$$F = h^m P(g/h), \qquad \gcd(g, h) = 1, \qquad \max(\deg g, \deg h) = 2,$$

where $\deg P = m > 1$ and $\deg F = 2m$. The last condition is automatic for irreducible $P$; in general the degree drops exactly when $h_2 \neq 0$ and $P(g_2/h_2) = 0$ [8, Section 2].

**Corollary 3.8** (Boundary for full-degree quadratic pullbacks)**.** *If a full-degree quadratic pullback is a four-zero tail palindrome, then it belongs either to the lists in Theorem 3.3 or to* (17)*, and its degree is* 2 *modulo* 4*. If it is irreducible, it is either* $f_{14,2,4}$ *or a member of* (10)*. Consequently, no sequence of quadratic pullbacks, each doubling the degree, can reach a four-zero tail palindrome after starting at an even degree.*

*Proof.* Write $g = \sum_{j=0}^{2} g_j x^j$, $h = \sum_{j=0}^{2} h_j x^j$, and put

$$a = g_2h_1 + g_1h_2, \quad b = g_0h_2 + g_2h_0, \quad c = g_1h_0 + g_0h_1.$$

The classical quadratic framework gives $b^2 + ac = 1$ and, in the separable case,

$$(ax+b)^{2m} F\left(\frac{bx+c}{ax+b}\right) = F(x) \qquad \text{[8, Section 4, Lemma 9]}.$$

In the separable case the possible triples are exactly $(1,0,1), (0,1,1), (1,1,0)$, corresponding to reciprocity, translation and $\mathcal{M}_{2m}$. There is no nontrivial scalar normalization over $\mathbb{F}_2$. In the inseparable case $g_1 = h_1 = 0$, the output is a square; its odd coefficients vanish, whereas a four-zero candidate has $[x^{2m-1}]F = 1$.

The reciprocal branch contains no four-zero candidate: combining $f^* = f$ with Lemma 2.1 forces $f = S_n$. The other two branches are classified by Theorems 3.3 and 3.7. The irreducible alternatives follow from their reducibility conclusions. All listed degrees are 2 modulo 4. Starting at an even degree, every subsequent doubled degree is divisible by four, proving the last assertion. □

The reduction to three involutions is classical; the complete support intersections impose the additional construction restrictions. Higher-degree maps, selected factors, degree drops and extension-field descent are outside this conclusion.

# 4 Independently verified finite data

All 161700 candidates at the 98 even degrees $6 \le n \le 200$ were independently retested. The total is

$$\sum_{\substack{6 \le n \le 200 \\ n \text{ even}}} N_4(n) = 2663, \qquad N_4(n) > 0 \text{ at every tested degree.}$$

The audit reproduces every historical complete count and every lexicographically first pair. For $202 \le n \le 300$, it retests the lexicographic prefix through the recorded first solution, covering 5410 additional candidates at 50 degrees. This verifies the first pair at each such degree; it does not provide complete counts beyond degree 200.

**Arithmetic verification and reproducibility.** The independent code constructs coefficients directly and tests irreducibility, followed by the standard exact-order conditions

$$x^{2^n-1} \equiv 1 \pmod f, \qquad x^{(2^n-1)/p} \not\equiv 1 \pmod f \quad (p \mid 2^n - 1 \text{ prime}). \tag{19}$$

It imports none of the historical polynomial routines and uses python-flint 0.9.0 with FLINT 3.6.0 [14]. All 148 integer order factorizations are checked by exact division with residual cofactor one. Their 484 distinct factors, of at most 144 bits, pass FLINT's rigorous `fmpz_is_prime` procedure. A standard-library implementation independently agrees on all 220 candidates through degree 24.

| $n$ | candidates | $N_4(n)$ | first $(i,j)$ |
|---|---|---|---|
| 6 | 1 | 1 | $(1,2)$ |
| 8 | 3 | 1 | $(2,3)$ |
| 10 | 6 | 2 | $(1,2)$ |
| 12 | 10 | 3 | $(1,3)$ |
| 20 | 36 | 1 | $(4,5)$ |
| 40 | 171 | 11 | $(1,4)$ |
| 62 | 435 | 16 | $(1,17)$ |
| 100 | 1176 | 31 | $(1,23)$ |
| 126 | 1891 | 55 | $(1,2)$ |
| 154 | 2850 | 45 | $(1,23)$ |
| 194 | 4560 | 20 | $(6,65)$ |
| 200 | 4851 | 17 | $(1,71)$ |

Table 1: Selected complete counts. All 98 rows and all successful position pairs through degree 200 are supplied with the verification data.

The folder `verification/primitivity/` supplies complete candidate verdicts, successful positions, exact order-test residues for the 148 first witnesses, recovered integer factorizations and input hashes. Historical inputs are preserved separately. Sixteen missing auxiliary entries for the first successful $j$ with $i = 1$ were corrected; complete counts and first pairs are unchanged. Software primality decisions are not exported independent certificates; no Lean formalization is claimed. The data and source are available from the author.

**Structural checks.** The accompanying `verification/` folders record the following scopes:

| Audit | Finite verification range |
|---|---|
| Reciprocal screen | 19600 candidates, $n \le 100$. |
| Coefficient quotient | 341376 candidates, $n \le 256$; full character checks for $k \le 20$. |
| Translated $Q$ | 341376 four-zero positions at output degrees $\le 256$; 1389888 two-zero positions at input degrees $\le 256$. |
| Independent $Q$ | Horner/Pascal: 4960 four-zero positions at output degrees $\le 64$; 72568 two-zero positions at input degrees $\le 96$; 1023 general tail palindromes to $n = 20$. |
| Translation | 341376 four-zero positions to $n = 256$; 4960 direct Horner checks to $n = 64$; 1023 general tail palindromes; 24 quadratic maps; 508 Artin–Schreier instances of input degrees 2–8. |
| Explicit B descent | Standard-library forward and inverse checks for $3 \le R \le 10$; no factorization search. |

The four translation outputs have independently certified factor degrees $2+12$, $12+18$, $6+24$ and $7+7+8+8$, with a repeated factor in the last. Earlier B-sequence factorization data cover only $3 \le R \le 10$: the first term is irreducible and the next seven reducible. The new descent checks and unchanged historical audits are distinguished in the supplied files. All-parameter results rest on the written proofs; these finite checks imply no assertion for later parameters.

## 5 Interpretation and remaining question

The main results give family-specific support restrictions within classical quadratic-transform theory. The reciprocal screen is necessary, and the classifications concern complete outputs of full

degree. Within this scope, any infinite irreducible output family would have to occur in (10); its existence remains unproved. The auxiliary coefficient quotient does not control multiplicative order or give a primitive-count estimate. The finite data support Conjecture 1.1 only over the stated ranges. A further substantive step would require an effective existence argument or a construction with controlled root order.

# A Auxiliary classical descriptions

These descriptions record coefficient multiplicities and an exact field parametrization. Neither is used to prove the support classifications.

## A.1 Coefficient quotient

Let $\rho$ reverse $\mathbb{F}_2^k$, and set

$$P = \{(A, \rho A) : A \in \mathbb{F}_2^k\}, \quad K = \{(C, C) : C = \rho C\}, \quad U = P/K, \quad r = \lfloor k/2 \rfloor.$$

The map $[(A, \rho A)] \mapsto A + \rho A$ identifies $U$ with $\mathrm{im}(I + \rho)$, of dimension $r$. Prepending a zero to $x \in \mathbb{F}_2^{k-1}$ defines $\Phi(x) = [(\widetilde{x}, \rho\widetilde{x})]$. The all-ones tail lies in $K$, so deleting positions $i, j$ gives class $\Phi(e_i + e_j)$. With zero-based coordinates and a suitable basis $z_1, \ldots, z_r$, the columns $u_i = \Phi(e_i)$ form the multiset

$$\{u_1, \ldots, u_{k-1}\} = \begin{cases} \{z_1, z_1, \ldots, z_{r-1}, z_{r-1}, z_r\}, & k = 2r, \\ \{z_1, z_1, \ldots, z_{r-1}, z_{r-1}, z_r, 0\}, & k = 2r + 1. \end{cases} \tag{20}$$

Indeed pair positions $i, k-1-i$; the boundary pair contributes the singleton $z_r$ and an unpaired middle position contributes zero. In particular $\Phi$ is onto.

For $s = \sum s_a z_a$, write $h = \sum_{a<r} s_a$ as an integer and $\varepsilon = k \bmod 2$. The ordinary coset weight enumerator is

$$A_s(z) = \sum_{\Phi(x)=s} z^{\mathrm{wt}(x)} = (1+z)^{\varepsilon} z^{s_r} (2z)^h (1+z^2)^{r-1-h}. \tag{21}$$

Each repeated column allows $00, 11$ for zero syndrome and $01, 10$ for nonzero syndrome; the singleton and optional zero column give the remaining factors. Thus $\mu(s) = [z^2]A_s(z)$ counts four-zero candidates in class $s$, and the number of occupied classes is

$$D(3) = 1, \qquad D(k) = 1 + \binom{\lfloor (k+1)/2 \rfloor}{2} \quad (k \geq 4). \tag{22}$$

This follows by counting the nonzero quadratic coefficients in (21); at $k = 3$ the zero class is unoccupied. For a character $\chi$ of $U$, put $t = \#\{i : \chi(u_i) = -1\}$ and $\ell = k - 1$. Then

$$\widehat{\mu}(\chi) = \sum_{i<j} \chi(u_i)\chi(u_j) = K_2(t; \ell) = \frac{(\ell - 2t)^2 - \ell}{2}. \tag{23}$$

These are classical direct-product weight-enumerator and Krawtchouk identities [7, Sections 3–4][6, Sections 1.1 and 2.1]. They do not decide irreducibility: at degree eight, $x^8 + x^7 + x^6 + x + 1$ is primitive, while $x^8 + x^7 + x^5 + x^2 + 1 = (x^2 + x + 1)(x^6 + x^4 + x^2 + x + 1)$ is reducible; their tail difference $0110\,0110$ belongs to $K$.

### A.2 Finite-field parametrization

Put $q = 2^k$, $\mathbb{K} = \mathbb{F}_q$, $\mathbb{L} = \mathbb{F}_{q^2}$ and $T = \{\gamma \in \mathbb{L}^* : \gamma^{q+1} = 1\}$. The coprime orders $q-1, q+1$ give $\mathbb{L}^* = \mathbb{K}^* \times T$. A primitive element is uniquely $\beta = \nu\gamma$ with $\mathrm{ord}(\nu) = q-1$ and $\mathrm{ord}(\gamma) = q+1$. Writing $\tau = \gamma + \gamma^{-1} \in \mathbb{K}$, the standard Dickson identity $D_a(u + u^{-1}) = u^a + u^{-a}$ and Frobenius pairing give

$$F_{\nu,\gamma}(x) = \prod_{i=0}^{k-1} \left( x^2 + \nu^{2^i} D_{2^i}(\tau) x + \nu^{2^{i+1}} \right). \tag{24}$$

Here $D_0 = 0, D_1 = t, D_{a+1} = tD_a + D_{a-1}$ in characteristic two [2]. Conversely, the two order conditions make this product the minimal polynomial of a primitive element. Expanding the product and comparing with (1) gives an exact coefficient-and-order system, without an existence conclusion.

## References


[1] S. D. Cohen, *Primitive polynomials with a prescribed coefficient*, Finite Fields Appl. **12** (2006), 425–491.

[2] R. Lidl and H. Niederreiter, *Finite Fields*, Encyclopedia of Mathematics and its Applications, vol. 20, 2nd ed., Cambridge University Press, 1997.

[3] R. G. Swan, *Factorization of polynomials over finite fields*, Pacific J. Math. **12** (1962), 1099–1106.

[4] H. M. Fredricksen, A. W. Hales, and M. M. Sweet, *A generalization of Swan's theorem*, Math. Comp. **46** (1986), 321–331.

[5] C. Defant, *Enumerating cliques in direct product graphs*, J. Combin. **11** (2020), 351–358.

[6] A. E. Brouwer, S. M. Cioabă, F. Ihringer, and M. McGinnis, *The smallest eigenvalues of Hamming graphs, Johnson graphs and other distance-regular graphs with classical parameters*, J. Combin. Theory Ser. B **133** (2018), 88–121. doi:10.1016/j.jctb.2018.04.005.

[7] G. D. Forney, Jr., *Codes on graphs: duality and MacWilliams identities*, IEEE Trans. Inform. Theory **57** (2011), 1382–1397. doi:10.1109/TIT.2011.2104994.

[8] S. Mattarei and M. Pizzato, *Generalizations of self-reciprocal polynomials*, Finite Fields Appl. **48** (2017), 271–288. doi:10.1016/j.ffa.2017.08.004.

[9] L. Reis, *Möbius-like maps on irreducible polynomials and rational transformations*, J. Pure Appl. Algebra **224** (2020), 169–180. doi:10.1016/j.jpaa.2019.05.001.

[10] A. F. Long, Jr., *Factorization of irreducible polynomials over a finite field with the substitution $x^{p^r} - x$ for $x$*, Duke Math. J. **40** (1973), 63–76. doi:10.1215/S0012-7094-73-04007-6.

[11] S. Agou, *Factorisation sur un corps fini $\mathbb{F}_{p^n}$ des polynômes composés $f(X^{p^r} - aX)$ lorsque $f(X)$ est un polynôme irréductible de $\mathbb{F}_{p^n}(X)$*, J. Number Theory **9** (1977), 229–239. Publisher page.

[12] G. Kapetanakis, *Prescribing coefficients of invariant irreducible polynomials*, J. Number Theory **180** (2017), 615–628. doi:10.1016/j.jnt.2017.05.003.

[13] S. Ugolini, *Sequences of binary irreducible polynomials*, Discrete Math. **313** (2013), 2656–2662. doi:10.1016/j.disc.2013.08.011.

[14] The python-flint and FLINT developers, *python-flint*, version 0.9.0, with FLINT version 3.6.0. Documentation; integer primality-testing documentation. Accessed 27 October 2026.