

# Correction of midterm exam n°2

## Exercise 1 (4,5 points)

1. Using an integration by parts, by setting  $u(x) = \sin(\ln(x))$  and  $v'(x) = 1$ , we get :

$$\begin{aligned} I &= \left[ x \sin(\ln(x)) \right]_1^e - \int_1^e \cos(\ln(x)) \, dx \\ &= e \sin(1) - \int_1^e \cos(\ln(x)) \, dx \end{aligned}$$

Using a second integration by parts, by setting  $u(x) = \cos(\ln(x))$  and  $v'(x) = 1$ , we get :

$$I = e \sin(1) - \left( \left[ x \cos(\ln(x)) \right]_1^e + I \right)$$

thus  $2I = 1 + e(\sin(1) - \cos(1))$ , that is to say  $I = \frac{1}{2}(1 + e(\sin(1) - \cos(1)))$ .

2. Using an integration by parts, by setting  $u(x) = \arctan(x)$  and  $v'(x) = 1$ , we get :

$$\begin{aligned} J &= \left[ x \arctan(x) \right]_0^1 - \int_0^1 \frac{x}{1+x^2} \, dx \\ &= \frac{\pi}{4} - \frac{1}{2} [\ln(1+x^2)]_0^1 \\ &= \frac{\pi}{4} - \frac{1}{2} \ln(2) \end{aligned}$$

3. Using the substitution  $u = \sqrt{x}$ ,  $x = u^2$  thus  $dx = 2u \, du$ .

We get :  $K = 2 \int_0^\pi u \cos(u) \, du$ .

Using an integration by parts, by setting  $\varphi(u) = u$  and  $\psi'(u) = \cos(u)$ , we get :

$$K = 2 \left( \left[ u \sin(u) \right]_0^\pi - \int_0^\pi \sin(u) \, du \right) = -2 \int_0^\pi \sin(u) \, du$$

Thus  $K = -4$ .

## Exercise 2 (3 points)

1. From the hypothesis we deduce that 
$$\left\{ \begin{array}{l} \frac{u_n}{u_{n-1}} \leq \frac{v_n}{v_{n-1}} \\ \vdots \\ \frac{u_1}{u_0} \leq \frac{v_1}{v_0} \end{array} \right.$$

By multiplying these inequalities of positive numbers, we get :

$$\frac{u_n}{u_{n-1}} \cdot \frac{u_{n-1}}{u_{n-2}} \cdot \dots \cdot \frac{u_1}{u_0} \leq \frac{v_n}{v_{n-1}} \cdot \frac{v_{n-1}}{v_{n-2}} \cdot \dots \cdot \frac{v_1}{v_0}$$

from which we deduce that  $\frac{u_n}{u_0} \leq \frac{v_n}{v_0}$ , that is to say  $0 < u_n \leq \frac{u_0}{v_0} v_n$ .

Therefore, using the squeeze theorem (sandwich theorem), if  $v_n \xrightarrow[n \rightarrow +\infty]{} 0$  then  $u_n \xrightarrow[n \rightarrow +\infty]{} 0$ .

2. Similarly,  $v_n \geq \frac{v_0}{u_0} u_n$ .

Thus, still using the squeeze theorem, if  $u_n \xrightarrow[n \rightarrow +\infty]{} +\infty$  then  $v_n \xrightarrow[n \rightarrow +\infty]{} +\infty$ .

### Exercise 3 (3 points)

- a. Let  $(u_n)$  be a sequence of real numbers, and  $\ell \in \mathbb{R}$ . Then the assertion « if  $(u_n)$  converges towards  $\ell$  then, for every  $n \in \mathbb{N}$ ,  $u_n \leq \ell$  » is equivalent to the assertion « if there exists  $n \in \mathbb{N}$  such that  $u_n > \ell$ , then  $(u_n)$  does not converge towards  $\ell$  ».
- b. If  $(u_n)$  is a nonzero geometric sequence with common ratio  $q \in \mathbb{R}^*$ , then  $\left(\frac{1}{u_n}\right)$  is a geometric sequence with common ratio  $\frac{1}{q}$ .
- c. If  $(u_n)$  is a bounded numerical sequence, there exists a subsequence of  $(u_n)$  that is convergent.
- d. Let  $(u_n)$  be a numerical sequence. Then  $(u_{6n})$  is a subsequence of  $(u_{2n})$ .
- e. Let  $(u_n)$  be a numerical sequence. Then  $(u_{3 \cdot 2^{n+1}})$  is a subsequence of  $(u_{6n})$ .
- f. none of the above

### Exercise 4 (3 points)

Let us prove that  $(u_n)$  is (strictly) increasing.

$$u_{n+1} - u_n = \sum_{k=0}^{2n+3} \frac{(-1)^k}{(2k)!} - \sum_{k=0}^{2n+1} \frac{(-1)^k}{(2k)!} = \frac{(-1)^{2n+2}}{(2(2n+2))!} + \frac{(-1)^{2n+3}}{(2(2n+3))!} = \frac{1}{(4n+4)!} - \frac{1}{(4n+6)!}$$

Yet

$$(4n+6)! > (4n+4)!$$

hence

$$\frac{1}{(4n+6)!} < \frac{1}{(4n+4)!}$$

Therefore  $u_{n+1} - u_n > 0$ , thus  $(u_n)$  is strictly increasing.

Let us prove that  $(v_n)$  is (strictly) decreasing.

$$v_{n+1} - v_n = u_{n+1} + \frac{1}{(4(n+1)+4)!} - u_n - \frac{1}{(4n+4)!}$$

By reusing the already calculated expression of  $u_{n+1} - u_n$ , we get :

$$v_{n+1} - v_n = \frac{1}{(4n+4)!} - \frac{1}{(4n+6)!} + \frac{1}{(4n+8)!} - \frac{1}{(4n+4)!} = \frac{1}{(4n+8)!} - \frac{1}{(4n+6)!}$$

Yet

$$(4n+8)! > (4n+6)!$$

hence

$$\frac{1}{(4n+8)!} < \frac{1}{(4n+6)!}$$

Therefore  $v_{n+1} - v_n < 0$  thus  $(v_n)$  is strictly decreasing.

Eventually,  $v_n - u_n = \frac{1}{(4n+4)!} \rightarrow 0$  thus  $(u_n)$  and  $(v_n)$  are adjacent sequences.

### Exercise 5 (2 points)

- $\ln(n!) = \ln(1) + \dots + \ln(n) = \sum_{k=1}^n \ln(k) \leq n \ln(n)$  as the function  $\ln$  is increasing.
- Thus, for every  $n \in \mathbb{N}^*$ ,  $0 \leq u_n \leq \frac{\ln(n)}{n}$ .

Using the squeeze theorem together with a result of compared growth,  $(u_n)_{n \in \mathbb{N}^*}$  converges towards 0.

## Exercise 6 (5,5 points)

1. As  $q \neq 1$ ,  $\sum_{k=1}^n q^{k-1} = \frac{1-q^n}{1-q}$
2. Using the previous question,

$$\sum_{k=1}^n \frac{1}{2^{k-1}} = \sum_{k=1}^n \left(\frac{1}{2}\right)^{k-1} = \frac{1 - \left(\frac{1}{2}\right)^n}{1 - \frac{1}{2}} = 2 \left(1 - \frac{1}{2^n}\right)$$

that is to say

$$\sum_{k=1}^n \frac{1}{2^{k-1}} = 2 - \frac{1}{2^{n-1}}$$

$$3. \text{ Let } k \geq 2. \text{ Then : } \begin{cases} 2 \geq 2 \\ 3 \geq 2 \\ \vdots \\ k \geq 2 \end{cases} \text{ hence } \begin{cases} \frac{1}{2} \leq \frac{1}{2} \\ \frac{1}{3} \leq \frac{1}{2} \\ \vdots \\ \frac{1}{k} \leq \frac{1}{2} \end{cases}$$

$$\text{Thus, } \frac{1}{k!} = \frac{1}{2 \times 3 \times \dots \times k} = \frac{1}{2} \times \frac{1}{3} \times \dots \times \frac{1}{k} \leq \frac{1}{2^{k-1}}$$

This property still holds when  $k = 1$  as  $\frac{1}{1!} = 1 \leq \frac{1}{2^0} = 1$ .

4. Let  $n \in \mathbb{N}$ . Then  $u_{n+1} - u_n = \frac{1}{(n+1)!} \geq 0$  thus  $(u_n)$  is increasing.
5. Using questions 2 and 3,

$$u_n = 1 + \sum_{k=1}^n \frac{1}{k!} \leq 1 + \sum_{k=1}^n \frac{1}{2^{k-1}} \leq 1 + 2 - \frac{1}{2^{n-1}} \leq 3$$

6.  $(u_n)$  is an increasing numerical sequence that is bounded above, thus it is convergent.