

Signal Processing

Fourier Analysis on Finite Abelian Groups

6.S096

January 30, 2024

- **Today's Content:** Fourier Analysis on Finite Abelian Group.

- **Today's Content:** Fourier Analysis on Finite Abelian Group.
- **Goal:** Decompose Functions on Finite Abelian Groups, analogous to the continuous case.

- **Today's Content:** Fourier Analysis on Finite Abelian Group.
- **Goal:** Decompose Functions on Finite Abelian Groups, analogous to the continuous case.
- We will mostly follow Bao Luong's eponymous textbook.
- Classes at MIT?

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

① **Commutativity:** $u + v = v + u$ for all $u, v \in V$

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

- ① **Commutativity:** $u + v = v + u$ for all $u, v \in V$
- ② **Associativity:** $(u + v) + w = u + (v + w)$ and $a \cdot (b \cdot v) = (a \cdot b) \cdot v$ for all $u, v, w \in V$ and all $a, b \in \mathbb{C}$

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

- ① **Commutativity:** $u + v = v + u$ for all $u, v \in V$
- ② **Associativity:** $(u + v) + w = u + (v + w)$ and $a \cdot (b \cdot v) = (a \cdot b) \cdot v$ for all $u, v, w \in V$ and all $a, b \in \mathbb{C}$
- ③ **Additive Identity:** There exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

- ① **Commutativity:** $u + v = v + u$ for all $u, v \in V$
- ② **Associativity:** $(u + v) + w = u + (v + w)$ and $a \cdot (b \cdot v) = (a \cdot b) \cdot v$ for all $u, v, w \in V$ and all $a, b \in \mathbb{C}$
- ③ **Additive Identity:** There exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$
- ④ **Additive Inverse:** For every $v \in V$, there exists $w \in V$ such that $v + w = 0$
- ⑤ **Multiplicative Identity:** $1 \cdot v = v$ for all $v \in V$

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

- ❶ **Commutativity:** $u + v = v + u$ for all $u, v \in V$
- ❷ **Associativity:** $(u + v) + w = u + (v + w)$ and $a \cdot (b \cdot v) = (a \cdot b) \cdot v$ for all $u, v, w \in V$ and all $a, b \in \mathbb{C}$
- ❸ **Additive Identity:** There exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$
- ❹ **Additive Inverse:** For every $v \in V$, there exists $w \in V$ such that $v + w = 0$
- ❺ **Multiplicative Identity:** $1 \cdot v = v$ for all $v \in V$
- ❻ **Distributive Properties:** $a \cdot (u + v) = a \cdot u + a \cdot v$ and $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in F$ and all $u, v \in V$

Definition of Vector Space

Definition (Vector Space)

[?] A vector space is a set V equipped with an $+$ and scalar multiplication such that

- ❶ **Commutativity:** $u + v = v + u$ for all $u, v \in V$
- ❷ **Associativity:** $(u + v) + w = u + (v + w)$ and $a \cdot (b \cdot v) = (a \cdot b) \cdot v$ for all $u, v, w \in V$ and all $a, b \in \mathbb{C}$
- ❸ **Additive Identity:** There exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$
- ❹ **Additive Inverse:** For every $v \in V$, there exists $w \in V$ such that $v + w = 0$
- ❺ **Multiplicative Identity:** $1 \cdot v = v$ for all $v \in V$
- ❻ **Distributive Properties:** $a \cdot (u + v) = a \cdot u + a \cdot v$ and $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in F$ and all $u, v \in V$

Example

The space of complex polynomials of degree at most n , denoted as $\mathbb{C}[x]_{\leq n}$, is a complex vector space. Addition is polynomial addition, and scalar multiplication is polynomial scalar multiplication.

Example

The space of complex polynomials of degree at most n , denoted as $\mathbb{C}[x]_{\leq n}$, is a complex vector space. Addition is polynomial addition, and scalar multiplication is polynomial scalar multiplication. This vector space has a basis $\{1, x, x^2, \dots, x^n\}$ and so it has dimension $n + 1$.

Example

The space of complex polynomials of degree at most n , denoted as $\mathbb{C}[x]_{\leq n}$, is a complex vector space. Addition is polynomial addition, and scalar multiplication is polynomial scalar multiplication. This vector space has a basis $\{1, x, x^2, \dots, x^n\}$ and so it has dimension $n + 1$.

We also introduce the notion

Inner Product Space

Let V be a complex vector space. An *inner product* on V is a function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$ that satisfies the following properties for all vectors \mathbf{u} , \mathbf{v} , and scalars $a, b \in \mathbb{C}$:

- 1 Conjugate Symmetry: $\langle \mathbf{u}, \mathbf{v} \rangle = \overline{\langle \mathbf{v}, \mathbf{u} \rangle}$.
- 2 Linearity in the First Argument: $\langle a\mathbf{u} + b\mathbf{v}, \mathbf{w} \rangle = a\langle \mathbf{u}, \mathbf{w} \rangle + b\langle \mathbf{v}, \mathbf{w} \rangle$.
- 3 Positive Definiteness: $\langle \mathbf{u}, \mathbf{u} \rangle \geq 0$ with equality if and only if $\mathbf{u} = \mathbf{0}$.

Last Ounce of Linear Algebra

Definition

A linear transformation of vector spaces is a map $T : V \rightarrow W$ which respects addition and scalar multiplication.

Last Ounce of Linear Algebra

Definition

A linear transformation of vector spaces is a map $T : V \rightarrow W$ which respects addition and scalar multiplication.

Definition

Let V and W be inner product spaces with inner products $\langle \cdot, \cdot \rangle_V$ and $\langle \cdot, \cdot \rangle_W$, respectively.

Last Ounce of Linear Algebra

Definition

A linear transformation of vector spaces is a map $T : V \rightarrow W$ which respects addition and scalar multiplication.

Definition

Let V and W be inner product spaces with inner products $\langle \cdot, \cdot \rangle_V$ and $\langle \cdot, \cdot \rangle_W$, respectively. A linear transformation $T : V \rightarrow W$ is called an *isometry* if, for all vectors $u, v \in V$, the following condition holds:

$$\langle T(u), T(v) \rangle_W = \langle u, v \rangle_V.$$

Last Ounce of Linear Algebra

Definition

A linear transformation of vector spaces is a map $T : V \rightarrow W$ which respects addition and scalar multiplication.

Definition

Let V and W be inner product spaces with inner products $\langle \cdot, \cdot \rangle_V$ and $\langle \cdot, \cdot \rangle_W$, respectively. A linear transformation $T : V \rightarrow W$ is called an *isometry* if, for all vectors $u, v \in V$, the following condition holds:

$$\langle T(u), T(v) \rangle_W = \langle u, v \rangle_V.$$

Another special property of inner-products is that they allow one to compute coordinates against a basis.

Last Ounce of Linear Algebra

Definition

A linear transformation of vector spaces is a map $T : V \rightarrow W$ which respects addition and scalar multiplication.

Definition

Let V and W be inner product spaces with inner products $\langle \cdot, \cdot \rangle_V$ and $\langle \cdot, \cdot \rangle_W$, respectively. A linear transformation $T : V \rightarrow W$ is called an *isometry* if, for all vectors $u, v \in V$, the following condition holds:

$$\langle T(u), T(v) \rangle_W = \langle u, v \rangle_V.$$

Another special property of inner-products is that they allow one to compute coordinates against a basis. If b_1, \dots, b_n is an orthonormal basis then

$$v = \sum_{i=1}^n \langle v, b_i \rangle b_i.$$

Functions on a Set

Definition

Consider a finite nonempty set S .

Functions on a Set

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Functions on a Set

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

$$\langle f, g \rangle = \sum_{s \in S} f(s) \overline{g(s)}.$$

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

$$\langle f, g \rangle = \sum_{s \in S} f(s) \overline{g(s)}.$$

Naturally, one can ask what is the dimension of V_S .

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

$$\langle f, g \rangle = \sum_{s \in S} f(s) \overline{g(s)}.$$

Naturally, one can ask what is the dimension of V_S . The dimension is exactly $|S|$: consider the basis of Dirac delta functions

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

$$\langle f, g \rangle = \sum_{s \in S} f(s) \overline{g(s)}.$$

Naturally, one can ask what is the dimension of V_S . The dimension is exactly $|S|$: consider the basis of Dirac delta functions

$$\delta_s(x) = \begin{cases} 1 & \text{if } x = s, \\ 0 & \text{if } x \neq s. \end{cases}$$

Functions on a Set

Definition

Consider a finite nonempty set S . Let V_S be the space of complex-valued functions on S , that is, the space of functions $S \rightarrow \mathbb{C}$.

Note that this is equivalent to specifying a tuple in $\mathbb{C}^{|S|}$. In fact, V_S is a complex vector space which can be equipped with an inner product

$$\langle f, g \rangle = \sum_{s \in S} f(s) \overline{g(s)}.$$

Naturally, one can ask what is the dimension of V_S . The dimension is exactly $|S|$: consider the basis of Dirac delta functions

$$\delta_s(x) = \begin{cases} 1 & \text{if } x = s, \\ 0 & \text{if } x \neq s. \end{cases}$$

Let Δ_S be the set of such δ_s functions. This implies that V_S is a $|S|$ -dimensional complex vector space.

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication.

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Definition (Characters)

A character of a group G is a group homomorphism χ from G to \mathbb{C}^* .

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Definition (Characters)

A character of a group G is a group homomorphism χ from G to \mathbb{C}^* .

The character corresponding to the trivial homomorphism is called the *trivial* (or *principal*) character.

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Definition (Characters)

A character of a group G is a group homomorphism χ from G to \mathbb{C}^* .

The character corresponding to the trivial homomorphism is called the *trivial* (or *principal*) character.

Example

If χ is a character on G and $g \in G$, then $g^{|G|} = 1$ and so $\chi(g)^{|G|} = 1$. What does this say about the image of χ ?

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Definition (Characters)

A character of a group G is a group homomorphism χ from G to \mathbb{C}^* .

The character corresponding to the trivial homomorphism is called the *trivial* (or *principal*) character.

Example

If χ is a character on G and $g \in G$, then $g^{|G|} = 1$ and so $\chi(g)^{|G|} = 1$. What does this say about the image of χ ?

Fact (Characters form a group)

The characters on a group G form a group under pointwise multiplication.

Characters

Recall that \mathbb{C}^* is the group of complex numbers under multiplication. Also, recall that an *abelian* group is a group G in which the group operation is commutative, so $xy = yx$ for all $x, y \in G$.

Definition (Characters)

A character of a group G is a group homomorphism χ from G to \mathbb{C}^* .

The character corresponding to the trivial homomorphism is called the *trivial* (or *principal*) character.

Example

If χ is a character on G and $g \in G$, then $g^{|G|} = 1$ and so $\chi(g)^{|G|} = 1$. What does this say about the image of χ ?

Fact (Characters form a group)

The characters on a group G form a group under pointwise multiplication.

Characters II

This group is called the *dual* group of G , and is denoted as \hat{G} .

Characters II

This group is called the *dual* group of G , and is denoted as \hat{G} .

Theorem

If G is a finite abelian group, then G and \hat{G} are isomorphic.

Let us sketch the proof of this result. For this, we will need the following result.

Characters II

This group is called the *dual* group of G , and is denoted as \hat{G} .

Theorem

If G is a finite abelian group, then G and \hat{G} are isomorphic.

Let us sketch the proof of this result. For this, we will need the following result.

Theorem (Structure Theorem for Finite Abelian Groups)

Every finite abelian group G can be expressed^a as the direct product of cyclic groups:

Characters II

This group is called the *dual* group of G , and is denoted as \hat{G} .

Theorem

If G is a finite abelian group, then G and \hat{G} are isomorphic.

Let us sketch the proof of this result. For this, we will need the following result.

Theorem (Structure Theorem for Finite Abelian Groups)

Every finite abelian group G can be expressed^a as the direct product of cyclic groups:

$$G \cong \mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2} \times \dots \times \mathbb{Z}_{n_k}$$

where n_1, n_2, \dots, n_k are positive integers whose product is $|G|$.

^ais isomorphic

Finishing the Proof

Moreover, one can verify that the group of characters on $\widehat{G_1 \times G_2}$ is isomorphic $\widehat{G_1} \times \widehat{G_2}$, so this boils down to the following result.

Lemma

The dual $\widehat{\mathbb{Z}_n}$ is isomorphic to \mathbb{Z}_n for all $n \geq 1$.

How do characters relate to each other?

Theorem (Orthogonality Relation I)

$$\sum_{g \in G} \chi(g) = \begin{cases} |G| & \text{if } \chi \text{ is trivial,} \\ 0 & \text{otherwise.} \end{cases}$$

How do characters relate to each other?

Theorem (Orthogonality Relation I)

$$\sum_{g \in G} \chi(g) = \begin{cases} |G| & \text{if } \chi \text{ is trivial,} \\ 0 & \text{otherwise.} \end{cases}$$

This gives rise to an orthogonality result.

Corollary (Orthogonality of Characters)

$$\langle \chi, \chi' \rangle = \begin{cases} |G| & \text{if } \chi = \chi', \\ 0 & \text{if } \chi \neq \chi'. \end{cases}$$

More orthogonality relations.

We can also compute a slightly different sum (where we fix the group element and vary the character).

More orthogonality relations.

We can also compute a slightly different sum (where we fix the group element and vary the character).

Corollary (Summing over characters)

If $g \in G$ then

$$\sum_{\chi \in \widehat{G}} \chi(g) = \begin{cases} |G| & \text{if } g = 1_G \\ 0 & \text{otherwise.} \end{cases}$$

More orthogonality relations.

We can also compute a slightly different sum (where we fix the group element and vary the character).

Corollary (Summing over characters)

If $g \in G$ then

$$\sum_{\chi \in \widehat{G}} \chi(g) = \begin{cases} |G| & \text{if } g = 1_G \\ 0 & \text{otherwise.} \end{cases}$$

Finally, one can also prove.

More orthogonality relations.

We can also compute a slightly different sum (where we fix the group element and vary the character).

Corollary (Summing over characters)

If $g \in G$ then

$$\sum_{\chi \in \widehat{G}} \chi(g) = \begin{cases} |G| & \text{if } g = 1_G \\ 0 & \text{otherwise.} \end{cases}$$

Finally, one can also prove.

Corollary

If $g \in G$ then

$$\sum_{g' \in G} \chi(g) \overline{\chi}(g') = \begin{cases} |G| & \text{if } g = g' \\ 0 & \text{otherwise.} \end{cases}$$

Character Matrix

What we just proved can be put together in a concise way. Let us enumerate the characters χ_1, \dots, χ_n .

Definition (Character Matrix)

Given a group G , we define its character matrix to be the matrix whose (s, t) entry is $\chi_s(t)$.

Character Matrix

What we just proved can be put together in a concise way. Let us enumerate the characters χ_1, \dots, χ_n .

Definition (Character Matrix)

Given a group G , we define its character matrix to be the matrix whose (s, t) entry is $\chi_s(t)$.

Example (Character Matrix of \mathbb{Z}_n)

Let $\zeta_n = e^{2\pi i/n}$. Then the character matrix of \mathbb{Z}_n is given by

$$\begin{bmatrix} 1 & 1 & \dots & \dots & 1 \\ 1 & \zeta_n & \zeta_n^2 & \dots & \zeta_n^{n-1} \\ 1 & \zeta_n^2 & \zeta_n^4 & \dots & \zeta_n^{2(n-1)} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 1 & \zeta_n^{(n-1)} & \zeta_n^{2(n-1)} & \dots & \zeta_n^{(n-1)(n-1)} \end{bmatrix}$$

Theorem

If G is a finite abelian group of order n , then matrix $\frac{1}{\sqrt{n}}X_G$ is a unitary matrix. That is, $X_G X_G^ = nI_n$, where I_n is the $n \times n$ identity matrix.*

Theorem

If G is a finite abelian group of order n , then matrix $\frac{1}{\sqrt{n}}X_G$ is a unitary matrix. That is, $X_G X_G^ = nI_n$, where I_n is the $n \times n$ identity matrix.*

One can also conclude the following result.

Theorem

If G is a finite abelian group of order n , then matrix $\frac{1}{\sqrt{n}}X_G$ is a unitary matrix. That is, $X_G X_G^ = nI_n$, where I_n is the $n \times n$ identity matrix.*

One can also conclude the following result.

Theorem

The characters form an orthogonal basis of V_G , and each character has norm \sqrt{n} .

Theorem

If G is a finite abelian group of order n , then matrix $\frac{1}{\sqrt{n}}X_G$ is a unitary matrix. That is, $X_G X_G^ = nI_n$, where I_n is the $n \times n$ identity matrix.*

One can also conclude the following result.

Theorem

The characters form an orthogonal basis of V_G , and each character has norm \sqrt{n} . That is, the normalized characters $\frac{1}{\sqrt{n}}\chi$ form an orthonormal basis of V_G .

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:**

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\},$

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\}$,
- **Normalized characters:**

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\},$
- **Normalized characters:** $B_G = \{B_g = \frac{1}{|G|}\chi_g : g \in G\}.$

Fourier Transform

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\}$,
- **Normalized characters:** $B_G = \{B_g = \frac{1}{|G|}\chi_g : g \in G\}$.

This allows us to define the *Fourier transform*.

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\},$
- **Normalized characters:** $B_G = \{B_g = \frac{1}{|G|}\chi_g : g \in G\}.$

This allows us to define the *Fourier transform*.

Definition

The unique linear operator on V_G sending each B_g to δ_g is known as the *Fourier transform*.

We have just seen two orthonormal bases for V_G :

- **Dirac deltas:** $\Delta_G = \{\delta_g : g \in G\},$
- **Normalized characters:** $B_G = \{B_g = \frac{1}{|G|}\chi_g : g \in G\}.$

This allows us to define the *Fourier transform*.

Definition

The unique linear operator on V_G sending each B_g to δ_g is known as the *Fourier transform*.

This operator is denoted as \mathcal{F} . We denote the Fourier transform (FT) of a function $f \in V_G$ as either $\mathcal{F}(f)$ or as \hat{f} .

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- 1 The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$. ^a

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- 1 *The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$.^a*
- 2 *\hat{f} is the unique complex-valued function in V_G whose value at g is the coefficient $\langle f, B_g \rangle$.*

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- 1 The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$.^a
- 2 \hat{f} is the unique complex-valued function in V_G whose value at g is the coefficient $\langle f, B_g \rangle$. That is: $\hat{f} = \sum_{g \in G} \langle f, B_g \rangle \delta_g$.

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- ① *The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$.^a*
- ② *\hat{f} is the unique complex-valued function in V_G whose value at g is the coefficient $\langle f, B_g \rangle$. That is: $\hat{f} = \sum_{g \in G} \langle f, B_g \rangle \delta_g$.*
- ③ *The FT of the FT is just the reversed signal: $\hat{\hat{f}}(x) = f(-x)$.*

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- ① *The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$.^a*
- ② *\hat{f} is the unique complex-valued function in V_G whose value at g is the coefficient $\langle f, B_g \rangle$. That is: $\hat{f} = \sum_{g \in G} \langle f, B_g \rangle \delta_g$.*
- ③ *The FT of the FT is just the reversed signal: $\hat{\hat{f}}(x) = f(-x)$.*
- ④ *The FT at the identity corresponds to the signal's average value:*

Properties of the Fourier Transform

We can prove some elegant properties of the FT.

Theorem (Properties of the Fourier Transform)

Let G be a finite abelian group.

- 1 The FT is an isometry, so $\langle f_1, f_2 \rangle = \langle \hat{f}_1, \hat{f}_2 \rangle$ for all $f_1, f_2 \in G$.^a
- 2 \hat{f} is the unique complex-valued function in V_G whose value at g is the coefficient $\langle f, B_g \rangle$. That is: $\hat{f} = \sum_{g \in G} \langle f, B_g \rangle \delta_g$.
- 3 The FT of the FT is just the reversed signal: $\hat{\hat{f}}(x) = f(-x)$.
- 4 The FT at the identity corresponds to the signal's average value: $\hat{f}(1_G) = \frac{1}{\sqrt{|G|}} \sum_{g \in G} f(g)$.
- 5 The FT on product of groups can be computed iteratively.

^athis is known as *Plancherel's identity*

Periods

Let us put this into context. Here is one natural way this construction arises. Consider a function $f : \mathbb{Z} \rightarrow \mathbb{C}$ which is periodic modulo n for some $n > 1$. This function descends uniquely to a function on \mathbb{Z}_n .

Lemma (Periods dividing Periods)

If a function on \mathbb{Z}_n is periodic, then its period divides n .

Theorem

If $f \in V_{\mathbb{Z}_n}$ is a nonconstant periodic function with period σ , and $f_\sigma \in V_{\mathbb{Z}_\sigma}$ is the corresponding restriction, then

Periods

Let us put this into context. Here is one natural way this construction arises. Consider a function $f : \mathbb{Z} \rightarrow \mathbb{C}$ which is periodic modulo n for some $n > 1$. This function descends uniquely to a function on \mathbb{Z}_n .

Lemma (Periods dividing Periods)

If a function on \mathbb{Z}_n is periodic, then its period divides n .

Theorem

If $f \in V_{\mathbb{Z}_n}$ is a nonconstant periodic function with period σ , and $f_\sigma \in V_{\mathbb{Z}_\sigma}$ is the corresponding restriction, then

$$\hat{f}(s) = \begin{cases} \sqrt{\frac{n}{\sigma}} \hat{f}_\sigma(m) & \text{if } s = m \frac{n}{\sigma} \text{ for some } 0 \leq m < \sigma, \\ 0 & \text{otherwise.} \end{cases}$$

Periods

Let us put this into context. Here is one natural way this construction arises. Consider a function $f : \mathbb{Z} \rightarrow \mathbb{C}$ which is periodic modulo n for some $n > 1$. This function descends uniquely to a function on \mathbb{Z}_n .

Lemma (Periods dividing Periods)

If a function on \mathbb{Z}_n is periodic, then its period divides n .

Theorem

If $f \in V_{\mathbb{Z}_n}$ is a nonconstant periodic function with period σ , and $f_\sigma \in V_{\mathbb{Z}_\sigma}$ is the corresponding restriction, then

$$\hat{f}(s) = \begin{cases} \sqrt{\frac{n}{\sigma}} \hat{f}_\sigma(m) & \text{if } s = m \frac{n}{\sigma} \text{ for some } 0 \leq m < \sigma, \\ 0 & \text{otherwise.} \end{cases}$$

It follows that \hat{f} has at most σ nonzero values.

① **Fourier Inversion:**

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$.

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.

- 1 **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- 2 **Convolution:**

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$.

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution,

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution, and a kind of orthogonality $B_s * B_t$ is nonzero exactly when $s = t$.

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution, and a kind of orthogonality $B_s * B_t$ is nonzero exactly when $s = t$. These elements form a ring with the identity being δ .

- 1 **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- 2 **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution, and a kind of orthogonality $B_s * B_t$ is nonzero exactly when $s = t$. These elements form a ring with the identity being δ .
Most important, the FT sends convolution to multiplication:

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution, and a kind of orthogonality $B_s * B_t$ is nonzero exactly when $s = t$. These elements form a ring with the identity being δ .
Most important, the FT sends convolution to multiplication:

$$\widehat{f_1 * f_2} = \hat{f}_1 \hat{f}_2$$

- ① **Fourier Inversion:** By the definition of the Fourier transform, one gets that $\check{f} = \sum_{g \in G} \langle f, \delta_g \rangle B_g$. It follows hence that $f = \sum_{g \in G} \langle \hat{f}, \delta_g \rangle B_g$.
- ② **Convolution:** $(f_1 * f_2)(x) := \frac{1}{\sqrt{|G|}} \sum_{g \in G} f_1(xg^{-1})f_2(g)$. The convolution is a symmetric bilinear transformation on V_G , satisfying associativity, commutativity, distribution, and a kind of orthogonality $B_s * B_t$ is nonzero exactly when $s = t$. These elements form a ring with the identity being δ .
Most important, the FT sends convolution to multiplication:

$$\widehat{f_1 * f_2} = \hat{f}_1 \hat{f}_2$$

- ③ **Iterated Fourier Transforms:** (See Luong's Section 4.6)