

Lecture 10: Orthogonality (Section 3.1)

Thang Huynh, UC San Diego

1/31/2018

Important geometric examples

► Example 1. $A = \begin{bmatrix} c & 0 \\ 0 & c \end{bmatrix}$

Important geometric examples

► **Example 1.** $A = \begin{bmatrix} c & 0 \\ 0 & c \end{bmatrix}$

stretches every vector in \mathbb{R}^2 by the same factor c .

Important geometric examples

► **Example 1.** $A = \begin{bmatrix} c & 0 \\ 0 & c \end{bmatrix}$

stretches every vector in \mathbb{R}^2 by the same factor c .

► **Example 2.** $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

Important geometric examples

► **Example 1.** $A = \begin{bmatrix} c & 0 \\ 0 & c \end{bmatrix}$

stretches every vector in \mathbb{R}^2 by the same factor c .

► **Example 2.** $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

maps $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \end{bmatrix}$

Important geometric examples

► **Example 1.** $A = \begin{bmatrix} c & 0 \\ 0 & c \end{bmatrix}$

stretches every vector in \mathbb{R}^2 by the same factor c .

► **Example 2.** $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

maps $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \end{bmatrix}$

reflects every vector in \mathbb{R}^2 through the line $y = x$.

Important geometric examples

► Example 3. $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

Important geometric examples

► **Example 3.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$
gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ 0 \end{bmatrix}$, i.e.

Important geometric examples

► **Example 3.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ 0 \end{bmatrix}$, i.e.

projects every vector in \mathbb{R}^2 onto the x -axis.

Important geometric examples

► **Example 3.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ 0 \end{bmatrix}$, i.e.

projects every vector in \mathbb{R}^2 onto the x -axis.

► **Example 4.** $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

Important geometric examples

► **Example 3.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ 0 \end{bmatrix}$, i.e.

projects every vector in \mathbb{R}^2 onto the x -axis.

► **Example 4.** $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} -y \\ x \end{bmatrix}$, i.e.

Important geometric examples

► **Example 3.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} x \\ 0 \end{bmatrix}$, i.e.

projects every vector in \mathbb{R}^2 onto the x -axis.

► **Example 4.** $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

gives the map $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} -y \\ x \end{bmatrix}$, i.e.

rotates every vector in \mathbb{R}^2 counter-clockwise by 90° .

Inner product and distances

► **Definition.** The **inner product** or **dot product** of \mathbf{v} and \mathbf{w} in \mathbb{R}^n is defined by

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n.$$

Inner product and distances

► **Definition.** The **inner product** or **dot product** of \mathbf{v} and \mathbf{w} in \mathbb{R}^n is defined by

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n.$$

► **Example.** Find the inner product of the following two vectors

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}.$$

Inner product and distances

► **Definition.** The **norm** or **length** of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + \dots + v_n^2}.$$

Inner product and distances

► **Definition.** The **norm** or **length** of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + \dots + v_n^2}.$$

► **Definition.** The **distance** between points \mathbf{v} and \mathbf{w} in \mathbb{R}^n is

$$\text{dist}(\mathbf{v}, \mathbf{w}) = \|\mathbf{v} - \mathbf{w}\|.$$

Inner product and distances

- **Definition.** The **norm** or **length** of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + \dots + v_n^2}.$$

- **Definition.** The **distance** between points \mathbf{v} and \mathbf{w} in \mathbb{R}^n is

$$\text{dist}(\mathbf{v}, \mathbf{w}) = \|\mathbf{v} - \mathbf{w}\|.$$

- **Example.** In \mathbb{R}^2 ,

$$\text{dist}\left(\begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix}\right) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Orthogonal vectors

► **Definition.** \mathbf{v} and \mathbf{w} in \mathbb{R}^n are **orthogonal** if $\mathbf{v} \cdot \mathbf{w} = 0$.

Orthogonal vectors

- **Definition.** \mathbf{v} and \mathbf{w} in \mathbb{R}^n are **orthogonal** if $\mathbf{v} \cdot \mathbf{w} = 0$.
- **Pythagoras.** \mathbf{v} and \mathbf{w} are orthogonal

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \mathbf{v} \cdot \mathbf{v} - 2\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{w}$$

$$\iff \mathbf{v} \cdot \mathbf{w} = 0.$$

Orthogonal vectors

► **Definition.** \mathbf{v} and \mathbf{w} in \mathbb{R}^n are **orthogonal** if $\mathbf{v} \cdot \mathbf{w} = 0$.

► **Pythagoras.** \mathbf{v} and \mathbf{w} are orthogonal

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \mathbf{v} \cdot \mathbf{v} - 2\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{w}$$

$$\iff \mathbf{v} \cdot \mathbf{w} = 0.$$

► **Example.** Are the following vectors orthogonal?

$$\bullet \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$$

$$\bullet \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Orthogonal vectors

► **Definition.** \mathbf{v} and \mathbf{w} in \mathbb{R}^n are **orthogonal** if $\mathbf{v} \cdot \mathbf{w} = 0$.

► **Pythagoras.** \mathbf{v} and \mathbf{w} are orthogonal

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \mathbf{v} \cdot \mathbf{v} - 2\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{w}$$

$$\iff \mathbf{v} \cdot \mathbf{w} = 0.$$

► **Example.** Are the following vectors orthogonal?

$$\bullet \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$$

$$\bullet \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Orthogonal vectors

Theorem. Suppose that $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal. Then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.

► Proof.

Orthogonal vectors

► **Example.** The vectors $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ are pairwise orthogonal and have length 1.

Orthogonal vectors

► **Example.** The vectors $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ are pairwise orthogonal and have length 1.

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

Orthogonal vectors

► **Example.** The vectors $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ are pairwise orthogonal and have length 1.

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

► **Solution.** $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}$ and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$.

Orthogonal vectors

► **Example.** The vectors $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ are pairwise orthogonal and have length 1.

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

► **Solution.** $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}$ and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$. Note that the two basis vectors are orthogonal $\begin{bmatrix} -2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \end{bmatrix} = 0$.

Orthogonal vectors

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

Orthogonal vectors

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

► **Solution.** $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$ and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

Orthogonal vectors

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

► **Solution.** $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$ and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

the vector in $N(A)$ is orthogonal to the vectors in $C(A^T)$

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Orthogonal vectors

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$. Find $N(A)$ and $C(A^T)$.

► **Solution.** $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$ and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

the vector in $N(A)$ is orthogonal to the vectors in $C(A^T)$

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Vectors in $N(A)$ are orthogonal to vectors in $C(A^T)$.

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W . (Is W^\perp a vector space?)

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W . (Is W^\perp a vector space?)

► **Example.** In previous example, $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$ and

$C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ are orthogonal subspaces.

The fundamental theorem, part II

Theorem. (Fundamental Theorem of Linear Algebra, Part II)

- $N(A)$ is orthogonal to $C(A^T)$. (The two spaces are orthogonal complements)
- $N(A^T)$ is orthogonal to $C(A)$. (The two spaces are orthogonal complements)

► Proof.

The fundamental theorem, part II

Theorem. (Fundamental Theorem of Linear Algebra, Part II)

- $N(A)$ is orthogonal to $C(A^T)$. (The two spaces are orthogonal complements)
- $N(A^T)$ is orthogonal to $C(A)$. (The two spaces are orthogonal complements)

► Proof.

Theorem. (Fundamental Theorem of Linear Algebra, Part I)

Let A be an $m \times n$ matrix of rank r .

- $\dim C(A) = \dim C(A^T) = r$.
- $\dim N(A) = n - r$
- $\dim N(A^T) = m - r$.

► **Example.** Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

► **Example.** Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

► **Example.** Let $V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}$. Find a basis for the orthogonal complement of V .

A new perspective on $Ax = b$

$Ax = b$ is solvable

$\iff b$ is in $C(A)$

$\iff b$ is orthogonal to $N(A^T)$

A new perspective on $Ax = b$

$Ax = b$ is solvable

$\Leftrightarrow b$ is in $C(A)$

$\Leftrightarrow b$ is orthogonal to $N(A^T)$

► **Example.** Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix}$. For which b does $Ax = b$ have a solution?