

# **Lecture 25: Jordan form; Singular Value Decompositions (Sections 5.6-6.3)**

---

Thang Huynh, UC San Diego

3/14/2018

## The Jordan form

Recall that if  $A$  is diagonalized,  $A$  is similar to a diagonal matrix  $\Lambda$  whose entries are eigenvalues of  $A$ .

## The Jordan form

Recall that if  $A$  is diagonalized,  $A$  is similar to a diagonal matrix  $\Lambda$  whose entries are eigenvalues of  $A$ .

How if  $A$  cannot be diagonalized?

## The Jordan form

If  $A$  has  $s$  independent eigenvectors, it is similar to a matrix with  $s$  blocks:

Jordan form

$$J = M^{-1}AM = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_s \end{bmatrix}.$$

► **Property.** Two matrices are similar if and only if they share the same Jordan form  $J$ .

## The Jordan form

If  $A$  has  $s$  independent eigenvectors, it is similar to a matrix with  $s$  blocks:

Jordan form  $J = M^{-1}AM = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_s \end{bmatrix}.$

Each  $J_i$  has only a single eigenvalue  $\lambda_i$  and one eigenvector:

Jordan block  $J_i = \begin{bmatrix} \lambda_i & 1 & & \\ & \lambda_i & \cdot & \\ & & \cdot & 1 \\ & & & \lambda_i \end{bmatrix}.$

► **Property.** Two matrices are similar if and only if they share the same Jordan form  $J$ .

## The Jordan form

► **Example.**  $T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ ,  $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$ , and  $B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  all lead to  $J = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ .

## The Jordan form

► **Example.**  $T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ ,  $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$ , and  $B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  all lead to  $J = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ .

These four matrices have eigenvalues 1 and 1 with only one eigenvector. How to find a matrix  $M$  such that  $M^{-1}TM = J$ ?

## The Jordan form

► **Example.**  $T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ ,  $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$ , and  $B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  all lead to  $J = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ .

These four matrices have eigenvalues 1 and 1 with only one eigenvector. How to find a matrix  $M$  such that  $M^{-1}TM = J$ ?

► **Definition.** If  $A$  is an  $n \times n$  matrix, a **generalized eigenvector** of  $A$  corresponding to the eigenvalue  $\lambda$  is a nonzero vector  $\mathbf{x}$  satisfying

$$(A - \lambda I)^p \mathbf{x} = 0$$

for some positive integer  $p$ .



## The Jordan form

$T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$  has an eigenvalue  $\lambda = 1$  of multiplicity 2. We want to find generalized eigenvectors from the equations:

## The Jordan form

$T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$  has an eigenvalue  $\lambda = 1$  of multiplicity 2. We want to find generalized eigenvectors from the equations:

- $(A - I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ , and
- $(A - I)^2\mathbf{x} = 0 \Rightarrow \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

## The Jordan form

$T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$  has an eigenvalue  $\lambda = 1$  of multiplicity 2. We want to find generalized eigenvectors from the equations:

- $(A - I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ , and
- $(A - I)^2\mathbf{x} = 0 \Rightarrow \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

$$M = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

## The Jordan form

$T = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$  has an eigenvalue  $\lambda = 1$  of multiplicity 2. We want to find generalized eigenvectors from the equations:

- $(A - I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ , and
- $(A - I)^2\mathbf{x} = 0 \Rightarrow \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

$$M = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$M^{-1}TM = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = J.$$

## The Jordan form

► **Example.** For  $A = \begin{bmatrix} 0 & 1 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ . Which of the following matrices are the Jordan forms of  $A$  and  $B$ ?

$$J_1 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \quad J_2 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad J_3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

## Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$
- columns of  $V$  ( $n$  by  $n$ ) are eigenvectors of  $A^TA$



## Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$
- columns of  $V$  ( $n$  by  $n$ ) are eigenvectors of  $A^TA$
- The  $r$  singular values on the diagonal of  $\Sigma$  ( $m$  by  $n$ ) are the square roots of the nonzero eigenvalues of both  $AA^T$  and  $A^TA$ .

# Singular Value Decompositions

► **Example.** Consider  $A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$ .

## Singular Value Decompositions

► **Example.** Consider  $A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$ . Then  $A$  has rank 2,  
and  $AA^T = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$  with eigenvalues  $\lambda = 3$  and 1.

## Singular Value Decompositions

► **Example.** Consider  $A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$ . Then  $A$  has rank 2,

and  $AA^T = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$  with eigenvalues  $\lambda = 3$  and 1.

$$A = U\Sigma V^T = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{6} & -2/\sqrt{6} & 1/\sqrt{6} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \end{bmatrix}.$$

## Least squares and the Pseudoinverse

A least squares solution  $\hat{\mathbf{x}}$  of the linear system

$$A\mathbf{x} = \mathbf{b}$$

is the one minimizing  $\|A\mathbf{x} - \mathbf{b}\|$ . To solve this, we can solve the associate normal system

$$A^T A \hat{\mathbf{x}} = A^T \mathbf{b}.$$

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**?

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ .



## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ . The optimal solution of the  $A\mathbf{x} = \mathbf{b}$  is the vector  $\hat{\mathbf{x}}$  that has the minimum length. We denote this solution by  $\mathbf{x}^\dagger$ , and the matrix which produces  $\mathbf{x}^\dagger$  from  $\mathbf{b}$  is called the **pseudoinverse** of  $A$  and is denoted by  $A^\dagger$ .

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ . The optimal solution of the  $A\mathbf{x} = \mathbf{b}$  is the vector  $\hat{\mathbf{x}}$  that has the minimum length. We denote this solution by  $\mathbf{x}^\dagger$ , and the matrix which produces  $\mathbf{x}^\dagger$  from  $\mathbf{b}$  is called the **pseudoinverse** of  $A$  and is denoted by  $A^\dagger$ .

$$\mathbf{x}^\dagger = A^\dagger \mathbf{b}.$$

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ?

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$ .

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$ .

The SVD of  $A$  is

$$A = [1][3 \ 0 \ 0] \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}.$$

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$ .

The SVD of  $A$  is

$$A = [1][3 \ 0 \ 0] \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}.$$

Then the pseudoinverse of  $A$  is

$$A^\dagger = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} \frac{1}{3} \\ 0 \\ 0 \end{bmatrix} [1] = \begin{bmatrix} -\frac{1}{9} \\ \frac{2}{9} \\ \frac{2}{9} \end{bmatrix}.$$