

Lecture 5: Vector spaces and subspaces

Thang Huynh, UC San Diego

1/19/2018

Vector spaces

► **Definition.** A **vector space** is a nonempty set V of elements, called **vectors**, which may be added and scaled.

Vector spaces

► **Definition.** A **vector space** is a nonempty set V of elements, called **vectors**, which may be added and scaled.

In other words, a vector space is a collection of vectors which can be added and scaled; subject to the usual rules you would hope for, namely associativity, commutativity, distributivity.

Vector spaces

- **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

Vector spaces

- ▶ **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)
- ▶ **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

Vector spaces

► **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

► **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

What is the zeros vector in $M_{2 \times 2}$?

Vector spaces

► **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

► **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

What is the zeros vector in $M_{2 \times 2}$?

► **Example.** Let \mathcal{P}_n be the set of all polynomials of degree at most $n \geq 0$. Is \mathcal{P}_n a vector space?

Vector spaces

► **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

► **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

What is the zeros vector in $M_{2 \times 2}$?

► **Example.** Let \mathcal{P}_n be the set of all polynomials of degree at most $n \geq 0$. Is \mathcal{P}_n a vector space?

Yes!

Vector spaces

► **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

► **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

What is the zeros vector in $M_{2 \times 2}$?

► **Example.** Let \mathcal{P}_n be the set of all polynomials of degree at most $n \geq 0$. Is \mathcal{P}_n a vector space?

Yes!

► **Example.** Let V be the set of all polynomials of degree exactly 3. Is V a vector space?

Vector spaces

► **Example.** The Euclidean space \mathbb{R}^n is a vector space. (Why?)

► **Example.** The space $M_{2 \times 2} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}$ is a vector space.

What is the zeros vector in $M_{2 \times 2}$?

► **Example.** Let \mathcal{P}_n be the set of all polynomials of degree at most $n \geq 0$. Is \mathcal{P}_n a vector space?

Yes!

► **Example.** Let V be the set of all polynomials of degree exactly 3. Is V a vector space?

No! For example, take $f(x) = x^3 + x$ and $g(x) = -x^3 + 5x^2 + 2$.

Both f and g are in V , but their sum $f(x) + g(x) = 5x^2 + x + 2$ is not in V .

Subspaces

► **Definition.** A subset W of a vector space V is a **subspace** of V if W itself is a vector space.

Subspaces

► **Definition.** A subset W of a vector space V is a **subspace** of V if W itself is a vector space.

How to check if a subset W is a subspace?

Subspaces

► **Definition.** A subset W of a vector space V is a **subspace** of V if W itself is a vector space.

How to check if a subset W is a subspace?

- W contains the zero vector 0 .
- W is closed under addition, i.e. if $\mathbf{u}, \mathbf{v} \in W$, then $\mathbf{u} + \mathbf{v} \in W$.
- W is closed under scaling, i.e. if $\mathbf{u} \in W$ and $c \in \mathbb{R}$, then $c\mathbf{u} \in W$.

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Yes!

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} x \\ x + 1 \end{bmatrix} : x \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^2 ?

Subspaces

► **Example.** Is $W = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^3 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$ a subspace of \mathbb{R}^2 ?

Yes!

► **Example.** Is $W = \left\{ \begin{bmatrix} x \\ x + 1 \end{bmatrix} : x \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^2 ?

No!

Spans of vectors are subspaces

The **span** of vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ is the set of all their linear combinations. We denote it by $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$.

Spans of vectors are subspaces

The **span** of vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ is the set of all their linear combinations. We denote it by $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$.

Theorem. If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are in a vector space V , then $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ is a subspace.

Spans of vectors are subspaces

► **Example.** Is $W = \left\{ \begin{bmatrix} a + 3b \\ 2a - b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^2 ?

Spans of vectors are subspaces

- **Example.** Is $W = \left\{ \begin{bmatrix} a + 3b \\ 2a - b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^2 ?
- **Example.** Is $W = \left\{ \begin{bmatrix} -a & 2b \\ a + b & 3a \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of $\mathcal{M}_{2 \times 2}$, the vector space of 2×2 matrices?

Spans of vectors are subspaces

- **Example.** Is $W = \left\{ \begin{bmatrix} a + 3b \\ 2a - b \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of \mathbb{R}^2 ?
- **Example.** Is $W = \left\{ \begin{bmatrix} -a & 2b \\ a + b & 3a \end{bmatrix} : a, b \in \mathbb{R} \right\}$ a subspace of $\mathcal{M}_{2 \times 2}$, the vector space of 2×2 matrices?

Spans of vectors are subspaces

► **Exercise.** Are the following vector spaces?

$$\bullet W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + 3b = 0, 2a - c = 1 \right\}.$$

$$\bullet W_2 = \left\{ \begin{bmatrix} a + c \\ -2b \\ b + 3c \\ c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}.$$

$$\bullet W_1 = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} : ab \geq 0 \right\}.$$

Null spaces

► **Definition.** The **null space** of a matrix A is

$$N(A) = \{\mathbf{v} : A\mathbf{v} = \mathbf{0}\}.$$

In other words, if A is $m \times n$, then its null space consists of those vectors $\mathbf{v} \in \mathbb{R}^n$ which solve the *homogeneous* equation $A\mathbf{x} = \mathbf{0}$.

Null spaces

► **Definition.** The **null space** of a matrix A is

$$N(A) = \{\mathbf{v} : A\mathbf{v} = \mathbf{0}\}.$$

In other words, if A is $m \times n$, then its null space consists of those vectors $\mathbf{v} \in \mathbb{R}^n$ which solve the *homogeneous* equation $A\mathbf{x} = \mathbf{0}$.

Theorem. If A is $m \times n$, then $N(A)$ is subspace of \mathbb{R}^n .

Null spaces

► **Definition.** The **null space** of a matrix A is

$$N(A) = \{\mathbf{v} : A\mathbf{v} = \mathbf{0}\}.$$

In other words, if A is $m \times n$, then its null space consists of those vectors $\mathbf{v} \in \mathbb{R}^n$ which solve the *homogeneous* equation $A\mathbf{x} = \mathbf{0}$.

Theorem. If A is $m \times n$, then $N(A)$ is subspace of \mathbb{R}^n .

► **Proof.**

Null spaces

► **Example.** Find an explicit description of $N(A)$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

Null spaces

► **Example.** Find an explicit description of $N(A)$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

► **Solution.** This is equivalent to finding the set of all solutions of $A\mathbf{x} = \mathbf{0}$.

Null spaces

► **Example.** Find an explicit description of $N(A)$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

► **Solution.** This is equivalent to finding the set of all solutions of $Ax = \mathbf{0}$.

$$\left[\begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccccc|c} \boxed{1} & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & \boxed{1} & -6 & -15 & 0 \end{array} \right].$$

Null spaces

► **Example.** Find an explicit description of $N(A)$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

► **Solution.** This is equivalent to finding the set of all solutions of $Ax = \mathbf{0}$.

$$\left[\begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccccc|c} \boxed{1} & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & \boxed{1} & -6 & -15 & 0 \end{array} \right].$$

Hence, x_1 and x_3 are pivot variables. Let $x_2 = u$, $x_4 = v$, and $x_5 = w$.

Null spaces

► **Example.** Find an explicit description of $N(A)$ where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

► **Solution.** This is equivalent to finding the set of all solutions of $Ax = \mathbf{0}$.

$$\left[\begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccccc|c} \boxed{1} & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & \boxed{1} & -6 & -15 & 0 \end{array} \right].$$

Hence, x_1 and x_3 are pivot variables. Let $x_2 = u$, $x_4 = v$, and $x_5 = w$.

The solutions of $A\mathbf{x} = \mathbf{0}$ is of the form

$$\begin{bmatrix} -2u - 13v - 33w \\ u \\ 6v + 15w \\ v \\ w \end{bmatrix} = u \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + v \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + w \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}.$$

The solutions of $A\mathbf{x} = \mathbf{0}$ is of the form

$$\begin{bmatrix} -2u - 13v - 33w \\ u \\ 6v + 15w \\ v \\ w \end{bmatrix} = u \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + v \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + w \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}.$$

$$N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

Column spaces

► **Definition.** The **column space**, $C(A)$, of A is the span of the vector columns of A , i.e., if $A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_n]$, then $C(A) = \text{span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$.

Column spaces

► **Definition.** The **column space**, $C(A)$, of A is the span of the vector columns of A , i.e., if $A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_n]$, then $C(A) = \text{span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$. Facts:

- \mathbf{b} is in $C(A)$ if and only if $A\mathbf{x} = \mathbf{b}$ has a solution. (Why?)

Column spaces

► **Definition.** The **column space**, $C(A)$, of A is the span of the vector columns of A , i.e., if $A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_n]$, then $C(A) = \text{span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$. Facts:

- \mathbf{b} is in $C(A)$ if and only if $A\mathbf{x} = \mathbf{b}$ has a solution. (Why?)
- If A is $m \times n$, then $C(A)$ is a subspace of \mathbb{R}^m . (Why?).

Column spaces

► **Example.** Find a matrix A such that $W = C(A)$ where

$$W = \left\{ \begin{bmatrix} 2x - y \\ 3y \\ 7x + y \end{bmatrix} : x, y \in \mathbb{R} \right\}.$$

$C(A)$ and solutions to $A\mathbf{x} = \mathbf{b}$

Theorem. Let \mathbf{x}_p be a solution of the equation $A\mathbf{x} = \mathbf{b}$. Then every solution to $A\mathbf{x} = \mathbf{b}$ is of the form $\mathbf{x} = \mathbf{x}_p + \mathbf{x}_n$, where \mathbf{x}_n is a solution to the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

$C(A)$ and solutions to $Ax = b$

► **Example.** Find a parametric description of the solutions to

$$\begin{bmatrix} 1 & 3 & 3 & 2 \\ 2 & 6 & 9 & 7 \\ -1 & -3 & 3 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \\ 5 \end{bmatrix}.$$