

Lecture 23: Similarity Transformations (Sections 5.6)

Thang Huynh, UC San Diego

3/9/2018

Unitary matrices

► **Example.** $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$ has eigenvalues e^{it} and e^{-it} .

Unitary matrices

► **Example.** $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$ has eigenvalues e^{it} and e^{-it} .

Orthogonal eigenvectors are $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$.

Unitary matrices

► **Example.** $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$ has eigenvalues e^{it} and e^{-it} .

Orthogonal eigenvectors are $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$.

► **Example.** (normalized) Fourier matrix

$$U = \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & w & \dots & w^{n-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & w^{n-1} & \dots & w^{(n-1)^2} \end{bmatrix}$$

where $w = e^{2\pi i/n}$.

Unitary matrices

► **Example.** $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$ has eigenvalues e^{it} and e^{-it} .

Orthogonal eigenvectors are $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$.

► **Example.** (normalized) Fourier matrix

$$U = \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & w & \dots & w^{n-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & w^{n-1} & \dots & w^{(n-1)^2} \end{bmatrix}$$

where $w = e^{2\pi i/n}$.

Row i of U^H times column j of U is

$$\frac{1}{n}(1 + W + W^2 + \dots + W^{n-1}) = \frac{W^n - 1}{W - 1} = 0,$$

where $W = w^{j-i}$.

Similar transformations

► **Definition.** Two $n \times n$ matrices A and B are called **similar** if $B = M^{-1}AM$ for some invertible $n \times n$ matrix M .

Similar transformations

- ▶ **Definition.** Two $n \times n$ matrices A and B are called **similar** if $B = M^{-1}AM$ for some invertible $n \times n$ matrix M .
- ▶ **Example.** If A can be diagonalized, Λ and A are similar.

Similar transformations

► **Property.** If A and B are similar, they have the **same eigenvalues**. Every eigenvector x of A corresponds to an eigenvector $M^{-1}x$ of B .

Similar transformations

► **Property.** If A and B are similar, they have the **same eigenvalues**. Every eigenvector \mathbf{x} of A corresponds to an eigenvector $M^{-1}\mathbf{x}$ of B .

$A - \lambda I$ and $B - \lambda I$ have the same determinant.

Similar transformations

► **Property.** If A and B are similar, they have the **same eigenvalues**. Every eigenvector \mathbf{x} of A corresponds to an eigenvector $M^{-1}\mathbf{x}$ of B .

$A - \lambda I$ and $B - \lambda I$ have the same determinant.

Similar transformations

► **Example.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ has eigenvalues 1. Each B is $M^{-1}AM$:

Similar transformations

► **Example.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ has eigenvalues 1. Each B is $M^{-1}AM$:

If $M = \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}$, then $B = \begin{bmatrix} 1 & b \\ 0 & 0 \end{bmatrix}$: triangular with $\lambda = 1$ and 0.

Similar transformations

► **Example.** $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ has eigenvalues 1. Each B is $M^{-1}AM$:

If $M = \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}$, then $B = \begin{bmatrix} 1 & b \\ 0 & 0 \end{bmatrix}$: triangular with $\lambda = 1$ and 0.

If $M = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$, then $B = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}$: triangular with $\lambda = 1$ and 0.

Triangular forms with a unitary M

Theorem. (Schur's Theorem) There is a unitary matrix U such that $U^{-1}AU = T$ is triangular. The eigenvalues of A appear along the diagonal of this similar matrix T .

Triangular forms with a unitary M

Theorem. (Schur's Theorem) There is a unitary matrix U such that $U^{-1}AU = T$ is triangular. The eigenvalues of A appear along the diagonal of this similar matrix T .

► **Example.** $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$ has the eigenvalue $\lambda = 1$ (twice).

$$U^{-1}AU = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = T.$$

Diagonalize symmetric and Hermitian matrices

Theorem. (Spectral Theorem) Every real symmetric A can be diagonalized by an orthogonal matrix Q . Every Hermitian matrix can be diagonalized by a unitary U :

$$\text{(real)} \quad Q^{-1}AQ = \Lambda \text{ or } A = Q\Lambda Q^T$$

$$\text{(complex)} \quad U^{-1}AU = \Lambda \text{ or } A = U\Lambda U^H$$

Diagonalize symmetric and Hermitian matrices

► **Example.** Consider $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

Diagonalize symmetric and Hermitian matrices

► **Example.** Consider $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Then A is symmetric with repeated eigenvalues $\lambda_1 = \lambda_2 = 1$ and $\lambda_3 = -1$.

Diagonalize symmetric and Hermitian matrices

► **Example.** Consider $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Then A is symmetric with repeated eigenvalues $\lambda_1 = \lambda_2 = 1$ and $\lambda_3 = -1$.

- For $\lambda_1 = \lambda_2 = 1$, eigenvectors $\mathbf{x}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and $\mathbf{x}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$
- For $\lambda_3 = -1$, eigenvector $\mathbf{x}_3 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$.

These are the columns of Q .

Normal matrices

► **Definition.** The matrix N is **normal** if it commutes with its conjugate transpose N^H : $NN^H = N^HN$.

Normal matrices

► **Definition.** The matrix N is **normal** if it commutes with its conjugate transpose N^H : $NN^H = N^HN$.
For such matrices, the triangular $T = U^{-1}NU$ is the diagonal Λ .
Normal matrices are exactly those that have a **complete set of orthonormal eigenvectors**.

Normal matrices

► **Definition.** The matrix N is **normal** if it commutes with its conjugate transpose N^H : $NN^H = N^HN$.

For such matrices, the triangular $T = U^{-1}NU$ is the diagonal Λ . Normal matrices are exactly those that have a **complete set of orthonormal eigenvectors**.

► **Example.**

- Symmetric and Hermitian matrices are normal.

Normal matrices

► **Definition.** The matrix N is **normal** if it commutes with its conjugate transpose N^H : $NN^H = N^HN$.

For such matrices, the triangular $T = U^{-1}NU$ is the diagonal Λ . Normal matrices are exactly those that have a **complete set of orthonormal eigenvectors**.

► **Example.**

- Symmetric and Hermitian matrices are normal.
- Orthogonal and unitary matrices are also normal.

Normal matrices

► **Property.** If N is normal, then so is the triangular $T = U^{-1}NU$.

Normal matrices

- ▶ **Property.** If N is normal, then so is the triangular $T = U^{-1}NU$.
- ▶ **Property.** A triangular T that is normal must be diagonal.
(Why?)