

Lecture 21: Linear Differential Equations; Complex Matrices (Sections 5.4--5.5)

Thang Huynh, UC San Diego

3/5/2018

Linear differential equations

- **Goal.** Solve the initial value problem

$$\mathbf{y}' = A\mathbf{y}, \quad \mathbf{y}(0) = \mathbf{y}_0.$$

Linear differential equations

► **Definition.** Let A be $n \times n$. The **matrix exponential** is

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$$

Linear differential equations

► **Definition.** Let A be $n \times n$. The **matrix exponential** is

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$$

Then

- $\frac{d}{dt}e^{At} = Ae^{At}.$

Linear differential equations

► **Definition.** Let A be $n \times n$. The **matrix exponential** is

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$$

Then

- $\frac{d}{dt}e^{At} = Ae^{At}.$
- $(e^{As})(e^{At}) = e^{A(s+t)}$
- $(e^{At})(e^{-At}) = I$

Linear differential equations

► **Definition.** Let A be $n \times n$. The **matrix exponential** is

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$$

Then

- $\frac{d}{dt}e^{At} = Ae^{At}$.
- $(e^{As})(e^{At}) = e^{A(s+t)}$
- $(e^{At})(e^{-At}) = I$
- Note that $e^{A+B} = e^A e^B$ is **not true** for matrices. For example,

$$A = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix}.$$

Linear differential equations

► **Definition.** Let A be $n \times n$. The **matrix exponential** is

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$$

Then

- $\frac{d}{dt}e^{At} = Ae^{At}$.
- $(e^{As})(e^{At}) = e^{A(s+t)}$
- $(e^{At})(e^{-At}) = I$
- Note that $e^{A+B} = e^A e^B$ is **not true** for matrices. For example,

$$A = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix}.$$

Theorem. Suppose $A = S\Lambda S^{-1}$. Then $e^A = Se^{\Lambda}S^{-1}$.

Linear differential equations

► **Example.** If $A = \begin{bmatrix} 2 & 0 \\ 0 & 5 \end{bmatrix}$, then

$$e^A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 0 \\ 0 & 5 \end{bmatrix} + \frac{1}{2!} \begin{bmatrix} 2^2 & 0 \\ 0 & 5^2 \end{bmatrix} + \dots = \begin{bmatrix} e^2 & 0 \\ 0 & e^5 \end{bmatrix}$$

$$e^{At} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2t & 0 \\ 0 & 5t \end{bmatrix} + \frac{1}{2!} \begin{bmatrix} (2t)^2 & 0 \\ 0 & (5t)^2 \end{bmatrix} + \dots = \begin{bmatrix} e^{2t} & 0 \\ 0 & e^{5t} \end{bmatrix}$$

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}$, $\mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}$, $\mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}, \mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

► **Solution.** The solution is $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}, \mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

► **Solution.** The solution is $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

Step 1: Diagonalize A

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}, \mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

► **Solution.** The solution is $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

Step 1: Diagonalize A

- $\lambda_1 = 1 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$
- $\lambda_2 = -1 \Rightarrow \mathbf{x}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Linear differential equations

The solution to $\mathbf{y}' = A\mathbf{y}, \mathbf{y}(0) = \mathbf{y}_0$ is

$$\mathbf{y}(t) = e^{At}\mathbf{y}_0.$$

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

► **Solution.** The solution is $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

Step 1: Diagonalize A

- $\lambda_1 = 1 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$
- $\lambda_2 = -1 \Rightarrow \mathbf{x}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Linear differential equations

Hence,

$$A = S\Lambda S^{-1}, \quad \text{where } S = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \text{ and } \Lambda = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Linear differential equations

Hence,

$$A = S\Lambda S^{-1}, \quad \text{where } S = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \text{ and } \Lambda = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Step 2: Compute the solution $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

$$\mathbf{y} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} e^t & 0 \\ 0 & e^{-t} \end{bmatrix} \frac{1}{2} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} e^t + e^{-t} \\ e^t - e^{-t} \end{bmatrix}.$$

Linear differential equations

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}.$$

Linear differential equations

- **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}.$$

- **Solution.** Step 1: Diagonalize A

Linear differential equations

► **Example.** Solve the differential equation

$$\mathbf{y}' = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}.$$

► **Solution.** Step 1: Diagonalize A

A has eigenvalues 2 and 4 (Why?).

- $\lambda = 2 \Rightarrow$ eigenspace $\text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$
- $\lambda = 4 \Rightarrow$ eigenspace $\text{span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}$

Linear differential equations

$$A = S\Lambda S^{-1}, \quad \text{where } S = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}, \Lambda = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix}.$$

Linear differential equations

$$A = S\Lambda S^{-1}, \quad \text{where } S = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}, \Lambda = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix}.$$

The solution is

$$\Rightarrow \mathbf{y} = e^{At}\mathbf{y}_0 = \begin{bmatrix} e^{2t} \\ e^{2t} + e^{4t} \\ e^{4t} \end{bmatrix}.$$

Complex Numbers and Their Conjugates

- **Complex conjugate** of $a + ib$ is $a - ib$ and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

Complex Numbers and Their Conjugates

- **Complex conjugate** of $a + ib$ is $a - ib$ and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

- Conjugate of a product (or sum) equals the product (or sum) of the conjugates:

$$\overline{(a + ib)(c + id)} = (\overline{a + ib})(\overline{c + id})$$

$$\overline{a + ib + c + id} = \overline{(a + c) + i(b + d)}$$

Complex Numbers and Their Conjugates

- **Complex conjugate** of $a + ib$ is $a - ib$ and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

- Conjugate of a product (or sum) equals the product (or sum) of the conjugates:

$$\overline{(a + ib)(c + id)} = (\overline{a + ib})(\overline{c + id})$$

$$\overline{a + ib} + \overline{c + id} = \overline{(a + c) + i(b + d)}$$

- Multiplying $a + ib$ by its conjugate produces $a^2 + b^2$:

$$(a + ib)(a - ib) = a^2 + b^2.$$

Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

► Length/norm of \mathbf{x}

$$\|\mathbf{x}\| = \sqrt{|x_1|^2 + |x_2|^2 + \dots + |x_n|^2}.$$

Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

► Length/norm of \mathbf{x}

$$\|\mathbf{x}\| = \sqrt{|x_1|^2 + |x_2|^2 + \dots + |x_n|^2}.$$

► Example.

$$\mathbf{x} = \begin{bmatrix} 1 \\ i \end{bmatrix} \Rightarrow \|\mathbf{x}\| = \sqrt{|1|^2 + |i|^2} = 2.$$

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products $\langle \mathbf{x}, \mathbf{y} \rangle$ and $\langle \mathbf{y}, \mathbf{x} \rangle$ of $\mathbf{x} = (1 + 3i, 3i)$ and $\mathbf{y} = (6 + 3i, 4 + i)$.

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products $\langle \mathbf{x}, \mathbf{y} \rangle$ and $\langle \mathbf{y}, \mathbf{x} \rangle$ of $\mathbf{x} = (1 + 3i, 3i)$ and $\mathbf{y} = (6 + 3i, 4 + i)$.

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products $\langle \mathbf{x}, \mathbf{y} \rangle$ and $\langle \mathbf{y}, \mathbf{x} \rangle$ of $\mathbf{x} = (1 + 3i, 3i)$ and $\mathbf{y} = (6 + 3i, 4 + i)$.

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

$$\langle \mathbf{y}, \mathbf{x} \rangle = (6 - 3i)(1 + 3i) + (4 - i)(3i) = 18 + 27i.$$

Lengths and Inner Products in the Complex Case

► Inner product of \mathbf{x} and \mathbf{y}

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products $\langle \mathbf{x}, \mathbf{y} \rangle$ and $\langle \mathbf{y}, \mathbf{x} \rangle$ of $\mathbf{x} = (1 + 3i, 3i)$ and $\mathbf{y} = (6 + 3i, 4 + i)$.

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

$$\langle \mathbf{y}, \mathbf{x} \rangle = (6 - 3i)(1 + 3i) + (4 - i)(3i) = 18 + 27i.$$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \mathbf{x} \rangle}$$

Conjugate transpose of a matrix A

► **Definition.** A conjugate transpose of a matrix A is \overline{A}^T (and denoted by A^H or A^*).

Conjugate transpose of a matrix A

► **Definition.** A conjugate transpose of a matrix A is \overline{A}^T (and denoted by A^H or A^*).

► **Example.**

$$\begin{bmatrix} 2+i & 3i \\ 4-i & 5 \\ 0 & 0 \end{bmatrix}^H = \begin{bmatrix} 2-i & 4+i & 0 \\ -3i & 5 & 0 \end{bmatrix}.$$