

# Lecture 17: Cofactor Expansion (Section 4.3)

---

Thang Huynh, UC San Diego

2/21/2018

# Review

Three most important properties of the determinants

►  $\det(I) = 1$ .

## Review

Three most important properties of the determinants

►  $\det(I) = 1$ .

► The determinant changes sign when two rows are exchanged.

$$\begin{vmatrix} c & d \\ a & b \end{vmatrix} = cb - ad = - \begin{vmatrix} a & b \\ c & d \end{vmatrix}.$$

► The determinant depends linearly on the first row.

## Review

Three most important properties of the determinants

►  $\det(I) = 1$ .

► The determinant changes sign when two rows are exchanged.

$$\begin{vmatrix} c & d \\ a & b \end{vmatrix} = cb - ad = - \begin{vmatrix} a & b \\ c & d \end{vmatrix}.$$

► The determinant depends linearly on the first row.

- Add vectors in row 1:  $\begin{vmatrix} a + a' & b + b' \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} a' & b' \\ c & d \end{vmatrix}$

## Review

Three most important properties of the determinants

►  $\det(I) = 1$ .

► The determinant changes sign when two rows are exchanged.

$$\begin{vmatrix} c & d \\ a & b \end{vmatrix} = cb - ad = - \begin{vmatrix} a & b \\ c & d \end{vmatrix}.$$

► The determinant depends linearly on the first row.

- Add vectors in row 1:  $\begin{vmatrix} a + a' & b + b' \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} a' & b' \\ c & d \end{vmatrix}$

- Multiply by  $t$  in row 1:  $\begin{vmatrix} ta & tb \\ c & d \end{vmatrix} = t \begin{vmatrix} a & b \\ c & d \end{vmatrix}$

## Cofactor expansion

► **Definition.** Let  $A = [a_{ij}]$  be an  $n \times n$  matrix. Denote by  $M_{ij}$  the submatrix of  $A$  obtained by deleting its row and column containing  $a_{ij}$ . Then  $\det(M_{ij})$  is called the **minor** of  $a_{ij}$ .

## Cofactor expansion

► **Definition.** Let  $A = [a_{ij}]$  be an  $n \times n$  matrix. Denote by  $M_{ij}$  the submatrix of  $A$  obtained by deleting its row and column containing  $a_{ij}$ . Then  $\det(M_{ij})$  is called the **minor** of  $a_{ij}$ .

► **Example.** Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ . Then

$$M_{11} = \begin{bmatrix} 5 & 6 \\ 8 & 9 \end{bmatrix} \text{ and } M_{23} = \begin{bmatrix} 1 & 2 \\ 7 & 8 \end{bmatrix}.$$

## Cofactor expansion

► **Definition.** Let  $A = [a_{ij}]$  be an  $n \times n$  matrix. Denote by  $M_{ij}$  the submatrix of  $A$  obtained by deleting its row and column containing  $a_{ij}$ . Then  $\det(M_{ij})$  is called the **minor** of  $a_{ij}$ .

► **Example.** Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ . Then

$$M_{11} = \begin{bmatrix} 5 & 6 \\ 8 & 9 \end{bmatrix} \text{ and } M_{23} = \begin{bmatrix} 1 & 2 \\ 7 & 8 \end{bmatrix}.$$

The minor of  $a_{11}$  is  $\det(M_{11}) = 5(9) - 8(6) = -3$  and the minor of  $a_{23}$  is  $\det(M_{23}) = -6$ .



## Cofactor expansion

► **Definition.** If we multiply the minor of  $a_{ij}$  by  $(-1)^{i+j}$ , then we arrive at the definition of the **cofactor**  $A_{ij}$  of  $a_{ij}$ :

$$A_{ij} = (-1)^{i+j} \det(M_{ij}).$$

## Cofactor expansion

► **Definition.** If we multiply the minor of  $a_{ij}$  by  $(-1)^{i+j}$ , then we arrive at the definition of the **cofactor**  $A_{ij}$  of  $a_{ij}$ :

$$A_{ij} = (-1)^{i+j} \det(M_{ij}).$$

► **Example.** From the previous example,  $A_{11} = (-1)^2(-3) = -3$  and  $A_{23} = (-1)^5(-6) = 6$ .

## Cofactor expansion

The method of cofactor expansion is given by the formulas

$$\det(A) = a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}$$

(expansion of  $\det(A)$  along  $i$ th row)

## Cofactor expansion

The method of cofactor expansion is given by the formulas

$$\det(A) = a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}$$

(expansion of  $\det(A)$  along  $i$ th row)

$$\det(A) = a_{1j}A_{1j} + a_{2j}A_{2j} + \dots + a_{nj}A_{nj}$$

(expansion of  $\det(A)$  along  $j$ th column).

## Cofactor expansion

► **Example.** For  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ .

## Cofactor expansion

► **Example.** For  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ .

$$\det(A) = a_{12}A_{12} + a_{22}A_{22} + a_{32}A_{32} = 1(-3) + 2(6) + 3(-3) = 0.$$

## Cofactor expansion

The method of cofactor expansion is especially applicable if a matrix has a row or a column with many zeros.

## Cofactor expansion

The method of cofactor expansion is especially applicable if a matrix has a row or a column with many zeros.

► **Example.** Compute the determinant of

$$A = \begin{bmatrix} 2 & -1 & 1 & 0 \\ 3 & 5 & 0 & -2 \\ 1 & 1 & 0 & -3 \\ 4 & 0 & 3 & -1 \end{bmatrix}.$$



## Cofactor expansion

The method of cofactor expansion is especially applicable if a matrix has a row or a column with many zeros.

► **Example.** Compute the determinant of

$$A = \begin{bmatrix} 2 & -1 & 1 & 0 \\ 3 & 5 & 0 & -2 \\ 1 & 1 & 0 & -3 \\ 4 & 0 & 3 & -1 \end{bmatrix}.$$

► **Solution.**

$$\begin{vmatrix} 2 & -1 & 1 & 0 \\ 3 & 5 & 0 & -2 \\ 1 & 1 & 0 & -3 \\ 4 & 0 & 3 & -1 \end{vmatrix} = 1 \cdot \begin{vmatrix} 3 & 5 & -2 \\ 1 & 1 & -3 \\ 4 & 0 & -1 \end{vmatrix} - 3 \cdot \begin{vmatrix} 3 & 5 & -2 \\ 1 & 1 & -3 \\ 4 & 0 & -1 \end{vmatrix} = -50 + 99 = 49.$$

## Why is the method of cofactor expansion not practical?

Because to compute a large  $n \times n$  determinant,

## Why is the method of cofactor expansion not practical?

Because to compute a large  $n \times n$  determinant,

- one reduces to  $n$  determinants of size  $(n - 1) \times (n - 1)$ ,
- then  $n(n - 1)$  determinants of size  $(n - 2) \times (n - 2)$ .
- and so on.

In the end, we have  $n! = n(n - 1) \cdots 3 \cdot 2 \cdot 1$  many numbers to add.

$\Rightarrow$  too much work.