

Lecture 19: Eigenvectors and eigenvalues; Diagonalization (Sections 5.1--5.2)

Thang Huynh, UC San Diego

2/28/2018

Eigenvectors and eigenvalues

A will be an $n \times n$ matrix.

► **Definition.** An **eigenvector** of A is a nonzero \mathbf{x} such that

$$A\mathbf{x} = \lambda\mathbf{x} \quad \text{for some scalar } \lambda.$$

The scalar λ is the corresponding **eigenvalue**.

Eigenvectors and eigenvalues

A will be an $n \times n$ matrix.

► **Definition.** An **eigenvector** of A is a nonzero \mathbf{x} such that

$$A\mathbf{x} = \lambda\mathbf{x} \quad \text{for some scalar } \lambda.$$

The scalar λ is the corresponding **eigenvalue**.

► **Definition.** Given λ , the set of all eigenvectors with eigenvalue λ is called the **eigenspace** of A corresponding to λ .

How to solve $A\mathbf{x} = \lambda\mathbf{x}$

Key observation:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$A\mathbf{x} - \lambda\mathbf{x} = 0$$

$$(A - \lambda I)\mathbf{x} = 0$$

This has a nonzero solution if and only if $\det(A - \lambda I) = 0$.

How to solve $A\mathbf{x} = \lambda\mathbf{x}$

Key observation:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$A\mathbf{x} - \lambda\mathbf{x} = 0$$

$$(A - \lambda I)\mathbf{x} = 0$$

This has a nonzero solution if and only if $\det(A - \lambda I) = 0$.

► **Recipe.** To find eigenvectors and eigenvalues of A .

- First, find the eigenvalues λ using $\det(A - \lambda I) = 0$.
- Then, for each eigenvalue λ , find corresponding eigenvectors by solving $(A - \lambda I)\mathbf{x} = 0$.

How to solve $Ax = \lambda x$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

How to solve $Ax = \lambda x$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial is

$$\det(A - \lambda I) = (3 - \lambda)(6 - \lambda)(2 - \lambda) \Rightarrow A \text{ has eigenvalues } 2, 3, 6.$$

How to solve $A\mathbf{x} = \lambda\mathbf{x}$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial is

$$\det(A - \lambda I) = (3 - \lambda)(6 - \lambda)(2 - \lambda) \Rightarrow A \text{ has eigenvalues } 2, 3, 6.$$

- For $\lambda_1 = 2$, $(A - 2I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = (2, -5/2, 1)^T$.

How to solve $A\mathbf{x} = \lambda\mathbf{x}$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial is

$$\det(A - \lambda I) = (3 - \lambda)(6 - \lambda)(2 - \lambda) \Rightarrow A \text{ has eigenvalues } 2, 3, 6.$$

- For $\lambda_1 = 2$, $(A - 2I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = (2, -5/2, 1)^T$.
- For $\lambda_2 = 3$, $(A - 3I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_2 = (1, 0, 0)^T$.

How to solve $A\mathbf{x} = \lambda\mathbf{x}$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial is

$$\det(A - \lambda I) = (3 - \lambda)(6 - \lambda)(2 - \lambda) \Rightarrow A \text{ has eigenvalues } 2, 3, 6.$$

- For $\lambda_1 = 2$, $(A - 2I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = (2, -5/2, 1)^T$.
- For $\lambda_2 = 3$, $(A - 3I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_2 = (1, 0, 0)^T$.
- For $\lambda_3 = 6$, $(A - 6I)\mathbf{x} = 0 \Rightarrow \mathbf{x}_3 = (2, 3, 0)^T$.

(Note that these three eigenvectors are linearly independent.)

Eigenvectors

Theorem. If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors A corresponding to *different eigenvalues*, then they are linearly independent.

Algebraic Multiplicity

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Algebraic Multiplicity

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial is

$\det(A - \lambda I) = (2 - \lambda)(\lambda - 2)(\lambda - 4)$. A has eigenvalues 2, 2, 4.

Since $\lambda = 2$ is a double root, it has **algebraic multiplicity** 2. (It's an exercise to find eigenvectors.)

Eigenvalues and Eigenvectors

An $n \times n$ matrix A has up to n different eigenvalues.

- For each eigenvalue λ , A has at least one eigenvector.
- If λ has multiplicity m , then A has up to m (independent) eigenvectors for λ .

Eigenvalues and Eigenvectors

Ideally, we would like to find a total of n (independent) eigenvectors of A .

Eigenvalues and Eigenvectors

Ideally, we would like to find a total of n (independent) eigenvectors of A . But there are two sources of trouble: eigenvalues can be

- complex numbers or
- repeated roots of characteristic polynomial.

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \text{ (Geometrically, what is the trouble?)}$$

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \text{ (Geometrically, what is the trouble?)}$$

► **Solution.** The characteristic polynomial

$$\det(A - \lambda I) = \lambda^2 + 1 \Rightarrow \lambda_1 = i \text{ and } \lambda_2 = -i.$$

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \text{ (Geometrically, what is the trouble?)}$$

► **Solution.** The characteristic polynomial

$$\det(A - \lambda I) = \lambda^2 + 1 \Rightarrow \lambda_1 = i \text{ and } \lambda_2 = -i.$$

- For $\lambda_1 = i$, $\begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x}_1 = \begin{bmatrix} i \\ 1 \end{bmatrix}.$

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \text{ (Geometrically, what is the trouble?)}$$

► **Solution.** The characteristic polynomial

$$\det(A - \lambda I) = \lambda^2 + 1 \Rightarrow \lambda_1 = i \text{ and } \lambda_2 = -i.$$

- For $\lambda_1 = i$, $\begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x}_1 = \begin{bmatrix} i \\ 1 \end{bmatrix}.$

- For $\lambda_2 = -i$, $\begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x}_2 = \begin{bmatrix} -i \\ 1 \end{bmatrix}.$

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

Eigenvalues and Eigenvectors

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

► **Solution.** The characteristic polynomial $\det(A - \lambda I) = (1 - \lambda)^2$.
So $\lambda = 1$ is the only eigenvalue (of multiplicity 2).

$$(A - \lambda I)\mathbf{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}\mathbf{x} = 0 \Rightarrow \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

So the eigenspace is $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

Diagonalization

Diagonal matrices are very easy to work with.

► **Example.** For instance, it is easy to compute their powers.

Let's consider $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$, then

Diagonalization

Diagonal matrices are very easy to work with.

► **Example.** For instance, it is easy to compute their powers.

Let's consider $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$, then

$$A^2 = \begin{bmatrix} 2^2 & & \\ & 3^2 & \\ & & 4^2 \end{bmatrix} \quad \text{and} \quad A^{100} = \begin{bmatrix} 2^{100} & & \\ & 3^{100} & \\ & & 4^{100} \end{bmatrix}.$$

Diagonalization

► **Example.** If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Diagonalization

► **Example.** If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$ Let's find eigenvalues and eigenvectors of A :

- $\lambda_1 = 4 \Rightarrow$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5 \Rightarrow$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Diagonalization

► **Example.** If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$ Let's find eigenvalues and eigenvectors of A :

- $\lambda_1 = 4 \Rightarrow$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5 \Rightarrow$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Key observation: if \mathbf{v} is an eigenvector of A corresponding to an eigenvalue λ ,

$$A^m \mathbf{v} = \lambda^m \mathbf{v}.$$

Diagonalization

Let $B = A^{100} = [\mathbf{b}_1 \ \mathbf{b}_2]$. Then $\mathbf{b}_1 = A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\mathbf{b}_2 = A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Diagonalization

Let $B = A^{100} = [\mathbf{b}_1 \ \mathbf{b}_2]$. Then $\mathbf{b}_1 = A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\mathbf{b}_2 = A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Therefore, finding A^{100} is equivalent to finding $A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and

$$A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Diagonalization

Let $B = A^{100} = [\mathbf{b}_1 \ \mathbf{b}_2]$. Then $\mathbf{b}_1 = A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\mathbf{b}_2 = A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Therefore, finding A^{100} is equivalent to finding $A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and

$$A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Observe that

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} = -\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \end{bmatrix} = -\mathbf{v}_1 + 2\mathbf{v}_2.$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = -4^{100} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \cdot 5^{100} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Diagonalization

Let $B = A^{100} = [\mathbf{b}_1 \ \mathbf{b}_2]$. Then $\mathbf{b}_1 = A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\mathbf{b}_2 = A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Therefore, finding A^{100} is equivalent to finding $A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and

$$A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Observe that

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} = -\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \end{bmatrix} = -\mathbf{v}_1 + 2\mathbf{v}_2.$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = -4^{100} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \cdot 5^{100} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Exercise: find $A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.