

# Lecture 18: Applications of Determinants; Eigenvectors and eigenvalues (Sections 4.3--5.1)

---

Thang Huynh, UC San Diego

2/26/2018

## Computation of $A^{-1}$

► Example.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

## Computation of $A^{-1}$

► Example.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

►  $C_{11} = d$  is the cofactor of  $a$ .

## Computation of $A^{-1}$

► Example.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

- $C_{11} = d$  is the cofactor of  $a$ .
- $C_{12} = -c$  is the cofactor of  $b$ .

## Computation of $A^{-1}$

► Example.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

- $C_{11} = d$  is the cofactor of  $a$ .
- $C_{12} = -c$  is the cofactor of  $b$ .
- $C_{21} = -b$  is the cofactor of  $c$ .

## Computation of $A^{-1}$

► Example.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

- $C_{11} = d$  is the cofactor of  $a$ .
- $C_{12} = -c$  is the cofactor of  $b$ .
- $C_{21} = -b$  is the cofactor of  $c$ .
- $C_{22} = a$  is the cofactor of  $d$ .

## Computation of $A^{-1}$

► **Example.**

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

- $C_{11} = d$  is the cofactor of  $a$ .
- $C_{12} = -c$  is the cofactor of  $b$ .
- $C_{21} = -b$  is the cofactor of  $c$ .
- $C_{22} = a$  is the cofactor of  $d$ .

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{\det(A)} \begin{bmatrix} C_{11} & C_{21} \\ C_{12} & C_{22} \end{bmatrix} = \frac{1}{\det(A)} C^T.$$

## Computation of $A^{-1}$

$A^{-1}$  divides (the transpose) of the cofactors of  $A$  by  $\det(A)$ .

$$A^{-1} = \frac{C^T}{\det(A)} \quad \text{means} \quad (A^{-1})_{ij} = \frac{C_{ji}}{\det(A)}.$$

(See the textbook for the proof.)



## Computation of $A^{-1}$

► **Example.** Find the inverse of the following matrix using cofactor formula:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

## Computation of $A^{-1}$

► **Example.** Find the inverse of the following matrix using cofactor formula:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

► **Solution.**

$$A^{-1} = \frac{C^T}{\det A} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

## The solution of $Ax = b$

Suppose  $A$  is invertible.

$$Ax = b \Rightarrow x = A^{-1}b = \frac{C^T b}{\det A}.$$

## The solution of $Ax = b$

Suppose  $A$  is invertible.

$$Ax = b \Rightarrow x = A^{-1}b = \frac{C^T b}{\det A}.$$

► **Cramer's rule.** The  $j$ th component of  $x = A^{-1}b$  is the ratio

$$x_j = \frac{\det B_j}{\det A},$$

where

$$B_j = \begin{bmatrix} a_{11} & a_{12} & b_1 & a_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & b_n & a_{nn} \end{bmatrix} \text{ has } b \text{ in column } j.$$

## The solution of $Ax = b$

► Example. Solve

$$x_1 + 3x_2 = 0$$

$$2x_1 + 4x_2 = 6$$

► Solution.

$$x_1 = \frac{\begin{vmatrix} 0 & 3 \\ 6 & 4 \end{vmatrix}}{\begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix}} = \frac{-18}{-2} = 9, \quad x_2 = \frac{\begin{vmatrix} 1 & 0 \\ 2 & 6 \end{vmatrix}}{\begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix}} = \frac{6}{-2} = -3.$$

## Volume of a box

The determinant equals the volume.

## Eigenvectors and eigenvalues

$A$  will be an  $n \times n$  matrix.

► **Definition.** An **eigenvector** of  $A$  is a nonzero  $x$  such that

$$Ax = \lambda x \quad \text{for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

## Eigenvectors and eigenvalues

$A$  will be an  $n \times n$  matrix.

► **Definition.** An **eigenvector** of  $A$  is a nonzero  $x$  such that

$$Ax = \lambda x \quad \text{for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

► **Example.** Verify that  $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .



## Eigenvectors and eigenvalues

$A$  will be an  $n \times n$  matrix.

► **Definition.** An **eigenvector** of  $A$  is a nonzero  $x$  such that

$$Ax = \lambda x \quad \text{for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

► **Example.** Verify that  $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

► **Solution.**

$$\begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 4 \\ -8 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ -2 \end{bmatrix}.$$

## Eigenvectors and eigenvalues

$A$  will be an  $n \times n$  matrix.

► **Definition.** An **eigenvector** of  $A$  is a nonzero  $x$  such that

$$Ax = \lambda x \quad \text{for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

► **Example.** Verify that  $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

► **Solution.**

$$\begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 4 \\ -8 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ -2 \end{bmatrix}.$$

► **Example.** Use your geometric understanding to find the eigenvectors and eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Eigenvectors and eigenvalues

- **Definition.** Given  $\lambda$ , the set of all eigenvectors with eigenvalue  $\lambda$  is called the **eigenspace** of  $A$  corresponding to  $\lambda$ .
- **Example.** Let  $P$  be the projection matrix corresponding to orthogonal projection onto the subspace  $V$ . What are the eigenvalues and eigenvectors of  $P$ ?

## Eigenvectors and eigenvalues

- ▶ **Definition.** Given  $\lambda$ , the set of all eigenvectors with eigenvalue  $\lambda$  is called the **eigenspace** of  $A$  corresponding to  $\lambda$ .
- ▶ **Example.** Let  $P$  be the projection matrix corresponding to orthogonal projection onto the subspace  $V$ . What are the eigenvalues and eigenvectors of  $P$ ?
- ▶ **Solution.**
  - For every vector  $x$  in  $V$ ,  $Px = x$ .

## Eigenvectors and eigenvalues

- ▶ **Definition.** Given  $\lambda$ , the set of all eigenvectors with eigenvalue  $\lambda$  is called the **eigenspace** of  $A$  corresponding to  $\lambda$ .
- ▶ **Example.** Let  $P$  be the projection matrix corresponding to orthogonal projection onto the subspace  $V$ . What are the eigenvalues and eigenvectors of  $P$ ?
- ▶ **Solution.**
  - For every vector  $\mathbf{x}$  in  $V$ ,  $P\mathbf{x} = \mathbf{x}$ . These are the eigenvectors with eigenvalue 1.
  - For every vector  $\mathbf{x}$  orthogonal to  $V$ ,  $P\mathbf{x} = 0$ .

## Eigenvectors and eigenvalues

- **Definition.** Given  $\lambda$ , the set of all eigenvectors with eigenvalue  $\lambda$  is called the **eigenspace** of  $A$  corresponding to  $\lambda$ .
- **Example.** Let  $P$  be the projection matrix corresponding to orthogonal projection onto the subspace  $V$ . What are the eigenvalues and eigenvectors of  $P$ ?
- **Solution.**
  - For every vector  $\mathbf{x}$  in  $V$ ,  $P\mathbf{x} = \mathbf{x}$ . These are the eigenvectors with eigenvalue 1.
  - For every vector  $\mathbf{x}$  orthogonal to  $V$ ,  $P\mathbf{x} = 0$ . These are the eigenvectors with eigenvalue 0.

## Eigenvectors and eigenvalues

- **Definition.** Given  $\lambda$ , the set of all eigenvectors with eigenvalue  $\lambda$  is called the **eigenspace** of  $A$  corresponding to  $\lambda$ .
- **Example.** Let  $P$  be the projection matrix corresponding to orthogonal projection onto the subspace  $V$ . What are the eigenvalues and eigenvectors of  $P$ ?
- **Solution.**
  - For every vector  $\mathbf{x}$  in  $V$ ,  $P\mathbf{x} = \mathbf{x}$ . These are the eigenvectors with eigenvalue 1.
  - For every vector  $\mathbf{x}$  orthogonal to  $V$ ,  $P\mathbf{x} = 0$ . These are the eigenvectors with eigenvalue 0.

The eigenspace of  $\lambda = 1$  is  $V$ . The eigenspace of  $\lambda = 0$  is  $V^\perp$ .

## How to solve $A\mathbf{x} = \lambda\mathbf{x}$

Key observation:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$A\mathbf{x} - \lambda\mathbf{x} = 0$$

$$(A - \lambda I)\mathbf{x} = 0$$

This has a nonzero solution if and only if  $\det(A - \lambda I) = 0$ .



## How to solve $A\mathbf{x} = \lambda\mathbf{x}$

Key observation:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$A\mathbf{x} - \lambda\mathbf{x} = 0$$

$$(A - \lambda I)\mathbf{x} = 0$$

This has a nonzero solution if and only if  $\det(A - \lambda I) = 0$ .

► **Recipe.** To find eigenvectors and eigenvalues of  $A$ .

- First, find the eigenvalues  $\lambda$  using  $\det(A - \lambda I) = 0$ .
- Then, for each eigenvalue  $\lambda$ , find corresponding eigenvectors by solving  $(A - \lambda I)\mathbf{x} = 0$ .

## How to solve $Ax = \lambda x$

► **Example.** Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}.$$

# Eigenvectors

**Theorem.** If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors  $A$  corresponding to different eigenvalues, then they are linearly independent.