

Lecture 16: Determinants (Section 4.1-4.2)

Thang Huynh, UC San Diego

2/16/2018

For the next few lectures, **all matrices are square!**

► Recall that

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

For the next few lectures, **all matrices are square!**

► Recall that

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

The **determinant** of

- a 2×2 matrix is $\det \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = ad - bc$,
- a 1×1 matrix is $\det([a]) = a$.

We will write both $\det \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right)$ and $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$

A is invertible $\iff \det(A) \neq 0$

The determinant is characterized by:

- the normalization $\det I = 1$,

A is invertible $\iff \det(A) \neq 0$

The determinant is characterized by:

- the normalization $\det I = 1$,
- and how it is affected by elementary row operations:

A is invertible $\iff \det(A) \neq 0$

The determinant is characterized by:

- the normalization $\det I = 1$,
- and how it is affected by elementary row operations:
 - (replacement) Add one row to a multiple of another row.
Does not change the determinant.

A is invertible $\iff \det(A) \neq 0$

The determinant is characterized by:

- the normalization $\det I = 1$,
- and how it is affected by elementary row operations:
 - (replacement) Add one row to a multiple of another row.
Does not change the determinant.
 - (interchange) Interchange two rows.
Reverses the sign of the determinant.

A is invertible $\iff \det(A) \neq 0$

The determinant is characterized by:

- the normalization $\det I = 1$,
- and how it is affected by elementary row operations:
 - (replacement) Add one row to a multiple of another row.
Does not change the determinant.
 - (interchange) Interchange two rows.
Reverses the sign of the determinant.
 - (scaling) Multiply all entries in a row by s .
Multiplies the determinant by s .

Examples

► **Example.** Compute $\begin{vmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 7 \end{vmatrix}$.

Examples

► **Example.** Compute $\begin{vmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 7 \end{vmatrix}$.

► **Solution.**

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 7 \end{vmatrix} \xrightarrow[R_2 \rightarrow \frac{1}{2}R_2]{=} 2 \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 7 \end{vmatrix} \xrightarrow[R_3 \rightarrow \frac{1}{7}R_3]{=} 14 \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 14.$$

Examples

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 7 \end{vmatrix}$.

Examples

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 7 \end{vmatrix}$.

► **Solution.**

$$\begin{vmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 7 \end{vmatrix} \xrightarrow[R_2 \rightarrow \frac{1}{2}R_2]{=} 2 \begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 7 \end{vmatrix} \xrightarrow[R_3 \rightarrow \frac{1}{7}R_3]{=} 14 \begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{vmatrix}$$

$$\xrightarrow[R_1 \rightarrow R_1 - 3R_3, R_2 \rightarrow R_2 - 2R_3]{=} 14 \begin{vmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} \xrightarrow[R_1 \rightarrow R_1 - 2R_2]{=} 14 \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 14$$

The determinant of a triangular matrix is the product of the diagonal entries

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$.

The determinant of a triangular matrix is the product of the diagonal entries

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$.

► **Solution.**

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} \xrightarrow{R_2 \rightarrow R_2 - 3R_1, R_3 \rightarrow R_3 - 2R_1} \begin{vmatrix} 1 & 2 & 0 \\ 0 & -7 & 2 \\ 0 & -4 & 1 \end{vmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - \frac{4}{7}R_2} \begin{vmatrix} 1 & 2 & 0 \\ 0 & -7 & 2 \\ 0 & 0 & -\frac{1}{7} \end{vmatrix}$$

$$= 1(-7)\left(-\frac{1}{7}\right) = 1.$$

The determinant of a triangular matrix is the product of the diagonal entries

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$.

► **Solution.**

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} \xrightarrow{R_2 \rightarrow R_2 - 3R_1, R_3 \rightarrow R_3 - 2R_1} \begin{vmatrix} 1 & 2 & 0 \\ 0 & -7 & 2 \\ 0 & -4 & 1 \end{vmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - \frac{4}{7}R_2} \begin{vmatrix} 1 & 2 & 0 \\ 0 & -7 & 2 \\ 0 & 0 & -\frac{1}{7} \end{vmatrix}$$

$$= 1(-7)\left(-\frac{1}{7}\right) = 1.$$

► **Example.** Prove the determinant formula for $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$.

► **Example.** Prove the determinant formula for $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$.

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix}$.

Properties of determinants

The following important properties follow from the behavior under row operations:

Properties of determinants

The following important properties follow from the behavior under row operations:

- $\det(A) = 0 \iff A$ is not invertible.

Properties of determinants

The following important properties follow from the behavior under row operations:

- $\det(A) = 0 \iff A$ is not invertible.

Why? Because $\det(A) = 0$ if and only if, in an echelon form, a diagonal entry is zero (that is, a pivot is missing).

Properties of determinants

The following important properties follow from the behavior under row operations:

- $\det(A) = 0 \iff A$ is not invertible.

Why? Because $\det(A) = 0$ if and only if, in an echelon form, a diagonal entry is zero (that is, a pivot is missing).

- $\det(AB) = \det(A) \det(B)$.

Properties of determinants

The following important properties follow from the behavior under row operations:

- $\det(A) = 0 \iff A$ is not invertible.

Why? Because $\det(A) = 0$ if and only if, in an echelon form, a diagonal entry is zero (that is, a pivot is missing).

- $\det(AB) = \det(A) \det(B)$.
- $\det(A^{-1}) = \frac{1}{\det(A)}$

Properties of determinants

The following important properties follow from the behavior under row operations:

- $\det(A) = 0 \iff A$ is not invertible.

Why? Because $\det(A) = 0$ if and only if, in an echelon form, a diagonal entry is zero (that is, a pivot is missing).

- $\det(AB) = \det(A) \det(B)$.
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$

► **Example.** Recall that if $AB = 0$, it does not follow that $A = 0$ or $B = 0$. However, show that $\det(A) = 0$ or $\det(B) = 0$.

- **Example.** Recall that if $AB = 0$, it does not follow that $A = 0$ or $B = 0$. However, show that $\det(A) = 0$ or $\det(B) = 0$.
- **Solution.** Follow from $\det(A) \det(B) = \det(AB) = 0$.

- **Example.** Recall that if $AB = 0$, it does not follow that $A = 0$ or $B = 0$. However, show that $\det(A) = 0$ or $\det(B) = 0$.
- **Solution.** Follow from $\det(A) \det(B) = \det(AB) = 0$.
- **Example.** Suppose A is a 3×3 matrix with $\det(A) = 5$. What is $\det(2A)$?

► **Example.** Recall that if $AB = 0$, it does not follow that $A = 0$ or $B = 0$. However, show that $\det(A) = 0$ or $\det(B) = 0$.

► **Solution.** Follow from $\det(A) \det(B) = \det(AB) = 0$.

► **Example.** Suppose A is a 3×3 matrix with $\det(A) = 5$. What is $\det(2A)$?

► **Solution.** A has three rows. Multiplying all 3 of them by 2 produces $2A$. Hence, $\det(2A) = 2^3 \det(A) = 40$.

A “bad” way to compute determinants

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$ by **cofactor expansion**.

A “bad” way to compute determinants

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$ by **cofactor expansion**.

► **Solution.** We expand by the second column:

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = -2 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & 1 \end{vmatrix} - 0 \begin{vmatrix} 1 & 0 \\ 3 & 2 \end{vmatrix} = -2(-1) + (-1)1 - 0 = 1.$$

A “bad” way to compute determinants

► **Example.** Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$ by **cofactor expansion**.

► **Solution.** We expand by the second column:

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = -2 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & 1 \end{vmatrix} - 0 \begin{vmatrix} 1 & 0 \\ 3 & 2 \end{vmatrix} = -2(-1) + (-1)1 - 0 = 1.$$

► **Solution.** We expand by the third column:

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix} + (-2) \begin{vmatrix} 1 & 2 \\ 2 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 3 & -1 \end{vmatrix} = 0 - 2(-4) + 1(-7) = 1.$$

Why is the method of cofactor expansion not practical?

Because to compute a large $n \times n$ determinant,

Why is the method of cofactor expansion not practical?

Because to compute a large $n \times n$ determinant,

- one reduces to n determinants of size $(n - 1) \times (n - 1)$,
- then $n(n - 1)$ determinants of size $(n - 2) \times (n - 2)$.
- and so on.

In the end, we have $n! = n(n - 1) \cdots 3 \cdot 2 \cdot 1$ many numbers to add.

\Rightarrow too much work.