

Lecture 11: Orthogonality (Section 3.1 - 3.2)

Thang Huynh, UC San Diego

2/5/2018

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W .

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W . (Is W^\perp a vector space?)

The fundamental theorem, part II

- **Definition.** Let W be a subspace of \mathbb{R}^n , and \mathbf{v} in \mathbb{R}^n .
- \mathbf{v} is **orthogonal** to W , if $\mathbf{v} \cdot \mathbf{w} = 0$ for all \mathbf{w} in W .
 - Another subspace V is **orthogonal** to W , if every vector in V is orthogonal to W .
 - The **orthogonal complement** of W is the space W^\perp of all vectors in \mathbb{R}^n that are orthogonal to W . (Is W^\perp a vector space?)

► **Example.** Consider $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$, then $N(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$

and $C(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ are orthogonal subspaces.

The fundamental theorem, part II

Theorem. (Fundamental Theorem of Linear Algebra, Part II)

- $N(A)$ is orthogonal to $C(A^T)$. (The two spaces are orthogonal complements)
- $N(A^T)$ is orthogonal to $C(A)$. (The two spaces are orthogonal complements)

► Proof.

Theorem. (Fundamental Theorem of Linear Algebra, Part I)

Let A be an $m \times n$ matrix of rank r .

- $\dim C(A) = \dim C(A^T) = r$.
- $\dim N(A) = n - r$
- $\dim N(A^T) = m - r$.

► **Example.** Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

► **Example.** Find all vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$.

► **Example.** Let $V = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b = 2c \right\}$. Find a basis for the orthogonal complement of V .

A new perspective on $Ax = b$

$Ax = b$ is solvable

$\Leftrightarrow b$ is in $C(A)$

$\Leftrightarrow b$ is orthogonal to $N(A^T)$

A new perspective on $Ax = b$

$Ax = b$ is solvable

$\Leftrightarrow b$ is in $C(A)$

$\Leftrightarrow b$ is orthogonal to $N(A^T)$

► **Example.** Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix}$. For which b does $Ax = b$ have a solution?

Orthogonal bases

- Recall that if $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal, then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.

Orthogonal bases

- ▶ Recall that if $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal, then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.
- ▶ **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthogonal basis** if the vectors are pairwise orthogonal.

Orthogonal bases

► Recall that if $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal, then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.

► **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthogonal basis** if the vectors are pairwise orthogonal.

► **Example.** The standard basis $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ is an orthogonal basis for \mathbb{R}^3 . (Why?)

Orthogonal bases

► Recall that if $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal, then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.

► **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthogonal basis** if the vectors are pairwise orthogonal.

► **Example.** The standard basis $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ is an

orthogonal basis for \mathbb{R}^3 . (Why?)

► **Example.** Are the vectors $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ an orthogonal

basis for \mathbb{R}^3 ?

Orthogonal bases

► Recall that if $\mathbf{v}_1, \dots, \mathbf{v}_n$ are nonzero and pairwise orthogonal, then $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent.

► **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthogonal basis** if the vectors are pairwise orthogonal.

► **Example.** The standard basis $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ is an

orthogonal basis for \mathbb{R}^3 . (Why?)

► **Example.** Are the vectors $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ an orthogonal

basis for \mathbb{R}^3 ? (Do we need to check that the three vectors are independent?)

Orthogonal bases

► **Example.** Suppose $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthogonal basis of V , and \mathbf{w} is in V . Find c_1, \dots, c_n such that

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n.$$

Orthogonal bases

► **Example.** Suppose $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthogonal basis of V , and \mathbf{w} is in V . Find c_1, \dots, c_n such that

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n.$$

► **Solution.** Take the dot product of \mathbf{v}_1 with both sides

$$\begin{aligned}\mathbf{v}_1 \cdot \mathbf{w} &= \mathbf{v}_1 \cdot (c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n) \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 + c_2 \mathbf{v}_1 \cdot \mathbf{v}_2 + \dots + c_n \mathbf{v}_1 \cdot \mathbf{v}_n \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1.\end{aligned}$$

Hence, $c_1 = \frac{\mathbf{v}_1 \cdot \mathbf{w}}{\mathbf{v}_1 \cdot \mathbf{v}_1}$.

Orthogonal bases

► **Example.** Suppose $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthogonal basis of V , and \mathbf{w} is in V . Find c_1, \dots, c_n such that

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n.$$

► **Solution.** Take the dot product of \mathbf{v}_1 with both sides

$$\begin{aligned}\mathbf{v}_1 \cdot \mathbf{w} &= \mathbf{v}_1 \cdot (c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n) \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 + c_2 \mathbf{v}_1 \cdot \mathbf{v}_2 + \dots + c_n \mathbf{v}_1 \cdot \mathbf{v}_n \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1.\end{aligned}$$

Hence, $c_1 = \frac{\mathbf{v}_1 \cdot \mathbf{w}}{\mathbf{v}_1 \cdot \mathbf{v}_1}$. In general, $c_j = \frac{\mathbf{v}_j \cdot \mathbf{w}}{\mathbf{v}_j \cdot \mathbf{v}_j}$.

Orthogonal bases

► **Example.** Express $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ in terms of the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

Orthogonal bases

► **Example.** Express $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ in terms of the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

► **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthonormal basis** if the vectors are orthogonal and have length 1.

Orthogonal bases

► **Example.** Express $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ in terms of the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

► **Definition.** A basis $\mathbf{v}_1, \dots, \mathbf{v}_n$ of a vector space V is an **orthonormal basis** if the vectors are orthogonal and have length 1.

► **Example.** The standard basis $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ is an orthonormal basis for \mathbb{R}^3 .

Orthogonal bases

If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthonormal basis of V , and \mathbf{w} is in V , then

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n \quad \text{with} \quad c_j = \mathbf{v}_j \cdot \mathbf{w}.$$

Orthogonal bases

If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthonormal basis of V , and \mathbf{w} is in V , then

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n \quad \text{with} \quad c_j = \mathbf{v}_j \cdot \mathbf{w}.$$

► **Example.** Is the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ orthonormal?

Orthogonal bases

If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthonormal basis of V , and \mathbf{w} is in V , then

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n \quad \text{with} \quad c_j = \mathbf{v}_j \cdot \mathbf{w}.$$

► **Example.** Is the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ orthonormal? If not, normalize the vectors to produce an orthonormal basis.

Orthogonal bases

If $\mathbf{v}_1, \dots, \mathbf{v}_n$ is an orthonormal basis of V , and \mathbf{w} is in V , then

$$\mathbf{w} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n \quad \text{with} \quad c_j = \mathbf{v}_j \cdot \mathbf{w}.$$

► **Example.** Is the basis $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ orthonormal? If not, normalize the vectors to produce an orthonormal basis.

► **Example.** Express $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ in terms of the basis

$$\left\{ \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector x onto vector y is

$$\hat{x} = \frac{x \cdot y}{y \cdot y} y.$$

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector x onto vector y is

$$\hat{x} = \frac{x \cdot y}{y \cdot y} y.$$

(Geometric meaning?)

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector x onto vector y is

$$\hat{x} = \frac{x \cdot y}{y \cdot y} y.$$

(Geometric meaning?)

- The vector \hat{x} is the closest vector x , which is in $\text{span}\{y\}$.

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector \mathbf{x} onto vector \mathbf{y} is

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}.$$

(Geometric meaning?)

- The vector $\hat{\mathbf{x}}$ is the closest vector \mathbf{x} , which is in $\text{span}\{\mathbf{y}\}$.
- Characterized by the error $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ is orthogonal to $\text{span}\{\mathbf{y}\}$.

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector \mathbf{x} onto vector \mathbf{y} is

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}.$$

(Geometric meaning?)

- The vector $\hat{\mathbf{x}}$ is the closest vector \mathbf{x} , which is in $\text{span}\{\mathbf{y}\}$.
- Characterized by the error $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ is orthogonal to $\text{span}\{\mathbf{y}\}$.
- To find the formula for $\hat{\mathbf{x}}$, start with $\hat{\mathbf{x}} = c\mathbf{y}$. Find c such that

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector \mathbf{x} onto vector \mathbf{y} is

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}.$$

(Geometric meaning?)

- The vector $\hat{\mathbf{x}}$ is the closest vector \mathbf{x} , which is in $\text{span}\{\mathbf{y}\}$.
- Characterized by the error $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ is orthogonal to $\text{span}\{\mathbf{y}\}$.
- To find the formula for $\hat{\mathbf{x}}$, start with $\hat{\mathbf{x}} = c\mathbf{y}$. Find c such that

$$(\mathbf{x} - \hat{\mathbf{x}}) \cdot \mathbf{y} = (\mathbf{x} - c\mathbf{y}) \cdot \mathbf{y} = \mathbf{x} \cdot \mathbf{y} - c\mathbf{y} \cdot \mathbf{y} = 0.$$

It follows that $c = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}}$.

Orthogonal projections

► **Definition.** The **orthogonal projection** of vector \mathbf{x} onto vector \mathbf{y} is

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}.$$

(Geometric meaning?)

- The vector $\hat{\mathbf{x}}$ is the closest vector \mathbf{x} , which is in $\text{span}\{\mathbf{y}\}$.
- Characterized by the error $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ is orthogonal to $\text{span}\{\mathbf{y}\}$.
- To find the formula for $\hat{\mathbf{x}}$, start with $\hat{\mathbf{x}} = c\mathbf{y}$. Find c such that

$$(\mathbf{x} - \hat{\mathbf{x}}) \cdot \mathbf{y} = (\mathbf{x} - c\mathbf{y}) \cdot \mathbf{y} = \mathbf{x} \cdot \mathbf{y} - c\mathbf{y} \cdot \mathbf{y} = 0.$$

It follows that $c = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}}$.

- \mathbf{x}^\perp is also called the component of \mathbf{x} orthogonal to \mathbf{y} .

Orthogonal projections

► **Example.** What is the orthogonal projection of $\mathbf{x} = \begin{bmatrix} -8 \\ 4 \end{bmatrix}$ onto

$$\mathbf{y} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}?$$

Orthogonal projections

► **Example.** What is the orthogonal projection of $\mathbf{x} = \begin{bmatrix} -8 \\ 4 \end{bmatrix}$ onto $\mathbf{y} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$?

► **Example.** What is the orthogonal projection of $\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}$ onto each of the vectors $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$?