

# Lecture 26: Singular Value Decompositions (Sections 6.3)

---

Thang Huynh, UC San Diego

3/16/2018

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$
- columns of  $V$  ( $n$  by  $n$ ) are eigenvectors of  $A^TA$

# Singular Value Decompositions

Recall we can decompose a matrix  $A$  as  $LU$  or  $QR$ .

(SVD) Any  $m \times n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

- columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$
- columns of  $V$  ( $n$  by  $n$ ) are eigenvectors of  $A^TA$
- The  $r$  singular values on the diagonal of  $\Sigma$  ( $m$  by  $n$ ) are the square roots of the nonzero eigenvalues of both  $AA^T$  and  $A^TA$ .

## SVD Theory

$$AV = U\Sigma \Rightarrow A\mathbf{v}_j = \sigma_j \mathbf{u}_j, \quad j = 1, 2, \dots, r.$$

- If  $\sigma_j = 0$ ,  $A\mathbf{v}_j = 0$  and  $\mathbf{v}_j$  is in  $N(A)$ .  
The corresponding  $\mathbf{u}_j$  is in  $N(A^T)$

## SVD Theory

$$AV = U\Sigma \Rightarrow A\mathbf{v}_j = \sigma_j \mathbf{u}_j, \quad j = 1, 2, \dots, r.$$

- If  $\sigma_j = 0$ ,  $A\mathbf{v}_j = 0$  and  $\mathbf{v}_j$  is in  $N(A)$ .  
The corresponding  $\mathbf{u}_j$  is in  $N(A^T)$
- Else,  $\mathbf{v}_j$  is in  $C(A^T)$  and the corresponding  $\mathbf{u}_j$  is in  $C(A)$ .

## SVD Theory

$$AV = U\Sigma \Rightarrow A\mathbf{v}_j = \sigma_j \mathbf{u}_j, \quad j = 1, 2, \dots, r.$$

- If  $\sigma_j = 0$ ,  $A\mathbf{v}_j = 0$  and  $\mathbf{v}_j$  is in  $N(A)$ .  
The corresponding  $\mathbf{u}_j$  is in  $N(A^T)$
- Else,  $\mathbf{v}_j$  is in  $C(A^T)$  and the corresponding  $\mathbf{u}_j$  is in  $C(A)$ .
- Number of nonzero  $\sigma_j = \text{rank of } A$ .



## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$ .

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$ . (Then  $A$  has rank 2.)

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$ . (Then  $A$  has rank 2.)

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix} \Rightarrow \lambda_1 = 8 \text{ and } \mathbf{v}_1 = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}; \lambda_2 = 2 \text{ and } \mathbf{v}_2 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}.$$

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$ . (Then  $A$  has rank 2.)

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix} \Rightarrow \lambda_1 = 8 \text{ and } \mathbf{v}_1 = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}; \lambda_2 = 2 \text{ and } \mathbf{v}_2 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}.$$

**Step 2:** compute  $A\mathbf{v}_i$ . If  $\sigma_i \neq 0$ , get  $\mathbf{u}_i$ . Else, find  $\mathbf{u}_i$  from  $N(A^T)$ .

$$A\mathbf{v}_1 = \begin{bmatrix} 2\sqrt{2} \\ 0 \end{bmatrix} = \sigma_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow \mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$A\mathbf{v}_2 = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix} = \sigma_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} \Rightarrow \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$ . (Then  $A$  has rank 2.)

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix} \Rightarrow \lambda_1 = 8 \text{ and } \mathbf{v}_1 = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}; \lambda_2 = 2 \text{ and } \mathbf{v}_2 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}.$$

**Step 2:** compute  $A\mathbf{v}_i$ . If  $\sigma_i \neq 0$ , get  $\mathbf{u}_i$ . Else, find  $\mathbf{u}_i$  from  $N(A^T)$ .

$$A\mathbf{v}_1 = \begin{bmatrix} 2\sqrt{2} \\ 0 \end{bmatrix} = \sigma_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow \mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$A\mathbf{v}_2 = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix} = \sigma_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} \Rightarrow \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$A = U\Sigma V^T = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2\sqrt{2} & 0 \\ 0 & \sqrt{2} \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}.$$

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ .

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . ( $A$  has rank 2 and  $A_{2 \times 3} V_{3 \times 3} = U_{2 \times 2} \Sigma_{2 \times 3}$ .)

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . ( $A$  has rank 2 and

$$A_{2 \times 3} V_{3 \times 3} = U_{2 \times 2} \Sigma_{2 \times 3}.)$$

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$



## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . ( $A$  has rank 2 and

$$A_{2 \times 3} V_{3 \times 3} = U_{2 \times 2} \Sigma_{2 \times 3}.)$$

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

$$\bullet \lambda_1 = 3 \Rightarrow \mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . ( $A$  has rank 2 and

$$A_{2 \times 3} V_{3 \times 3} = U_{2 \times 2} \Sigma_{2 \times 3}.)$$

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

$$\begin{aligned} \bullet \lambda_1 = 3 &\Rightarrow \mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \\ \bullet \lambda_2 = 1 &\Rightarrow \mathbf{v}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \end{aligned}$$

## Singular Value Decompositions

► **Example.** Find SVD of  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . ( $A$  has rank 2 and

$$A_{2 \times 3} V_{3 \times 3} = U_{2 \times 2} \Sigma_{2 \times 3}.$$

► **Solution.** **Step 1:** Find eigenvalues and eigenvectors of  $A^T A$ :

$$A^T A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

$$\bullet \lambda_1 = 3 \Rightarrow \mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

$$\bullet \lambda_2 = 1 \Rightarrow \mathbf{v}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \quad \lambda_3 = 0 \Rightarrow \mathbf{v}_3 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

## Singular Value Decompositions

**Step 2:** compute  $A\mathbf{v}_i$ . If  $\sigma_i \neq 0$ , get  $\mathbf{u}_i$ . Else, find  $\mathbf{u}_i$  from  $N(A^T)$ .

$$A\mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 3 \\ 3 \end{bmatrix} = \sigma_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

## Singular Value Decompositions

**Step 2:** compute  $A\mathbf{v}_i$ . If  $\sigma_i \neq 0$ , get  $\mathbf{u}_i$ . Else, find  $\mathbf{u}_i$  from  $N(A^T)$ .

$$A\mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 3 \\ 3 \end{bmatrix} = \sigma_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$A\mathbf{v}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \sigma_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

## Singular Value Decompositions

**Step 2:** compute  $A\mathbf{v}_i$ . If  $\sigma_i \neq 0$ , get  $\mathbf{u}_i$ . Else, find  $\mathbf{u}_i$  from  $N(A^T)$ .

$$A\mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{bmatrix} 3 \\ 3 \end{bmatrix} = \sigma_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$A\mathbf{v}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \sigma_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

## Least squares and the pseudoinverse

A least squares solution  $\hat{\mathbf{x}}$  of the linear system

$$A\mathbf{x} = \mathbf{b}$$

is the one minimizing  $\|A\mathbf{x} - \mathbf{b}\|$ . To solve this, we can solve the associate normal system

$$A^T A \hat{\mathbf{x}} = A^T \mathbf{b}.$$

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$



## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**?

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ .

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ . The optimal solution of the  $A\mathbf{x} = \mathbf{b}$  is the vector  $\hat{\mathbf{x}}$  that has the minimum length. We denote this solution by  $\mathbf{x}^\dagger$ , and the matrix which produces  $\mathbf{x}^\dagger$  from  $\mathbf{b}$  is called the **pseudoinverse** of  $A$  and is denoted by  $A^\dagger$ .

## Least squares and the pseudoinverse

- If  $A$  has full rank,  $A^T A$  is invertible and the system  $A\mathbf{x} = \mathbf{b}$  has **unique** least squares solution

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}.$$

- But how if  $A$  **does not have full rank**? The least squares problem still has a solution, but it is no longer unique, i.e., there are many  $\mathbf{x}$  that minimize  $\|A\mathbf{x} - \mathbf{b}\|$ . The optimal solution of the  $A\mathbf{x} = \mathbf{b}$  is the vector  $\hat{\mathbf{x}}$  that has the minimum length. We denote this solution by  $\mathbf{x}^\dagger$ , and the matrix which produces  $\mathbf{x}^\dagger$  from  $\mathbf{b}$  is called the **pseudoinverse** of  $A$  and is denoted by  $A^\dagger$ .

$$\mathbf{x}^\dagger = A^\dagger \mathbf{b}.$$

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ?

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$ .

## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$ .

The SVD of  $A$  is

$$A = [1][3 \ 0 \ 0] \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}.$$



## Least squares and the pseudoinverse

How to find  $A^\dagger$ ? If the SVD of  $A$  is  $A = U\Sigma V^T$ , then

$$A^\dagger = V\Sigma^\dagger U^T.$$

The singular values  $\sigma_1, \dots, \sigma_r$  are on the diagonal of  $\Sigma$ , and their reciprocals  $\frac{1}{\sigma_1}, \dots, \frac{1}{\sigma_r}$  are on the diagonal of  $\Sigma^\dagger$ .

► **Example.** Find the pseudoinverse of  $A = [-1 \ 2 \ 2]$ .

The SVD of  $A$  is

$$A = [1][3 \ 0 \ 0] \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}.$$

Then the pseudoinverse of  $A$  is

$$A^\dagger = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} \frac{1}{3} \\ 0 \\ 0 \end{bmatrix} [1] = \begin{bmatrix} -\frac{1}{9} \\ \frac{2}{9} \\ \frac{2}{9} \end{bmatrix}.$$