

# Lecture 22: Complex Matrices (Sections 5.5)

---

Thang Huynh, UC San Diego

3/7/2018

## Complex Numbers and Their Conjugates

- **Complex conjugate** of  $a + ib$  is  $a - ib$  and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

## Complex Numbers and Their Conjugates

- **Complex conjugate** of  $a + ib$  is  $a - ib$  and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

- Conjugate of a product (or sum) equals the product (or sum) of the conjugates:

$$\overline{(a + ib)(c + id)} = (\overline{a + ib})(\overline{c + id})$$

$$\overline{a + ib + c + id} = \overline{(a + c) + i(b + d)}$$

## Complex Numbers and Their Conjugates

- **Complex conjugate** of  $a + ib$  is  $a - ib$  and is denoted by

$$(a + ib)^* = \overline{a + ib} = a - ib.$$

- Conjugate of a product (or sum) equals the product (or sum) of the conjugates:

$$\overline{(a + ib)(c + id)} = (\overline{a + ib})(\overline{c + id})$$

$$\overline{a + ib} + \overline{c + id} = \overline{(a + c) + i(b + d)}$$

- Multiplying  $a + ib$  by its conjugate produces  $a^2 + b^2$ :

$$(a + ib)(a - ib) = a^2 + b^2.$$

## Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

## Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

### ► Length/norm of $\mathbf{x}$

$$\|\mathbf{x}\| = \sqrt{|x_1|^2 + |x_2|^2 + \dots + |x_n|^2}.$$

## Lengths and Inner Products in the Complex Case

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{with components } x_j = a_j + ib_j.$$

### ► Length/norm of $\mathbf{x}$

$$\|\mathbf{x}\| = \sqrt{|x_1|^2 + |x_2|^2 + \dots + |x_n|^2}.$$

### ► Example.

$$\mathbf{x} = \begin{bmatrix} 1 \\ i \end{bmatrix} \Rightarrow \|\mathbf{x}\| = \sqrt{|1|^2 + |i|^2} = 2.$$

## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$



## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products  $\langle \mathbf{x}, \mathbf{y} \rangle$  and  $\langle \mathbf{y}, \mathbf{x} \rangle$  of  $\mathbf{x} = (1 + 3i, 3i)$  and  $\mathbf{y} = (6 + 3i, 4 + i)$ .

## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products  $\langle \mathbf{x}, \mathbf{y} \rangle$  and  $\langle \mathbf{y}, \mathbf{x} \rangle$  of  $\mathbf{x} = (1 + 3i, 3i)$  and  $\mathbf{y} = (6 + 3i, 4 + i)$ .

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products  $\langle \mathbf{x}, \mathbf{y} \rangle$  and  $\langle \mathbf{y}, \mathbf{x} \rangle$  of  $\mathbf{x} = (1 + 3i, 3i)$  and  $\mathbf{y} = (6 + 3i, 4 + i)$ .

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

$$\langle \mathbf{y}, \mathbf{x} \rangle = (6 - 3i)(1 + 3i) + (4 - i)(3i) = 18 + 27i.$$

## Lengths and Inner Products in the Complex Case

► Inner product of  $\mathbf{x}$  and  $\mathbf{y}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \bar{\mathbf{x}}^T \mathbf{y} = \bar{x}_1 y_1 + \cdots + \bar{x}_n y_n.$$

► Example.

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\bar{x}_1 x_1 + \cdots + \bar{x}_n x_n} = \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2}.$$

► Example. Find inner products  $\langle \mathbf{x}, \mathbf{y} \rangle$  and  $\langle \mathbf{y}, \mathbf{x} \rangle$  of  $\mathbf{x} = (1 + 3i, 3i)$  and  $\mathbf{y} = (6 + 3i, 4 + i)$ .

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{(1 + 3i)}(6 + 3i) + \overline{(3i)}(4 + i) = (1 - 3i)(6 + 3i) - 3i(4 + i) = 18 - 27i.$$

$$\langle \mathbf{y}, \mathbf{x} \rangle = (6 - 3i)(1 + 3i) + (4 - i)(3i) = 18 + 27i.$$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \mathbf{x} \rangle}$$

## Conjugate transposes of a matrix $A$ and a vector

► **Definition.** A conjugate transpose of a matrix  $A$  is  $\overline{A}^T$  (and denoted by  $A^H$  or  $A^*$ ).

## Conjugate transposes of a matrix $A$ and a vector

► **Definition.** A conjugate transpose of a matrix  $A$  is  $\overline{A}^T$  (and denoted by  $A^H$  or  $A^*$ ).

► **Example.**

$$\begin{bmatrix} 2+i & 3i \\ 4-i & 5 \\ 0 & 0 \end{bmatrix}^H = \begin{bmatrix} 2-i & 4+i & 0 \\ -3i & 5 & 0 \end{bmatrix}.$$

## Conjugate transposes of a matrix $A$ and a vector

► **Definition.** A conjugate transpose of a matrix  $A$  is  $\bar{A}^T$  (and denoted by  $A^H$  or  $A^*$ ).

► **Example.**

$$\begin{bmatrix} 2+i & 3i \\ 4-i & 5 \\ 0 & 0 \end{bmatrix}^H = \begin{bmatrix} 2-i & 4+i & 0 \\ -3i & 5 & 0 \end{bmatrix}.$$

► **Definition.** A conjugate transpose of a vector  $x$  is  $\bar{x}^T$  and denoted by  $x^H$ .



## Conjugate transposes of a matrix $A$ and a vector

► **Definition.** A conjugate transpose of a matrix  $A$  is  $\overline{A}^T$  (and denoted by  $A^H$  or  $A^*$ ).

► **Example.**

$$\begin{bmatrix} 2+i & 3i \\ 4-i & 5 \\ 0 & 0 \end{bmatrix}^H = \begin{bmatrix} 2-i & 4+i & 0 \\ -3i & 5 & 0 \end{bmatrix}.$$

► **Definition.** A conjugate transpose of a vector  $x$  is  $\overline{x}^T$  and denoted by  $x^H$ .

► **Example.** If  $x = \begin{bmatrix} 1+i \\ 2-i \\ 3+4i \end{bmatrix}$ , then

$$x^H = (1-i, 2+i, 3-4i).$$

## Conjugate transposes of a matrix $A$ and a vector

- $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^H \mathbf{y}$ .  
 $\mathbf{x}$  and  $\mathbf{y}$  are orthogonal if  $\mathbf{x}^H \mathbf{y} = 0$ .

## Conjugate transposes of a matrix $A$ and a vector

- $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^H \mathbf{y}$ .  
 $\mathbf{x}$  and  $\mathbf{y}$  are orthogonal if  $\mathbf{x}^H \mathbf{y} = 0$ .
- $\|\mathbf{x}\|^2 = \mathbf{x}^H \mathbf{x} = |x_1|^2 + \cdots + |x_n|^2$ .

## Conjugate transposes of a matrix $A$ and a vector

- $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^H \mathbf{y}$ .  
 $\mathbf{x}$  and  $\mathbf{y}$  are orthogonal if  $\mathbf{x}^H \mathbf{y} = 0$ .
- $\|\mathbf{x}\|^2 = \mathbf{x}^H \mathbf{x} = |x_1|^2 + \cdots + |x_n|^2$ .
- $(AB)^H = B^H A^H$ .

# Hermitian Matrices

► **Definition.**  $A$  is a Hermitian matrix if  $A = A^H$ .

# Hermitian Matrices

► **Definition.**  $A$  is a Hermitian matrix if  $A = A^H$ .

► **Example.**

$$A = \begin{bmatrix} 2 & 3 - 3i \\ 3 + 3i + 5 & \end{bmatrix} = A^H.$$

# Hermitian Matrices

► **Definition.**  $A$  is a Hermitian matrix if  $A = A^H$ .

► **Example.**

$$A = \begin{bmatrix} 2 & 3 - 3i \\ 3 + 3i + 5 & \end{bmatrix} = A^H.$$

The diagonal entries of a Hermitian matrix must be real. And

$$a_{ij} = \overline{a_{ji}}.$$

# Hermitian Matrices

► **Property.** If  $A = A^H$ , then for all complex vectors  $\mathbf{x}$ ,  $\mathbf{x}^H A \mathbf{x}$  is real.



# Hermitian Matrices

- ▶ **Property.** If  $A = A^H$ , then for all complex vectors  $\mathbf{x}$ ,  $\mathbf{x}^H A \mathbf{x}$  is real.
- ▶ **Property.** If  $A = A^H$ , every eigenvalue is real.

# Hermitian Matrices

- ▶ **Property.** If  $A = A^H$ , then for all complex vectors  $\mathbf{x}$ ,  $\mathbf{x}^H A \mathbf{x}$  is real.
- ▶ **Property.** If  $A = A^H$ , every eigenvalue is real.
- ▶ **Property.** Two eigenvectors of a real symmetric matrix or a Hermitian matrix, if they come from different eigenvalues, are orthogonal to one another.

## Hermitian Matrices

► Example.

$$(A - 8I)\mathbf{x} = \begin{bmatrix} -6 & 3 - i \\ 3 + 3i & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x} = \begin{bmatrix} 1 \\ 1 + i \end{bmatrix}$$

$$(A + I)\mathbf{y} = \begin{bmatrix} 3 & 3 - i \\ 3 + 3i & 6 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{y} = \begin{bmatrix} 1 - i \\ -1 \end{bmatrix}.$$

## Hermitian Matrices

► Example.

$$(A - 8I)\mathbf{x} = \begin{bmatrix} -6 & 3 - i \\ 3 + 3i & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x} = \begin{bmatrix} 1 \\ 1 + i \end{bmatrix}$$
$$(A + I)\mathbf{y} = \begin{bmatrix} 3 & 3 - i \\ 3 + 3i & 6 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{y} = \begin{bmatrix} 1 - i \\ -1 \end{bmatrix}.$$

These two eigenvectors are orthogonal

$$\mathbf{x}^H \mathbf{y} = [1 \ 1 - i] \begin{bmatrix} 1 - i \\ -1 \end{bmatrix} = 0.$$

## Hermitian Matrices

► **Example.**

$$(A - 8I)\mathbf{x} = \begin{bmatrix} -6 & 3 - i \\ 3 + 3i & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{x} = \begin{bmatrix} 1 \\ 1 + i \end{bmatrix}$$
$$(A + I)\mathbf{y} = \begin{bmatrix} 3 & 3 - i \\ 3 + 3i & 6 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \mathbf{y} = \begin{bmatrix} 1 - i \\ -1 \end{bmatrix}.$$

These two eigenvectors are orthogonal

$$\mathbf{x}^H \mathbf{y} = [1 \ 1 - i] \begin{bmatrix} 1 - i \\ -1 \end{bmatrix} = 0.$$

► **Property.** (Spectral Theorem) A real symmetric matrix can be factored into  $A = Q\Lambda Q^T$ . Its orthonormal eigenvectors are in the orthogonal matrix  $Q$  and its eigenvalues are in  $\Lambda$ .

# Unitary Matrices

► **Definition.** A complex matrix  $U$  with orthonormal columns is called a unitary matrix, i.e.,

$$U^H U = I \quad U U^H = I, \quad \text{or } U^H = U^{-1}.$$

# Unitary Matrices

► **Definition.** A complex matrix  $U$  with orthonormal columns is called a unitary matrix, i.e.,

$$U^H U = I \quad U U^H = I, \quad \text{or } U^H = U^{-1}.$$

► **Property.**  $(U\mathbf{x})^H (U\mathbf{y}) = \mathbf{x}^H U^H U \mathbf{y} = \mathbf{x}^H \mathbf{y}$  and lengths are preserved by  $U$ :

$$\|U\mathbf{x}\| = \|\mathbf{x}\|.$$

# Unitary Matrices

► **Definition.** A complex matrix  $U$  with orthonormal columns is called a unitary matrix, i.e.,

$$U^H U = I \quad U U^H = I, \quad \text{or } U^H = U^{-1}.$$

► **Property.**  $(U\mathbf{x})^H (U\mathbf{y}) = \mathbf{x}^H U^H U \mathbf{y} = \mathbf{x}^H \mathbf{y}$  and lengths are preserved by  $U$ :

$$\|U\mathbf{x}\| = \|\mathbf{x}\|.$$

► **Property.** Every eigenvalue of  $U$  has absolute value  $|\lambda| = 1$ .



# Unitary Matrices

► **Definition.** A complex matrix  $U$  with orthonormal columns is called a unitary matrix, i.e.,

$$U^H U = I \quad U U^H = I, \quad \text{or } U^H = U^{-1}.$$

► **Property.**  $(U\mathbf{x})^H (U\mathbf{y}) = \mathbf{x}^H U^H U \mathbf{y} = \mathbf{x}^H \mathbf{y}$  and lengths are preserved by  $U$ :

$$\|U\mathbf{x}\| = \|\mathbf{x}\|.$$

► **Property.** Every eigenvalue of  $U$  has absolute value  $|\lambda| = 1$ .

► **Property.** Eigenvectors corresponding to different eigenvalues are orthonormal.

## Unitary Matrices

► **Example.**  $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$  has eigenvalues  $e^{it}$  and  $e^{-it}$ .

## Unitary Matrices

► **Example.**  $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$  has eigenvalues  $e^{it}$  and  $e^{-it}$ .

Orthogonal eigenvectors are  $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$  and  $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$ .

## Unitary Matrices

► **Example.**  $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$  has eigenvalues  $e^{it}$  and  $e^{-it}$ .

Orthogonal eigenvectors are  $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$  and  $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$ .

► **Example.** (normalized) Fourier matrix

$$U = \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & w & \dots & w^{n-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & w^{n-1} & \dots & w^{(n-1)^2} \end{bmatrix}$$

where  $w = e^{2\pi i/n}$ .

## Unitary Matrices

► **Example.**  $U = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$  has eigenvalues  $e^{it}$  and  $e^{-it}$ .

Orthogonal eigenvectors are  $\mathbf{x} = \begin{bmatrix} 1 \\ -i \end{bmatrix}$  and  $\mathbf{y} = \begin{bmatrix} 1 \\ i \end{bmatrix}$ .

► **Example.** (normalized) Fourier matrix

$$U = \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & w & \dots & w^{n-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & w^{n-1} & \dots & w^{(n-1)^2} \end{bmatrix}$$

where  $w = e^{2\pi i/n}$ .

Row  $i$  of  $U^H$  times column  $j$  of  $U$  is

$$\frac{1}{n}(1 + W + W^2 + \dots + W^{n-1}) = \frac{W^n - 1}{W - 1} = 0,$$

where  $W = w^{j-i}$ .