

Lecture 8: Four Fundamental Subspaces and Linear Transformations (Section 2.4 and 2.6)

Thang Huynh, UC San Diego

1/26/2018

The four fundamental subspaces

► **Definition.** The **row space** of A is the column space of A^T .

The four fundamental subspaces

- ▶ **Definition.** The **row space** of A is the column space of A^T .
- ▶ **Definition.** The **left null space** of A is the null space of A^T .

Why "left"?

The four fundamental subspaces

- ▶ **Definition.** The **row space** of A is the column space of A^T .
 - ▶ **Definition.** The **left null space** of A is the null space of A^T .
- Why "left"? \mathbf{y} is in $N(A^T)$ if and only if $\mathbf{y}^T A = 0$.

The four fundamental subspaces

► **Definition.** The **row space** of A is the column space of A^T .

► **Definition.** The **left null space** of A is the null space of A^T .

Why "left"? \mathbf{y} is in $N(A^T)$ if and only if $\mathbf{y}^T A = 0$.

► **Definition.** The **rank** of a matrix A is the number of its pivots.

The four fundamental subspaces

► **Example.** Find a basis for $C(A)$ and $C(A^T)$ where

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

The four fundamental subspaces

► **Example.** Find a basis for $C(A)$ and $C(A^T)$ where

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

► **Solution.** To find $C(A)$, we can just use the echelon form of A . Likewise, we can also obtain $C(A^T)$ for an echelon form of A^T . But, it's not necessary!

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B.$$

The rank of A is 2. And $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$ form a basis for $C(A)$.

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B.$$

The rank of A is 2. And $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$ form a basis for $C(A)$.

Recall that $C(A) \neq C(B)$. (We performed row operations).
However, $C(A^T) = C(B^T)$.

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B.$$

The rank of A is 2. And $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$ form a basis for $C(A)$.

Recall that $C(A) \neq C(B)$. (We performed row operations).

However, $C(A^T) = C(B^T)$.

Row space is preserved by elementary row operations.

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B.$$

The rank of A is 2. And $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$ form a basis for $C(A)$.

Recall that $C(A) \neq C(B)$. (We performed row operations).

However, $C(A^T) = C(B^T)$.

Row space is preserved by elementary row operations.

Hence, $\left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -1 \\ -5 \end{bmatrix} \right\}$ form a basis for $C(A^T)$.

The four fundamental subspaces

Theorem. (Fundamental Theorem of Linear Algebra, Part I)

Let A be an $m \times n$ matrix of rank r .

- $\dim C(A) = r$
- $\dim C(A^T) = r$
- $\dim N(A) = n - r$
- $\dim N(A^T) = m - r$

The four fundamental subspaces

The column and row space always have the same dimension!

That is, A and A^T have the same rank.

► **Example.**

$$\begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$$

The four fundamental subspaces

The column and row space always have the same dimension!

That is, A and A^T have the same rank.

► **Example.**

$$\begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$$

- $\dim C(A) = 3$

The four fundamental subspaces

The column and row space always have the same dimension!

That is, A and A^T have the same rank.

► **Example.**

$$\begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$$

- $\dim C(A) = 3$
- $\dim C(A^T) = 3$

The four fundamental subspaces

The column and row space always have the same dimension!

That is, A and A^T have the same rank.

► **Example.**

$$\begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$$

- $\dim C(A) = 3$
- $\dim C(A^T) = 3$
- $\dim N(A) = 1$

The four fundamental subspaces

The column and row space always have the same dimension!

That is, A and A^T have the same rank.

► **Example.**

$$\begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$$

- $\dim C(A) = 3$
- $\dim C(A^T) = 3$
- $\dim N(A) = 1$
- $\dim N(A^T) = 0$

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .
- c) A has rank n .

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .
- c) A has rank n .
- d) The columns of A span \mathbb{R}^n .

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .
- c) A has rank n .
- d) The columns of A span \mathbb{R}^n .
- e) The columns of A are independent.

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .
- c) A has rank n .
- d) The columns of A span \mathbb{R}^n .
- e) The columns of A are independent.
- f) For every \mathbf{b} , the system $A\mathbf{x} = \mathbf{b}$ has a unique solution.

Existence of inverses

Let A be an $n \times n$ matrix. Then the following statements are equivalent:

- a) A is invertible.
- b) A is row equivalent to I_n .
- c) A has rank n .
- d) The columns of A span \mathbb{R}^n .
- e) The columns of A are independent.
- f) For every \mathbf{b} , the system $A\mathbf{x} = \mathbf{b}$ has a unique solution.

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{v}$?

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{0}$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{0\}$.

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{0}$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{0\}$. True!

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = 0$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{0\}$. True!
- An $n \times n$ matrix A is invertible if and only if the rows of A span \mathbb{R}^n .

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = 0$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{0\}$. True!
- An $n \times n$ matrix A is invertible if and only if the rows of A span \mathbb{R}^n . True!

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = \mathbf{0}$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{\mathbf{0}\}$. True!
- An $n \times n$ matrix A is invertible if and only if the rows of A span \mathbb{R}^n . True!
- An $n \times n$ matrix A is invertible if and only if the rows of A are independent.

► **Example.** Suppose A is a 5×5 matrix, and that \mathbf{v} is a vector in \mathbb{R}^5 which is not a linear combination of the columns of A . What can you say about the number of solutions to $A\mathbf{x} = 0$.

► **Example.** True or false?

- An $n \times n$ matrix A is invertible if and only if $N(A) = \{0\}$. True!
- An $n \times n$ matrix A is invertible if and only if the rows of A span \mathbb{R}^n . True!
- An $n \times n$ matrix A is invertible if and only if the rows of A are independent. True!

Linear transformations

Consider vector spaces V and W .

► **Definition.** A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \text{ in } V \text{ and all } c, d \text{ in } \mathbb{R}.$$

Linear transformations

Consider vector spaces V and W .

► **Definition.** A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \text{ in } V \text{ and all } c, d \text{ in } \mathbb{R}.$$

► **Example.** Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation from \mathbb{R}^n to \mathbb{R}^m . Why?

Linear transformations

Consider vector spaces V and W .

► **Definition.** A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \text{ in } V \text{ and all } c, d \text{ in } \mathbb{R}.$$

► **Example.** Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation from \mathbb{R}^n to \mathbb{R}^m . Why?

► **Example.** Let \mathcal{P}_n be the vector space of all polynomials of degree at most n . Consider the map $T : \mathcal{P}_n \rightarrow \mathcal{P}_{n-1}$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

This map is a linear transformation! Why?

Representing linear maps by matrices

Let $\mathbf{x}_1, \dots, \mathbf{x}_n$ be a basis for V . A linear map $T : V \rightarrow W$ is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$.

Representing linear maps by matrices

Let $\mathbf{x}_1, \dots, \mathbf{x}_n$ be a basis for V . A linear map $T : V \rightarrow W$ is determined by the values $T(\mathbf{x}_1), \dots, T(\mathbf{x}_n)$.

► **Definition.** (From linear maps to matrices)

Let $\mathbf{x}_1, \dots, \mathbf{x}_n$ be a basis for V , and $\mathbf{y}_1, \dots, \mathbf{y}_m$ a basis for W .

The **matrix representing** T with respect to these bases

- has n columns (one for each of the \mathbf{x}_j),
- the j -th column has m entries a_{1j}, \dots, a_{mj} determined by

$$T(\mathbf{x}_j) = a_{1j}\mathbf{y}_1 + \dots + a_{mj}\mathbf{y}_m.$$

Representing linear maps by matrices

► **Example.** Let $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$. Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

What is the matrix A representing T with respect to the standard bases?

► **Example.** Let $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$. Let T be the linear map such that

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 0 \\ 7 \end{bmatrix}.$$

What is the matrix B representing T with respect to the following bases?

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix} \text{ for } \mathbb{R}^2, \quad \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ for } \mathbb{R}^3.$$

► **Example.** Let $T : \mathcal{P}_3 \rightarrow \mathcal{P}_2$ be the linear map given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

What is the matrix A representing T with respect to the standard bases?