



Algorithm Design XI

Linear Programming I

Guoqiang Li
School of Software



SHANGHAI JIAO TONG
UNIVERSITY

An Introduction to Linear Programming



A linear programming problem gives a set of variables, and assigns real values to them so as to



A **linear programming problem** gives a set of **variables**, and assigns **real values** to them so as to

- 1 satisfy a set of **linear equations** and/or **linear inequalities** involving these variables, and
- 2 maximize or minimize a given **linear objective function**.

Example: Profit Maximization



A boutique chocolatier has two products:

Example: Profit Maximization



A boutique chocolatier has two products:

- triangular chocolates, called **Pyramide**,
- and the more decadent and deluxe **Pyramide Nuit**.

Example: Profit Maximization



A boutique chocolatier has two products:

- triangular chocolates, called **Pyramide**,
- and the more decadent and deluxe **Pyramide Nuit**.

Q: How much of each should it produce to **maximize** profits?

Example: Profit Maximization

A boutique chocolatier has two products:

- triangular chocolates, called **Pyramide**,
- and the more decadent and deluxe **Pyramide Nuit**.

Q: How much of each should it produce to **maximize** profits?

- Every box of **Pyramide** has a profit of \$1.
- Every box of **Nuit** has a profit of \$6.
- The daily demand is limited to at most 200 boxes of **Pyramide** and 300 boxes of **Nuit**.
- The current workforce can produce a total of at most 400 boxes of chocolate per day.

LP Formulation



Objective function $\max x_1 + 6x_2$

Constraints $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 \leq 400$

$x_1, x_2 \geq 0$

LP Formulation



Objective function $\max x_1 + 6x_2$

Constraints $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 \leq 400$

$x_1, x_2 \geq 0$

A linear equation in x_1 and x_2 defines a line in the two-dimensional (2D) plane, and a linear inequality designates a half-space, the region on one side of the line.

LP Formulation



Objective function $\max x_1 + 6x_2$

Constraints $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 \leq 400$

$x_1, x_2 \geq 0$

A linear equation in x_1 and x_2 defines a line in the two-dimensional (2D) plane, and a linear inequality designates a half-space, the region on one side of the line.

The set of all feasible solutions of this linear program is the intersection of five half-spaces.

LP Formulation



Objective function $\max x_1 + 6x_2$

Constraints $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 \leq 400$

$x_1, x_2 \geq 0$

A linear equation in x_1 and x_2 defines a line in the two-dimensional (2D) plane, and a linear inequality designates a half-space, the region on one side of the line.

The set of all feasible solutions of this linear program is the intersection of five half-spaces.


It is a convex polygon.

凸边界

The Convex Polygon



SHANGHAI JIAO TONG
UNIVERSITY



The Optimal Solution



We want to find the point in this polygon at which the objective function is **maximized**.

The Optimal Solution



We want to find the point in this polygon at which the objective function is **maximized**.

The points with a profit of c dollars lie on the line $x_1 + 6x_2 = c$, which has a **slope** of $-1/6$.

The Optimal Solution



We want to find the point in this polygon at which the objective function is **maximized**.

The points with a profit of c dollars lie on the line $x_1 + 6x_2 = c$, which has a **slope** of $-1/6$.

As c increases, this “**profit line**” moves parallel to itself, up and to the right.

The Optimal Solution



We want to find the point in this polygon at which the objective function is **maximized**.

The points with a profit of c dollars lie on the line $x_1 + 6x_2 = c$, which has a **slope** of $-1/6$.

As c increases, this “**profit line**” moves parallel to itself, up and to the right.

Since the goal is to **maximize** c , we must move the line as far up as possible, while still touching the **feasible region**.

The Optimal Solution



We want to find the point in this polygon at which the objective function is **maximized**.


The points with a profit of c dollars lie on the line $x_1 + 6x_2 = c$, which has a **slope** of $-1/6$.

As c increases, this “**profit line**” moves **parallel to itself, up and to the right**.

Since the goal is to **maximize** c , we must move the line as far up as possible, **while still touching the feasible region**.

The optimum solution will be the very last feasible point that the profit line sees and must **therefore be a vertex of the polygon**.

The Convex Polygon



The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The only exceptions are cases in which there is **no optimum**; this can happen in two ways:

The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The only exceptions are cases in which there is **no optimum**; this can happen in two ways:

- 1 The linear program is **infeasible**; that is, the constraints are so tight that it is impossible to satisfy all of them.

The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The only exceptions are cases in which there is **no optimum**; this can happen in two ways:

- 1 The linear program is **infeasible**; that is, the constraints are so tight that it is impossible to satisfy all of them.
 - For instance, $x \leq 1$, $x \geq 2$.

The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The only exceptions are cases in which there is **no optimum**; this can happen in two ways:

- 1 The linear program is **infeasible**; that is, the constraints are so tight that it is impossible to satisfy all of them.
 - For instance, $x \leq 1$, $x \geq 2$.
- 2 The constraints are so loose that the feasible region is **unbounded**, and it is possible to achieve arbitrarily high objective values.

The Optimal Solution



It is a general rule of linear programs that the optimum is achieved at a vertex of the feasible region.

The only exceptions are cases in which there is no optimum; this can happen in two ways:

- 1 The linear program is infeasible; that is, the constraints are so tight that it is impossible to satisfy all of them.
 - For instance, $x \leq 1, x \geq 2$.
- 2 The constraints are so loose that the feasible region is unbounded, and it is possible to achieve arbitrarily high objective values.
 - For instance, $\max x_1 + x_2$
 - $x_1, x_2 \geq 0$

一般都在顶点处取到，但是有特例，如果约束条件没有解或者不会构成一个边界那么就不会在顶点处取到最优解。

Solving Linear Programs



SHANGHAI JIAO TONG
UNIVERSITY

Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

Solving Linear Programs



SHANGHAI JIAO TONG
UNIVERSITY

Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

This algorithm starts at a **vertex**,

Solving Linear Programs

Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

This algorithm starts at a **vertex**, and repeatedly looks for an **adjacent vertex** of better objective value.

Solving Linear Programs



Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

This algorithm starts at a **vertex**, and repeatedly looks for an **adjacent vertex** of better objective value.

It does **hill-climbing** on the vertices of the polygon,

Solving Linear Programs

Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

This algorithm starts at a **vertex**, and repeatedly looks for an **adjacent vertex** of better objective value.

It does **hill-climbing** on the vertices of the polygon, walking from neighbor to neighbor so as to **steadily increase profit** along the way.

Solving Linear Programs



(单纯形法)

Linear programs (LPs) can be solved by the **simplex method**, devised by **George Dantzig** in 1947.

This algorithm starts at a **vertex**, and repeatedly looks for an **adjacent vertex** of better objective value.

It does **hill-climbing** on the vertices of the polygon, walking **from neighbor to neighbor so as to steadily increase profit along the way.**

Upon reaching a vertex that has no better neighbor, simplex declares it to be optimal and halts.

Solving Linear Programs



SHANGHAI JIAO TONG
UNIVERSITY

Q: Why does this local test imply **global optimality**?

Solving Linear Programs




SHANGHAI JIAO TONG
UNIVERSITY

Q: Why does this local test imply **global optimality**?

By simple **geometry**. Since all the vertex's neighbors lie below the line, the rest of the **feasible polygon** must also lie below this line.

The Example



More Products



SHANGHAI JIAO TONG
UNIVERSITY

The chocolatier introduces a third and even more exclusive chocolates, called **Pyramide Luxe**. One box of these will bring in a profit of \$13.

More Products



The chocolatier introduces a third and even more exclusive chocolates, called **Pyramide Luxe**. One box of these will bring in a profit of \$13.

Let x_1, x_2, x_3 denote the number of boxes of each chocolate produced daily, with x_3 referring to **Luxe**.



The chocolatier introduces a third and even more exclusive chocolates, called **Pyramide Luxe**. One box of these will bring in a profit of \$13.

Let x_1, x_2, x_3 denote the number of boxes of each chocolate produced daily, with x_3 referring to **Luxe**.

The old constraints on x_1 and x_2 persist. The labor restriction now extends to x_3 as well: the sum of all three variables is at most 400.



The chocolatier introduces a third and even more exclusive chocolates, called **Pyramide Luxe**. One box of these will bring in a profit of \$13.

Let x_1, x_2, x_3 denote the number of boxes of each chocolate produced daily, with x_3 referring to **Luxe**.

The old constraints on x_1 and x_2 persist. The labor restriction now extends to x_3 as well: the sum of all three variables is at most 400.

Nuit and **Luxe** require the same packaging machinery. Luxe uses it **three times** as much, which imposes another constraint $x_2 + 3x_3 \leq 600$.



$$\max x_1 + 6x_2 + 13x_3$$

$$x_1 \leq 200$$

$$x_2 \leq 300$$

$$x_1 + x_2 + x_3 \leq 400$$

$$x_2 + 3x_3 \leq 600$$

$$x_1, x_2, x_3 \geq 0$$



The space of solutions is now **three-dimensional**.



The space of solutions is now **three-dimensional**.

Each **linear equation** defines a **3D plane**, and **each inequality** a **half-space** on one side of the plane.



The space of solutions is now **three-dimensional**.

Each **linear equation** defines a **3D plane**, and **each inequality** a **half-space** on one side of the plane.

The feasible region is an intersection of seven half-spaces, a **polyhedron**.



The space of solutions is now **three-dimensional**.

Each **linear equation** defines a **3D plane**, and **each inequality** a **half-space** on one side of the plane.

The feasible region is an intersection of seven half-spaces, a **polyhedron**.

A profit of c corresponds to the plane $x_1 + 6x_2 + 13x_3 = c$.



The space of solutions is now **three-dimensional**.


Each **linear equation** defines a **3D plane**, and **each inequality** a **half-space** on one side of the plane.

The feasible region is an intersection of seven half-spaces, a **polyhedron**.

A profit of c corresponds to the plane $x_1 + 6x_2 + 13x_3 = c$.

As c increases, this profit-plane moves parallel to itself, further into the positive **orthant** until it no longer touches the feasible region.

The Example





The point of final contact is the **optimal vertex**: $(0, 300, 100)$, with total **profit** \$3100.



The point of final contact is the **optimal vertex**: $(0, 300, 100)$, with total **profit** \$3100.

Q: How would the **simplex** algorithm behave on this modified problem?




The point of final contact is the **optimal vertex**: $(0, 300, 100)$, with total **profit** \$3100.

Q: How would the **simplex** algorithm behave on this modified problem?

A possible **trajectory**

$$\frac{(0, 0, 0)}{\$0} \rightarrow \frac{(200, 0, 0)}{\$200} \rightarrow \frac{(200, 200, 0)}{\$1400} \rightarrow \frac{(200, 0, 200)}{\$2800} \rightarrow \frac{(0, 300, 100)}{\$3100}$$

The Example



Integer Linear Programming and Rounding

Example: Production Planning

The company makes **handwoven carpets**, a product for which the demand is extremely seasonal.

Example: Production Planning

The company makes **handwoven carpets**, a product for which the demand is extremely seasonal.

Our analyst has just obtained demand estimates for all months of the next calendar year:

d_1, d_2, \dots, d_{12} , ranging from 440 to 920.

Example: Production Planning

The company makes **handwoven carpets**, a product for which the demand is extremely seasonal.

Our analyst has just obtained demand estimates for all months of the next calendar year:

d_1, d_2, \dots, d_{12} , ranging from 440 to 920.

Currently with 30 employees, each of whom makes 20 carpets per month and gets a monthly salary of \$2000.

Example: Production Planning



The company makes **handwoven carpets**, a product for which the demand is extremely seasonal.

Our analyst has just obtained demand estimates **for all months of the next calendar year:**
 d_1, d_2, \dots, d_{12} , ranging from 440 to 920.

Currently with **30** employees, each of whom makes **20** carpets per month and gets a monthly salary of **\$2000**.

With no initial surplus of carpets.

Example: Production Planning



Q: How can we handle the **fluctuations in demand**? There are three ways:

Example: Production Planning



Q: How can we handle the **fluctuations in demand**? There are three ways:

- 1 **Overtime**. Overtime pay is **80%** more than regular pay. Workers can put in at most **30%** overtime.

Example: Production Planning



Q: How can we handle the **fluctuations in demand**? There are three ways:

- 1 **Overtime**. Overtime pay is **80%** more than regular pay. Workers can put in at most **30%** overtime.
- 2 **Hiring and firing**, costing **\$320** and **\$400**, respectively, per worker.

Example: Production Planning



Q: How can we handle the **fluctuations in demand**? There are three ways:

- 1 **Overtime**. Overtime pay is **80%** more than regular pay. Workers can put in at most **30%** overtime.
- 2 **Hiring and firing**, costing **\$320** and **\$400**, respectively, per worker.
- 3 **Storing surplus production**, costing **\$8** per carpet per month. Currently without stored carpets on hand, and without any carpets stored at the end of year.



- w_i = number of workers during i -th month; $w_0 = 30$.
- x_i = number of carpets made during i -th month.
- o_i = number of carpets made by overtime in month i .
- h_i, f_i = number of workers hired and fired, respectively, at beginning of month i .
- s_i = number of carpets stored at end of month i ; $s_0 = 0$.

LP Formulation



SHANGHAI JIAO TONG
UNIVERSITY

LP Formulation



All variables must be **nonnegative**:

$$w_i, x_i, o_i, h_i, f_i, s_i \geq 0, i = 1, \dots, 12$$

LP Formulation



All variables must be **nonnegative**:

$$w_i, x_i, o_i, h_i, f_i, s_i \geq 0, i = 1, \dots, 12$$

The total number of carpets made per month consists of regular production plus overtime:

$$x_i = 20w_i + o_i$$

$$i = 1, \dots, 12.$$



All variables must be **nonnegative**:

$$w_i, x_i, o_i, h_i, f_i, s_i \geq 0, i = 1, \dots, 12$$

The total number of carpets made per month consists of regular production plus overtime:

$$x_i = 20w_i + o_i$$

$$i = 1, \dots, 12.$$

The number of workers can potentially change at the start of each month:

$$w_i = w_{i-1} + h_i - f_i$$

LP Formulation



SHANGHAI JIAO TONG
UNIVERSITY



The number of carpets stored at the end of each month is what we started with, plus the number we made, minus the demand for the month:

$$s_i = s_{i-1} + x_i - d_i$$



The number of carpets stored at the end of each month is what we started with, plus the number we made, minus the demand for the month:

$$s_i = s_{i-1} + x_i - d_i$$

And overtime is limited:

$$o_i \leq 6w_i$$



The **objective function** is to minimize the total cost:

$$\min 2000 \sum_i w_i + 320 \sum_i h_i + 400 \sum_i f_i + 8 \sum_i s_i + 180 \sum_i o_i$$

180是怎么来的？

一天正常工时为200/3元，则加班时一天的工资为120元，平均每一条毛毯生产时间为2/3天，所以生产 o_i 条加班毛毯需要花费： $120 * o_i / (2/3) = 180 * o_i$ 。

Integer Linear Programming



SHANGHAI JIAO TONG
UNIVERSITY

The optimum solution might turn out to be **fractional**;



The optimum solution might turn out to be **fractional**; for instance, it might involve hiring **10.6** workers in the month of March.



The optimum solution might turn out to be **fractional**; for instance, it might involve hiring **10.6** workers in the month of March.

This number would have to be **rounded** to either **10** or **11** in order to make sense, and the overall cost would then increase correspondingly.

The optimum solution might turn out to be **fractional**; for instance, it might involve hiring **10.6** workers in the month of March.

This number would have to be rounded to either 10 or 11 in order to make sense, and the overall cost would then increase correspondingly.

In the example, most of the variables take on fairly large values, and thus **rounding** is unlikely to affect things too much.

Integer Linear Programming



SHANGHAI JIAO TONG
UNIVERSITY

There are other **LPs**, in which rounding decisions have to be made very carefully to end up with an integer solution of reasonable quality.

Integer Linear Programming



There are other **LPs**, in which rounding decisions have to be made very carefully to end up with an integer solution of reasonable quality.

There is a tension in linear programming between the ease of **obtaining fractional solutions** and the **desirability of integer ones**.

Integer Linear Programming



There are other LPs, in which rounding decisions have to be made very carefully to end up with an integer solution of reasonable quality.

There is a tension in linear programming between the ease of obtaining fractional solutions and the desirability of integer ones.

In NP problems, finding the optimum integer solution of an LP is an important but very hard problem, called integer linear programming.

Product Planning Revisit



SHANGHAI JIAO TONG
UNIVERSITY

Recall:

$$\begin{aligned}\max \quad & x_1 + 6x_2 \\ & x_1 \leq 200 \\ & x_2 \leq 300 \\ & x_1 + x_2 \leq 400 \\ & x_1, x_2 \geq 0\end{aligned}$$

Product Planning Revisit



SHANGHAI JIAO TONG
UNIVERSITY

Recall:

$$\begin{aligned}\max \quad & x_1 + 6x_2 \\ & x_1 \leq 200 \\ & x_2 \leq 300 \\ & x_1 + x_2 \leq 400 \\ & x_1, x_2 \geq 0\end{aligned}$$

Simplex declares the optimum solution to be $(x_1, x_2) = (100, 300)$, with objective value 1900.

Can this answer be checked somehow?



Recall:

$$\begin{aligned}\max \quad & x_1 + 6x_2 \\ & x_1 \leq 200 \\ & x_2 \leq 300 \\ & x_1 + x_2 \leq 400 \\ & x_1, x_2 \geq 0\end{aligned}$$

Simplex declares the optimum solution to be $(x_1, x_2) = (100, 300)$, with objective value 1900.

Can this answer be checked somehow?

We take the first inequality and add it to six times the second inequality:

$$x_1 + 6x_2 \leq 2000$$



Multiplying the three inequalities by 0, 5, and 1, respectively, and adding them up yields

$$x_1 + 6x_2 \leq 1900$$

Multipliers



Let's investigate the issue by describing what we expect of these three multipliers, call them y_1 , y_2 , y_3 .

| Multiplier | Inequality | | | |
|------------|------------|-------|--------|------------|
| y_1 | x_1 | | \leq | 200 |
| y_2 | | x_2 | \leq | 300 |
| y_3 | x_1 | + | x_2 | \leq 400 |

Multipliers



Let's investigate the issue by describing what we expect of these three multipliers, call them y_1 , y_2 , y_3 .

| Multiplier | Inequality |
|------------|----------------------|
| y_1 | $x_1 \leq 200$ |
| y_2 | $x_2 \leq 300$ |
| y_3 | $x_1 + x_2 \leq 400$ |

These y_i 's must be **nonnegative**,

Multipliers



Let's investigate the issue by describing what we expect of these three multipliers, call them y_1 , y_2 , y_3 .

| Multiplier | Inequality |
|------------|----------------------|
| y_1 | $x_1 \leq 200$ |
| y_2 | $x_2 \leq 300$ |
| y_3 | $x_1 + x_2 \leq 400$ |

These y_i 's must be **nonnegative**, otherwise they are unqualified to multiply inequalities.

Multipliers



Let's investigate the issue by describing what we expect of these three multipliers, call them y_1 , y_2 , y_3 .

| Multiplier | Inequality |
|------------|----------------------|
| y_1 | $x_1 \leq 200$ |
| y_2 | $x_2 \leq 300$ |
| y_3 | $x_1 + x_2 \leq 400$ |

These y_i 's must be **nonnegative**, otherwise they are unqualified to multiply inequalities.

After the multiplication and addition steps, we get the bound:

$$(y_1 + y_3)x_1 + (y_2 + y_3)x_2 \leq 200y_1 + 300y_2 + 400y_3$$

Multipliers

Let's investigate the issue by describing what we expect of these three multipliers, call them y_1 , y_2 , y_3 .

| Multiplier | Inequality |
|------------|----------------------|
| y_1 | $x_1 \leq 200$ |
| y_2 | $x_2 \leq 300$ |
| y_3 | $x_1 + x_2 \leq 400$ |

These y_i 's must be **nonnegative**, otherwise they are unqualified to multiply inequalities.

After the multiplication and addition steps, we get the bound:

$$(y_1 + y_3)x_1 + (y_2 + y_3)x_2 \leq 200y_1 + 300y_2 + 400y_3$$

We want the left-hand side to look like the **objective function** $x_1 + 6x_2$ so that the right-hand side is an upper bound on the **optimum solution**.



$$x_1 + 6x_2 \leq 200y_1 + 300y_2 + 400y_3$$

if

$$y_1, y_2, y_3 \geq 0$$

$$y_1 + y_3 \geq 1$$

$$y_2 + y_3 \geq 6$$

The Dual Program



SHANGHAI JIAO TONG
UNIVERSITY

The Dual Program



We can easily find y 's that satisfy the inequalities on the right by simply making them large enough, for example $(y_1, y_2, y_3) = (5, 3, 6)$.

The Dual Program



We can easily find y 's that satisfy the inequalities on the right by simply making them large enough, for example $(y_1, y_2, y_3) = (5, 3, 6)$.

These particular multipliers tell us that the optimum solution of the LP is at most

$$200 \cdot 5 + 300 \cdot 3 + 400 \cdot 6 = 4300$$

The Dual Program



We can easily find y 's that satisfy the inequalities on the right by simply making them large enough, for example $(y_1, y_2, y_3) = (5, 3, 6)$.

These particular multipliers tell us that the optimum solution of the LP is at most

$$200 \cdot 5 + 300 \cdot 3 + 400 \cdot 6 = 4300$$

What we want is a bound as tight as possible, so we minimize

$$200y_1 + 300y_2 + 400y_3$$

subject to the preceding inequalities. This is a new linear program!

The Dual Program



$$\min 200y_1 + 300y_2 + 400y_3$$

$$y_1 + y_3 \geq 1$$

$$y_2 + y_3 \geq 6$$

$$y_1, y_2, y_3 \geq 0$$

The Dual Program



$$\begin{aligned}\min \quad & 200y_1 + 300y_2 + 400y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + y_3 \geq 6 \\ & y_1, y_2, y_3 \geq 0\end{aligned}$$

Any feasible value of this dual LP is an upper bound on the original primal LP.

The Dual Program



$$\begin{aligned}\min \quad & 200y_1 + 300y_2 + 400y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + y_3 \geq 6 \\ & y_1, y_2, y_3 \geq 0\end{aligned}$$

Any feasible value of this **dual LP** is an **upper bound** on the **original primal LP**.

If we find a pair of primal and dual feasible values that are equal, then they must both be **optimal**.

The Dual Program



$$\begin{aligned}\min & 200y_1 + 300y_2 + 400y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + y_3 \geq 6 \\ & y_1, y_2, y_3 \geq 0\end{aligned}$$

Any feasible value of this **dual LP** is an **upper bound** on the **original primal LP**.

If we find a pair of primal and dual feasible values that are equal, then they must both be **optimal**.

Here is just such a pair:

- **Primal:** $(x_1, x_2) = (100, 300)$;
- **Dual:** $(y_1, y_2, y_3) = (0, 5, 1)$.

The Dual Program



$$\begin{aligned}\min \quad & 200y_1 + 300y_2 + 400y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + y_3 \geq 6 \\ & y_1, y_2, y_3 \geq 0\end{aligned}$$

Any feasible value of this **dual LP** is an **upper bound** on the **original primal LP**.

If we find a pair of primal and dual feasible values that are equal, then they must both be **optimal**.

Here is just such a pair:

- **Primal:** $(x_1, x_2) = (100, 300)$;
- **Dual:** $(y_1, y_2, y_3) = (0, 5, 1)$.

They both have value 1900 and certify each other's optimality.



Matrix-Vector Form and Its Dual

关系：原问题与对偶问题的最值类型相反；原问题每有一个约束条件，对偶问题就有一个优化变量；原问题有一个优化变量，对偶问题就有一个约束条件。

Primal LP

$$\begin{aligned} \max \quad & c^T x \\ & Ax \leq b \\ & x \geq 0 \end{aligned}$$

Dual LP

$$\begin{aligned} \min \quad & y^T b \\ & y^T A \geq c^T \\ & y \geq 0 \end{aligned}$$

Primal LP:

$$\begin{aligned} \max \quad & c_1 x_1 + \cdots + c_n x_n \\ & a_{i1} x_1 + \cdots + a_{in} x_n \leq b_i \quad \text{for } i \in I \\ & a_{i1} x_1 + \cdots + a_{in} x_n = b_i \quad \text{for } i \in E \\ & x_j \geq 0 \quad \text{for } j \in N \end{aligned}$$

Dual LP:

$$\begin{aligned} \min \quad & b_1 y_1 + \cdots + b_m y_m \\ & a_{1j} y_1 + \cdots + a_{mj} y_m \geq c_j \quad \text{for } j \in N \\ & a_{1j} y_1 + \cdots + a_{mj} y_m = c_j \quad \text{for } j \notin N \\ & y_i \geq 0 \quad \text{for } i \in I \end{aligned}$$

Matrix-Vector Form and Its Dual



SHANGHAI JIAO TONG
UNIVERSITY

$$\max x_1 + 6x_2$$

$$x_1 \leq 200$$

$$x_2 \leq 300$$

$$x_1 + x_2 \leq 400$$

$$x_1, x_2 \geq 0$$

$$\min 200y_1 + 300y_2 + 400y_3$$

$$y_1 + y_3 \geq 1$$

$$y_2 + y_3 \geq 6$$

$$y_1, y_2, y_3 \geq 0$$



Theorem (Duality)

*If a linear program has a **bounded optimum**, then so does its **dual**, and the two optimum values coincide.*

若一个线性规划问题有界，那么他的对偶问题也有界，并且两者的最优解对应的最优数值应该相同。

Complementary Slackness



The number of variables in the dual is equal to that of constraints in the primal and the number of constraints in the dual is equal to that of variables in the primal.

Complementary Slackness



The number of variables in the dual is equal to that of constraints in the primal and the number of constraints in the dual is equal to that of variables in the primal.

An inequality constraint has slack if the slack variable is positive.

Complementary Slackness

The number of variables in the dual is equal to that of constraints in the primal and the number of constraints in the dual is equal to that of variables in the primal.

An inequality constraint has slack if the slack variable is positive.

The **complementary slackness** refers to a relationship between the slackness in a primal constraint and the associated dual variable.



$$\begin{aligned}\max & x_1 + 6x_2 \\ & x_1 \leq 200 \\ & x_2 \leq 300 \\ & x_1 + x_2 \leq 400 \\ & x_1, x_2 \geq 0\end{aligned}$$

$$x_1 = 100, x_2 = 300$$

$$\begin{aligned}\min & 200y_1 + 300y_2 + 400y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + y_3 \geq 6 \\ & y_1, y_2, y_3 \geq 0\end{aligned}$$

$$y_1 = 0, y_2 = 5, y_3 = 1$$

Theorem

Assume LP problem (P) has a solution x^* and its dual problem (D) has a solution y^* .

- 1 If $x_j^* > 0$, then the j -th constraint in (D) is binding.
- 2 If the j -th constraint in (D) is not binding, then $x_j^* = 0$.
- 3 If $y_i^* > 0$, then the i -th constraint in (P) is binding.
- 4 If the i -th constraint in (P) is not binding, then $y_i^* = 0$.

e. g. : $6x_1 + 12x_2 + 6$

$x_1, x_2 \geq 0$

$x_1 + x_2 \leq 2$

$x_1 + 2x_2 \leq 3$

$x_1 - 3x_2 \leq 5$

的对偶？

由于对偶实质上就是在找极值

等价的等价问题，所以只需要先将+6

移除，之后在对偶之后的问题上面在加6就可以了。