



Math for the people, by the people.

## idempotency

|                  |                     |
|------------------|---------------------|
| Canonical name   | Idempotency         |
| Date of creation | 2013-03-22 12:27:31 |
| Last modified on | 2013-03-22 12:27:31 |
| Owner            | yark (2760)         |
| Last modified by | yark (2760)         |
| Numerical id     | 21                  |
| Author           | yark (2760)         |
| Entry type       | Definition          |
| Classification   | msc 20N02           |
| Related topic    | BooleanRing         |
| Related topic    | PeriodOfMapping     |
| Related topic    | Idempotent2         |
| Defines          | idempotent          |

If  $(S, *)$  is a magma, then an element  $x \in S$  is said to be *idempotent* if  $x * x = x$ . For example, every identity element is idempotent, and in a group this is the only idempotent element. An idempotent element is often just called an idempotent.

If every element of the magma  $(S, *)$  is idempotent, then the binary operation  $*$  (or the magma itself) is said to be idempotent. For example, the  $\wedge$  and  $\vee$  operations in a <http://planetmath.org/Lattice> are idempotent, because  $x \wedge x = x$  and  $x \vee x = x$  for all  $x$  in the lattice.

A function  $f: D \rightarrow D$  is said to be idempotent if  $f \circ f = f$ . (This is just a special case of the first definition above, the magma in question being  $(D^D, \circ)$ , the monoid of all functions from  $D$  to  $D$  with the operation of function composition.) In other words,  $f$  is idempotent if and only if repeated application of  $f$  has the same effect as a single application:  $f(f(x)) = f(x)$  for all  $x \in D$ . An idempotent linear transformation from a vector space to itself is called a projection.