



Math for the people, by the people.

Young's projection operators

Canonical name	YoungsProjectionOperators
Date of creation	2013-03-22 16:48:25
Last modified on	2013-03-22 16:48:25
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	16
Author	rspuzio (6075)
Entry type	Definition
Classification	msc 20C30
Classification	msc 11P99
Classification	msc 05A17

Associated to a Young tableau with n boxes, we have two elements of the group ring of the permutation group on n symbols. To construct the operators, we first construct the antisymmetrizing operators corresponding to the columns and the symmetrizing operators corresponding to the rows. Then one operator corresponding to the tableau consists of the product of the symmetrizing operators corresponding to the rows multiplied by the product of the antisymmetrizing operators corresponding to the columns and the other consists of the product of the antisymmetrizing operators corresponding to the columns multiplied by the symmetrizing operators corresponding to the rows.

How this works may be illustrated with a simple example. Consider the tableau

$$\begin{array}{cc} 1 & 2 \\ 3 & 4 \end{array}$$

Corresponding to the first row, we have the symmetrization operator

$$1 + (12).$$

Corresponding to the second row, we have the symmetrization operator

$$1 + (34)$$

Multiplying these two symmetrization operators (the order does not matter because they involve permutations of different elements) produces

$$1 + (12) + (34) + (12)(34)$$

Corresponding to the first column, we have the antisymmetrization operator

$$1 - (13).$$

Corresponding to the second column, we have the antisymmetrization operator

$$1 - (24).$$

Multiplying these two antisymmetrization operators (the order does not matter because they involve permutations of different elements) produces

$$1 - (13) - (24) + (13)(24).$$

To obtain one Young projector, we multiply the product of the symmetrization operators by the product of the antisymmetrization operators.

$$(1 + (12) + (34) + (12)(34))(1 - (13) - (24) + (13)(24)) =$$

$$\begin{aligned} 1 &+ (12) + (34) + (12)(34) - (13) - (123) - (134) - (1234) - \\ (24) &- (142) - (243) - (1432) + (13)(24) + (1423) + (1324) + (14)(23) \end{aligned}$$

To obtain the other projector, we multiply in the other order.

$$(1 - (13) - (24) + (13)(24))(1 + (12) + (34) + (12)(34)) =$$

$$\begin{aligned} 1 &+ (12) + (34) + (12)(34) - (13) - (132) - (143) - (1432) - \\ (24) &- (124) - (234) - (1234) + (13)(24) + (1324) + (1423) + (14)(23) \end{aligned}$$