



planetmath.org

Math for the people, by the people.

isotope of a groupoid

Canonical name	IsotopeOfAGroupoid
Date of creation	2013-03-22 18:35:54
Last modified on	2013-03-22 18:35:54
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	8
Author	CWoo (3771)
Entry type	Definition
Classification	msc 20N02
Classification	msc 20N05
Synonym	isotopism
Synonym	homotopism
Defines	isotopy
Defines	isotope
Defines	homotopy
Defines	homotope
Defines	isotopic
Defines	homotopic
Defines	principal isotopy
Defines	principal isotope

Let G, H be <http://planetmath.org/Groupoidgroupoids>. An *isotopy* ϕ from G to H is an ordered triple: $\phi = (f, g, h)$, of bijections from G to H , such that

$$f(a)g(b) = h(ab) \quad \text{for all } a, b \in G.$$

H is called an *isotope* of G (or H is *isotopic* to G) if there is an isotopy $\phi : G \rightarrow H$.

Some easy examples of isotopies:

1. If $f : G \rightarrow H$ is an isomorphism, $(f, f, f) : G \rightarrow H$ is an isotopy. By abuse of language, we write $f = (f, f, f)$. In particular, $(1_G, 1_G, 1_G) : G \rightarrow G$ is an isotopy.
2. If $\phi = (f, g, h) : G \rightarrow H$ is an isotopy, then so is

$$\phi^{-1} := (f^{-1}, g^{-1}, h^{-1}) : H \rightarrow G,$$

for if $f^{-1}(a) = c$ and $g^{-1}(b) = d$, then $ab = f(c)g(d) = h(cd)$, so that $f^{-1}(a)g^{-1}(b) = cd = h^{-1}(ab)$

3. If $\phi = (f, g, h) : G \rightarrow H$ and $\gamma = (r, s, t) : H \rightarrow K$ are isotopies, then so is

$$\gamma \circ \phi := (r \circ f, s \circ g, t \circ h) : G \rightarrow K,$$

for $(r \circ f)(a)(s \circ g)(b) = r(f(a))s(g(b)) = t(f(a)g(b)) = t(h(ab)) = (t \circ h)(ab)$.

From the examples above, it is easy to see that “groupoids being isotopic” on the class of groupoids is an equivalence relation, and that an isomorphism class is contained in an isotopic class. In fact, the containment is strict. For an example of non-isomorphic isotopic groupoids, see the reference below. However, if G is a groupoid with unity and G is isotopic to a semigroup S , then it is isomorphic to S . Other conditions making isotopic groupoids isomorphic can be found in the reference below.

An isotopy of the form $(f, g, 1_H) : G \rightarrow H$ is called a *principal isotopy*, where 1_H is the identity function on H . H is called a *principal isotope* of G . If H is isotopic to G , then H is isomorphic to a principal isotope K of G .

Proof. Suppose $(f, g, h) : G \rightarrow H$ is an isotopy. To construct K , start with elements of G , which will form the underlying set of K . The binary operation on K is defined by

$$a \cdot b := (f^{-1} \circ h)(a)(g^{-1} \circ h)(b).$$

Then \cdot is well-defined, since f, g are bijective, for all pairs of elements of G . Hence K is a groupoid. Furthermore, $(f^{-1} \circ h, g^{-1} \circ h, 1_K) : G \rightarrow K$ is an isotopy by definition, so that K is a principal isotope of G . Finally, $h(a \cdot b) = h(f^{-1}(h(a))g^{-1}(h(b))) = f(f^{-1}(h(a)))g(g^{-1}(h(b))) = h(a)h(b)$, showing that $h : K \rightarrow H$ is a bijective homomorphism, and hence an isomorphism. \square

Remark. In the literature, the definition of an isotope is sometimes limited to quasigroups. However, this is not necessary, as the follow proposition suggests:

Proposition 1. *Any isotope of a quasigroup is a quasigroup.*

Proof. Suppose $(f, g, h) : G \rightarrow H$ is an isotopy, and G a quasigroup. Pick $x, z \in H$. Let $a, c \in G$ be such that $f(a) = x$ and $h(c) = z$. Let $b \in G$ be such that $ab = c$. Set $y = g(b) \in H$. Then $xy = f(a)g(b) = h(ab) = h(c) = z$. Similarly, there is $t \in H$ such that $tx = z$. Hence H is a quasigroup. \square

On the other hand, an isotope of a loop may not be a loop. Nevertheless, we sometimes say that an isotope of a loop L as a loop isotopic to L .

References

- [1] R. H. Bruck: *A Survey of Binary Systems*. Springer-Verlag. New York (1966).