



planetmath.org

Math for the people, by the people.

adjoining an identity to a semigroup

Canonical name	AdjoiningAnIdentityToASemigroup
Date of creation	2013-03-22 13:01:19
Last modified on	2013-03-22 13:01:19
Owner	mclase (549)
Last modified by	mclase (549)
Numerical id	5
Author	mclase (549)
Entry type	Definition
Classification	msc 20M99
Related topic	Semigroup
Related topic	Monoid

It is possible to formally adjoin an identity element to any semigroup to make it into a monoid.

Suppose  $S$  is a semigroup without an identity, and consider the set  $S \cup \{1\}$  where  $1$  is a symbol not in  $S$ . Extend the semigroup operation from  $S$  to  $S \cup \{1\}$  by additionally defining:

$$s \cdot 1 = s = 1 \cdot s, \text{ for all } s \in S \cup 1$$

It is easy to verify that this defines a semigroup (associativity is the only thing that needs to be checked).

As a matter of notation, it is customary to write  $S^1$  for the semigroup  $S$  with an identity adjoined in this manner, if  $S$  does not already have one, and to agree that  $S^1 = S$ , if  $S$  does already have an identity.

Despite the simplicity of this construction, however, it rarely allows one to simplify a problem by considering monoids instead of semigroups. As soon as one starts to look at the structure of the semigroup, it is almost invariably the case that one needs to consider subsemigroups and ideals of the semigroup which do not contain the identity.