



planetmath.org

Math for the people, by the people.

mixed group

Canonical name	MixedGroup
Date of creation	2013-03-22 18:42:32
Last modified on	2013-03-22 18:42:32
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	7
Author	CWoo (3771)
Entry type	Definition
Classification	msc 20N99
Defines	kernel

A *mixed group* is a partial groupoid G such that G contains a non-empty subset K , called the *kernel* of G , with the following conditions:

1. if $a, b \in G$, then ab is defined iff $a \in K$,
2. if $a, b \in K$ and $c \in G$, then $(ab)c = a(bc)$,
3. if $a \in K$, then $K \subseteq aK \cap Ka$,
4. if $a \in K$ and $b \in G$ such that $ab = b$, then $ac = c$ for all $c \in G$.

Mixed groups are generalizations of groups, as the following proposition illustrates:

Proposition 1. *If $K = G$, then G is a group.*

Proof. G is a groupoid by condition 1, and a semigroup by condition 2.

Now, by condition 3, given $a \in G$, there is $b \in G$ such that $ba = a$, so that $bc = c$ for all $c \in G$ by condition 4. In other words, b is a left identity of G . Again, by condition 3, for every $a \in G$, there is a $d \in G$ such that $b = da$. So $ad = a(bd) = a(da)d = (ad)^2$, so, by condition 4, $adx = x$ for all $x \in G$. In particular, set $x = a$, we get $a = (ad)a = a(da) = ab$. Hence, b is a two-sided identity, and G is a monoid.

Finally, by condition 3, for every $a \in G$, there are $c, d \in G$, such that $b = ac = da$. So, $c = bc = (da)c = d(ac) = db = d$, showing that a has a two-sided inverse. This means that G is a group. \square

For a non-trivial example of a mixed group, let G be a group and H a subgroup of G . Define a new multiplication \cdot on G as follows: $a \cdot b$ is defined iff $a \in H$, and if $a \cdot b$ is defined, it is defined as ab , the group multiplication of a and b . Then (G, \cdot) is a mixed group. Clearly, associativity of \cdot is automatically satisfied. Next, pick any $a \in H$, then, for any $b \in H$, $a^{-1} \cdot b$ and $b \cdot a^{-1}$ are both elements of H , so that $b \in a \cdot H \cap H \cdot a$, and condition 3 is also satisfied. Finally, if $a \in H$ and $b \in G$ such that $a \cdot b = b$, then a is the multiplicative identity of G , clearly $a \cdot c = c$ for all $c \in G$.

References

- [1] R. H. Bruck, *A Survey of Binary Systems*, Springer-Verlag, 1966

- [2] R. Baer, *Zur Einordnung der Theorie der Mischgruppen in die Gruppentheorie*, S.-B. Heidelberg. Akad. Wiss., Math.-naturwiss. Kl. 1928, 4, 13 pp
- [3] R. Baer, *Über die Zerlegungen einer Mischgruppe nach einer Untermischgruppe*, S.-B. Heidelberg. Akad. Wiss., Math.-naturwiss. Kl. 1928, 5, 13 pp
- [4] A. Loewy, *Über abstrakt definierte Transmutationssysteme oder Mischgruppen*, J. reine angew. Math. 157, pp 239-254, 1927